Cognitive behaviour therapy for excessive worry over health in medical patients

Submission date	Recruitment status	[X] Prospectively registered		
17/06/2008	No longer recruiting	[X] Protocol		
Registration date	Overall study status	Statistical analysis plan		
20/06/2008	Completed	[X] Results		
Last Edited	Condition category	Individual participant data		
17/05/2016	Mental and Behavioural Disorders			

Plain English summary of protocol

Background and study aims

Hypochondriasis, or health anxiety, is a significant problem in all parts of clinical practice, but has been largely ignored by services until recently. Our studies, and those of others, suggest that between 1 in 4 and 1 in 10 of all those attending medical clinics have significant health anxiety and this accounts for both considerable suffering and unnecessary use of resources. One of the characteristics of those with severe health anxiety is that it persists and leads to frequent medical consultations in both primary and secondary care. This represents a significant drain on health service resources at a time of considerable cost pressures, and, in addition, the symptoms of health anxiety are highly troubling and disturbing, are associated with much time off work, and adversely affect social functioning. In a study of treatment in a genitourinary medicine clinic, we found that a short course of cognitive behaviour therapy (CBT) was highly effective at reducing health anxiety, and that these benefits are maintained over a period of one year. We also found that the number of consultations in both general practice and the clinics fell in those in the active treatment group over one year but the savings made were small as the costs of investigations are relatively low. We are now planning a larger study of CBT to treat health anxiety in cardiology, respiratory medicine, gastroenterology and endocrinology clinics, where such anxiety is more common than in genuitourinary medicine.

Who can participate?

Patients aged between 16 and 75 attending four medical specialty clinics (in cardiology, respiratory medicine, gastroenterology and endocrinology) who have significant health anxiety

What does the study involve?

Participants are randomly allocated to one of two groups. One group is treated with CBT adapted for health anxiety in the form of between 5 and 10 one-hour sessions, which will address abnormal worries about health and ways of overcoming them. The other group is treated with a single 45-minute explanatory interview describing the nature of health anxiety and how it tends to be perpetuated. We measure patients' functioning, anxiety and quality of life, as well as health service and employment costs, at the start of the study and at 6 monthly intervals for two years, to see whether the costs of treatment are offset by savings on attendance and investigations over 2 years.

What are the possible benefits and risks of participating? Not provided at time of registration

Where is the study run from? Imperial College London (UK)

When is the study starting and how long is it expected to run for? October 2008 to September 2012

Who is funding the study? Health Technology Assessment Programme (UK)

Who is the main contact? Prof. Peter Tyrer p.tyrer@imperial.ac.uk

Contact information

Type(s)

Scientific

Contact name

Prof Peter Tyrer

Contact details

Department of Psychological Medicine St Dunstan's Road London United Kingdom W6 8RP +44 (0)207 386 1237 p.tyrer@imperial.ac.uk

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers HTA 07/01/26

Study information

Scientific Title

Cognitive behaviour therapy for Health Anxiety in Medical Patients

Acronym

CHAMP

Study objectives

Cognitive behaviour therapy adapted for health anxiety is a cost-effective way of treating patients with significant health anxiety in medical clinics.

More details can be found at: http://www.nets.nihr.ac.uk/projects/hta/070126 Protocol can be found at: http://www.nets.nihr.ac.uk/__data/assets/pdf_file/0017/51731/PRO-07-01-26.pdf

Ethics approval required

Old ethics approval format

Ethics approval(s)

Nottingham Research Ethics Committee 1, 13/06/2008, ref: 08/H0403/56

Study design

Randomised controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Not specified

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Hypochondriasis/mental health

Interventions

Arm 1: Cognitive behaviour therapy adapted for health anxiety (Warwick/Salkovskis model). Treatment will be given for between 5 and 10 one hour sessions for each patient, and will address abnormal worries about health and ways of overcoming them.

Arm 2: Single explanatory interview of the nature of health anxiety. This will be a single interview of 45 minutes describing the nature of health anxiety and how it tends to be perpetuated.

Duration of interventions: up to 6 months Duration of follow-up: 2 years

Intervention Type

Other

Phase

Not Specified

Primary outcome measure

Reduction in health anxiety scores 1 year after randomisation.

Secondary outcome measures

- 1. Reduction in health anxiety scores 2 years after randomisation
- 2. Presence or absence of diagnosis of hypochondriasis at 1 and 2 years after randomisation
- 3. Reduction in generalised anxiety and depression, assessed using the Hospital Anxiety and Depression Scale (HADS) at 1 and 2 years after randomisation
- 4. Improvement in social function and quality of life, assessed using the Social Functioning Questionnaire (SFQ) and Eurogol EQ-5D at 1 and 2 years after randomisation
- 5. Total costs of health care (primary and secondary) 1 and 2 years after randomisation
- 6. Change in presenteeism and absenteeism at work 1 and 2 years after randomisation

Overall study start date

01/10/2008

Completion date

30/09/2012

Eligibility

Key inclusion criteria

Patients attending four medical specialty clinics (in cardiology, respiratory medicine, gastroenterology and endocrinology) who:

- 1. Have significant health anxiety (score of 20 or more on the Health Anxiety Inventory)
- 2. Are aged between 16 and 75 (both males and females)
- 3. Are living independently
- 4. Are permanent residents in the area
- 5. Have sufficient understanding of English to read and complete the questionnaires
- 6. Give written consent for the interviews, including audiotaping of half the sessions
- 7. Give written consent for access to their medical records

Participant type(s)

Patient

Age group

Adult

Sex

Both

Target number of participants

465

Key exclusion criteria

Patients who are:

- 1. Considered too ill in medical terms by their consultants to be considered for the study
- 2. In the process of being investigated for significant pathology and for whom cognitive behaviour therapy might confuse or cause distress
- 3. Those who have significant cognitive impairment

Date of first enrolment

01/10/2008

Date of final enrolment

30/09/2012

Locations

Countries of recruitment

England

United Kingdom

Study participating centre Imperial College London

London United Kingdom W6 8RP

Sponsor information

Organisation

Imperial College London (UK)

Sponsor details

Charing Cross Campus
St Dunstan's Road
London
England
United Kingdom
W6 8RP
+44 (0)207 386 1233
gary.roper@imperial.ac.uk

Sponsor type

University/education

Website

http://www3.imperial.ac.uk

ROR

https://ror.org/041kmwe10

Funder(s)

Funder type

Government

Funder Name

Health Technology Assessment Programme

Alternative Name(s)

NIHR Health Technology Assessment Programme, HTA

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Protocol article	protocol	14/06/2011		Yes	No
Results article	results	18/01/2014		Yes	No