

# Effect of hyaluronic acid on the oral mucosa healing after diode laser biopsies

<b>Submission date</b> 29/08/2024	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 06/09/2024	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 06/09/2024	<b>Condition category</b> Oral Health	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Oral mucosa can be affected by lesions which interfere with many functions such as eating, speaking, and swallowing. Oral fibroma is one of these lesions that appears commonly in the oral cavity. It is the most common reactive lesion in the oral cavity. Fibroma results from a chronic repair process that includes granulation tissue and scar formation resulting in a fibrous submucosal mass. This lesion doesn't have a risk of malignancy transformation. The most common sites in the oral cavity are the tongue, buccal mucosa, and lower labial mucosa. An excisional biopsy is the treatment choice for irritation fibroma, either by surgical scalpel, electric scalpel, cryosurgery, or diode laser. Diode laser based on many studies is a very sufficient way of taking excisional biopsies, however, it has a thermal effect on the oral mucosa that affects mucosal healing, which makes it take longer to heal compared to the surgical conventional way. Accordingly, hyaluronic acid (HA) is now used to accelerate the mucosal healing process after excision using lasers in general.

### Who can participate?

Patients aged between 20 and 40 years old who have oral fibroma and systemic diseases free

### What does this study involve?

Participants are randomly allocated into three groups: Group A (HA gel + diode laser), Group B (HA mouthwash + diode laser), and Group C (control group, diode laser only). HA use instructions are given to the first two groups, and oral hygiene instructions are given to all groups.

### What are the possible benefits and risks of participating?

Benefits may include healing acceleration due to the HA application after taking the excisional biopsy of the oral fibroma using a diode laser.

Possible risks might be experiencing some pain and discomfort after taking the biopsy, and a slow healing process due to the thermal effect of the diode laser and the ineffectiveness of the HA.

### Where is the study run from?

Damascus University, Syria

When is the study starting and how long is it expected to run for?  
June 2023 to October 2025

Who is funding the study?  
Damascus University, Syria

Who is the main contact?  
Dr. Amr Alyafi, amr.alyafi@hotmail.com, amr.alyafi97@damascusuniversity.edu.sy

## Contact information

### Type(s)

Public, Scientific, Principal investigator

### Contact name

Dr Amr Alyafi

### ORCID ID

<https://orcid.org/0009-0001-8761-8019>

### Contact details

Telyani  
Damascus  
Syria  
4671  
+963947880112  
amr.alyafi97@damascusuniversity.edu.sy

## Additional identifiers

### Clinical Trials Information System (CTIS)

Nil known

### ClinicalTrials.gov (NCT)

Nil known

### Protocol serial number

5289

## Study information

### Scientific Title

Studying the effect of hyaluronic acid on mucosal healing after excisional oral biopsies with 810 nm diode laser

### Study objectives

h0: There are no statistical differences between the three studied groups. (HA gel group, HA mouthwash group, and only diode laser group)

h1: There are statistically significant differences between the three studied groups.

## **Ethics approval required**

Ethics approval required

## **Ethics approval(s)**

approved 07/08/2023, Scientific Ethics Committee at Damascus University (Baramkeh, Damascus, 4671, Syria; +9631133923223; ap.srd@damascusuniversity.edu.sy), ref: 3437

## **Study design**

Randomized controlled trial

## **Primary study design**

Interventional

## **Study type(s)**

Other, Treatment, Efficacy

## **Health condition(s) or problem(s) studied**

Mucosal healing after oral excisional biopsies

## **Interventions**

This study is designed to evaluate the effect of hyaluronic acid (HA; Mouthwash - Gel) on mucosal healing after oral excisional biopsies with an 810 nm diode laser.

A simple method of randomisation is used that involves writing the group name (A, B, C) on a piece of paper inside three envelopes, and letting the patient choose the envelope randomly.

A brief methodology for each treatment arm:

-Group A (HA gel group + diode laser):

After taking the excisional biopsy with a diode laser (with parameters of 810 nm, 4W) which takes approximately less than 5 minutes, the patients will be given topical hyaluronic acid gel 0.2% to apply on the wound 3 times daily for 10 days. After each appliance, the patient should not eat or drink for an hour.

-Group B (HA mouthwash group + diode laser):

After taking the excisional biopsy with a diode laser (with parameters of 810 nm, 4W) which takes approximately less than 5 minutes, the patients will be given hyaluronic acid mouthwash 0.025% to rinse with 3 times daily out of brushing times for 10 days, by keeping it for a minute in the mouth and then spitting it out. After each rinsing the patient should not eat or drink for an hour.

-Group C (diode laser only):

After taking the excisional biopsy with a diode laser (with parameters of 810 nm, 4W) which takes approximately less than 5 minutes, the wound will be left to heal spontaneously without allying any drug.

Percentage healing index (PHI: is an index that helps determine the healing process by taking a picture of the excision wound after the procedure immediately and after every follow-up visit, and then using the Adobe Photoshop application on PC to measure the excision wound area in square millimeters.

## **Intervention Type**

Procedure/Surgery

**Primary outcome(s)**

Excision wound area measured using the Percentage Healing Index (PHI) after the excision immediately (T0), at day 4 (T1), week 1 (T2), week 2 (T3), and 1 month (T4)

**Key secondary outcome(s)**

Pain is measured using a Visual Analogue Scale (VAS) at day 4, week 1, week 2, and 1 month

**Completion date**

24/10/2025

## Eligibility

**Key inclusion criteria**

1. Patients that have oral fibromas on the non-keratinized oral mucosa
2. Excisional biopsy wounds with a diameter of less than 10mm

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Lower age limit**

20 years

**Upper age limit**

40 years

**Sex**

All

**Key exclusion criteria**

1. People who have systemic diseases
2. Smokers
3. Alcoholics

**Date of first enrolment**

01/07/2023

**Date of final enrolment**

01/07/2025

## Locations

Countries of recruitment

Syria

**Study participating centre**

Oral Medicine Department, Faculty of Dental Medicine, Damascus University

Mezzeh highway

Damascus

Syria

4671

## Sponsor information

**Organisation**

Damascus University

**ROR**

<https://ror.org/03m098d13>

## Funder(s)

**Funder type**

University/education

**Funder Name**

Damascus University

**Alternative Name(s)**

University of Damascus, , DU

**Funding Body Type**

Government organisation

**Funding Body Subtype**

Universities (academic only)

**Location**

Syria

## Results and Publications

Individual participant data (IPD) sharing plan

The data sets will be generated and analyzed during the current study and will be available upon request from Amr Alyafi, amr.alayafi@hotmail.com, amr.alayafi97@damascusuniversity.edu.sy

**IPD sharing plan summary**

Available on request

**Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Participant information sheet</a>			02/09/2024	No	Yes
<a href="#">Participant information sheet</a>			02/09/2024	No	Yes
<a href="#">Participant information sheet</a>	Participant information sheet	11/11/2025	11/11/2025	No	Yes