Autologous intra-carotid arterial transfusion of peripheral blood stem cells improves brain ischemia and reperfusion in patients with chronic stroke

Submission date 11/04/2016	Recruitment status No longer recruiting	 Prospectively registered Protocol
Registration date 28/06/2016	Overall study status Ongoing	 Statistical analysis plan Results
Last Edited 06/06/2024	Condition category Circulatory System	 Individual participant data Record updated in last year

Plain English summary of protocol

Background and study aims

Stroke is a leading cause of death and disability worldwide, and is the third highest cause of death in Taiwan. About 85% of strokes are ischemic strokes. Ischemic strokes happen when the arteries that supply the brain with oxygen (the carotid arteries) become narrowed (stenosis) or blocked (occluded) by a sticky substance called plaque that builds up on the artery walls (atherosclerosis), causing reduced blood flow (ischemia) to the brain. Atherosclerosis in the carotid arteries is one of the biggest causes of recurrent stroke, and so reducing carotid stenosis is an important part of stroke prevention. There is growing evidence that stem cell therapy using endothelial progenitor cells (EPCs) (stem cells from bone marrow) can help to regenerate the lining of blood vessels. Previous studies have shown that increasing levels of EPCs circulating in the blood is related to improvement in the recovery of those with ischemic stroke. Currently, there is little research looking at the safety and effectiveness of using EPC therapy for treating ischemic stroke. The aim of this study is to find out whether ECP therapy is an effective treatment for patients with brain ischemia following an ischemic stroke.

Who can participate?

Adults aged between 45 and 80 who had an ischaemic stroke at least six months ago.

What does the study involve?

All participants receive eight injections of granulocyte-colony stimulating factor (a protein which stimulates the bone marrow to produce stem cells and release them into the bloodstream) every 12 hours for four days. On the fifth day, patients have a tube placed into the right femoral vein (main vein in the thigh) so that circulating EPCs can be collected. These EPCs are then injected back into the patients into the internal carotid artery (in the neck). Patients are then followed up for five years through a combination of interviews and medical record reviews, in order to find out how effective the treatment has been.

What are the possible benefits and risks of participating? Participants may benefit from improvement to their disability. There are risks of side effects from the EPC therapy, including deterioration of brain function, recurrent stroke, heart problems, blockage of arteries, bleeding, anemia, deterioration of kidney function, gastrointestinal (gut) complications, electrolyte (minerals in the body) imbalance, sepsis (blood poisoning) and cancer.

Where is the study run from? Kaohsiung Chang Gung Memorial Hospital (Taiwan)

When is the study starting and how long is it expected to run for? January 2014 to July 2026

Who is funding the study? National Science Council (Taiwan)

Who is the main contact? Dr Hon-Kan Yip han.gung@msa.hinet.net

Contact information

Type(s) Public

Contact name Dr Hon-Kan Yip

Contact details

University College of Medicine 123, Ta Pei Road Niao Sung District Kaohsiung City Taiwan 83302 +886 7 7317123 ext. 8300 han.gung@msa.hinet.net

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers N/A

Study information

Scientific Title

Autologous intra-carotid arterial transfusion of peripheral blood stem cells (PBSC) improves brain ischemia and reperfusion – A safety and tolerability study

Study objectives

Peripheral Blood Stem Cells transplantation for the patients with brain ischemia may be a more effective therapy than traditional drug therapy.

Ethics approval required Old ethics approval format

Ethics approval(s) Chang Gung Memorial Hospital, 24/11/2015, ref: 104-6798C

Study design Prospective single-centre non-randomised study

Primary study design Interventional

Secondary study design Non randomised study

Study setting(s) Hospital

Study type(s) Treatment

Participant information sheet

No specific participant information sheet available, please use the contact details below to request a further information.

Health condition(s) or problem(s) studied

Chronic ischemic stroke

Interventions

All participants receive eight granulocyte-colony stimulating factor (G-CSF) injections of 5 µg/kg sc q12h over a course of four days after admission and endothelial progenitor cell (EPC) collection at day 5 for 3 hours from double lumen sheath inserted from right femoral vein using a machine (COBE Spectra 6.1). The collected EPC will be immediately transfused back to patients themselves with a dose of 3x10(7) EPC once through internal carotid artery of infarct side using a 6F JR catheter through left radial arterial sheath. The participants will be followed up at Neurology and Cardiovascular outpatients clinic for safety and efficacy monitoring for 5 years.

Intervention Type

Procedure/Surgery

Primary outcome measure

Safety and tolerability of intracarotid artery autologous endothelial progenitor cells injection is measured through patient interviews and reviewing medical records throughout the five years of the study

Secondary outcome measures

 Recurrent stroke or death rate is measured by reviewing medical records at 90 days
 Disability is measured using the modified Rankin Scale score, Barthel Index WAIS-III and CASI National Institutes of Health Stroke Scale (NIHSS) at 24 hours, 72 hours, 30 days and 90 days
 Brain reperfusion status is measured using Magnetic resonance imaging (Arterial Spin Labeling (ASL) and Dynamic Contrast Enhancement) at 24-72 hours post-therapy and 90 days
 Brain perfusion defects are evaluated using radionuclear medicine image (brain Tc-99m scan) 24-72 hours post therapy and 90 days

5. Adverse event rate is measured through patient interviews and reviewing medical records at 5 years

Overall study start date

01/01/2014

Completion date

31/07/2026

Eligibility

Key inclusion criteria

 Aged between 45 and 80 years
 Chronic ischemic stroke (onset more than 6 months ago) over middle cerebral artery territory (NIHSS score 9-16)
 Not suitable for surgery or carotid artery stenting

Participant type(s)

Patient

Age group

Adult

Sex Both

Target number of participants 10 patients

Total final enrolment 28

Key exclusion criteria

- 1. Age less than 45 years old or more than 80 years old
- 2. Patients received t-PA and anticoagulation
- 3. Non-middle cerebral artery territory stroke or hemorrhagic stroke
- 4. Pregnant women

- 5. Patients with adventitious agents (like HIV virus)
- 6. Myocardial infarction (MI) within 3 months
- 7. Severe aortic stenosis or mitral stenosis
- 8. Congestive heart failure, New York Heart Association functional class (NYHA Fc.)IV
- 9. Malignancy or other severe disease with life span less than one year
- 10. Chronic kidney disease with CCr<20ml/min and end stage renal disease
- 11. Join other clinical trials
- 12. Patients can not receive regular follow-up
- 13. Other brain disease (tumor, degenerative disease, infective disease)

Date of first enrolment 01/01/2016

Date of final enrolment 31/12/2019

Locations

Countries of recruitment Taiwan

Study participating centre Kaohsiung Chang Gung Memorial Hospital 123, Ta Pei Road Niao Sung District Kaohsiung Taiwan 83302

Sponsor information

Organisation National Science Council

Sponsor details

No. 106 HoPing E. Road Sec.2 Taipei Taiwan 10622 +886 2 2737 7992 misservice@most.gov.tw

Sponsor type

Government

Website

https://www.most.gov.tw/

ROR https://ror.org/02kv4zf79

Funder(s)

Funder type Government

Funder Name National Science Council

Alternative Name(s) National Science Council, Taiwan, National Science Council of Taiwan, NSC

Funding Body Type Government organisation

Funding Body Subtype National government

Location Taiwan

Results and Publications

Publication and dissemination plan Planned publication in a peer reviewed journal.

Intention to publish date 31/07/2028

Individual participant data (IPD) sharing plan

IPD sharing plan summary Data sharing statement to be made available at a later date