

Epirubicin plus tamoxifen versus tamoxifen alone in post-menopausal node positive primary breast cancer

Submission date 19/08/2002	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 19/08/2002	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 12/08/2008	Condition category Cancer	<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr - -

Contact details
UKCCCR Register Co-ordinator
MRC Clinical Trials Unit
222 Euston Road
London
United Kingdom
NW1 2DA

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers
ICCG/4/87

Study information

Scientific Title

Study objectives

Not provided at time of registration

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Health condition(s) or problem(s) studied

Breast cancer

Interventions

1. Regimen A: tamoxifen 20 mg daily for 4 years
2. Regimen B: tamoxifen 20 mg daily for 4 years plus chemotherapy, single agent epirubicin repeated every 4 weeks for six cycles

Intervention Type

Other

Phase

Not Specified

Primary outcome measure

Not provided at time of registration

Secondary outcome measures

Not provided at time of registration

Overall study start date

01/01/1996

Completion date

28/02/1998

Eligibility

Key inclusion criteria

1. Post-menopausal as determined by:
 - 1.1. Last menstrual cycle 12 months before surgery
 - 1.2. Patients any age with previous bilateral oophorectomy
 - 1.3. Patients aged greater than 50 years who have had hysterectomy without oophorectomy as long as the reason for surgery was not a malignancy
2. Aged less than 75 years
3. Histologically proven non metastatic T1-T3 with at least one involved ipsilateral axillary node
4. Adequate renal and haematological function
5. No prior history of malignant breast tumours
6. No bilateral malignancy or inflammatory breast cancer
7. No previous or concomitant malignancy, except squamous or basal cell carcinoma of the skin which has been effectively treated or carcinoma of the cervix in situ which has been treated operatively only
8. No non malignant systemic disease
9. No definite indication for chemotherapy

Participant type(s)

Patient

Age group

Adult

Sex

Female

Target number of participants

604

Key exclusion criteria

Not provided at time of registration

Date of first enrolment

01/01/1996

Date of final enrolment

28/02/1998

Locations

Countries of recruitment

Belgium

England

France

Greece

Netherlands

Spain

United Kingdom

Study participating centre
UKCCCR Register Co-ordinator
London
United Kingdom
NW1 2DA

Sponsor information

Organisation

Pharmacia Ltd & Upjohn (UK)

Sponsor details

Davy Avenue
Milton Keynes
United Kingdom
MK5 8PH
+44 (0)1908 661101
info@adresco.co.uk

Sponsor type

Industry

Website

<http://www.pharmacia.com>

ROR

<https://ror.org/04x4v8p40>

Funder(s)

Funder type

Industry

Funder Name

Pharmacia and Upjohn Ltd (UK)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date**Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/07/1999		Yes	No