

When drinking water to prevent fainting, does the temperature matter?

Submission date 04/11/2021	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 14/12/2021	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 18/07/2022	Condition category Other	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

The aim of this study is to determine whether drinking water can improve orthostatic tolerance in healthy control volunteers. Orthostatic tolerance refers to the ability to maintain adequate blood pressure when standing. In some people blood pressure can fall when standing, which can lead to dizzy spells or fainting episodes. Drinking water can boost blood pressure and make fainting episodes less likely to occur. However, it is not clear whether the temperature of the water has an impact on the blood pressure response. This study will test whether warm water and cold water have the same effect on blood pressure responses.

Who can participate?

Healthy English-speaking men and women aged 19-50 years

What does the study involve?

Participants will be asked to undergo a "tilt test". This test measures blood pressure control and susceptibility to fainting spells. Participants will undergo this test on three separate days. On each day they will be given a drink of water: either a 500 ml drink of room temperature water, a 500 ml drink of ice-cold water, or a 500 ml drink of warm water. The order in which they receive the water will be determined randomly.

Before the start of the test one of the study team will ask participants some questions about their medical history and their general health. There will also be questions about heart disease risk factors such as smoking, exercise levels, and alcohol consumption. They will measure the participants' height and weight. Participants will be asked to empty their bladder before testing. The researchers will take a urine measurement of the amount of salt in the urine – a marker of salt intake and hydration.

For the test procedure, participants will be asked to lie down on a bed while we attach monitoring equipment to their body. This will include:

1. An electrocardiogram (ECG). This is a monitor that will measure their heart beat (how fast and how regularly the heart is beating). The researchers will attach three adhesive electrodes (stickers) to the skin of the participant's chest and connect them to the ECG machine. If participants have a hairy chest, it may be necessary to shave three small areas of their chest in order to help the electrodes stick to their skin. An alcohol swab will be used to clean the skin before electrode placement.

2. A blood pressure monitor. A small Velcro cuff will be placed around the middle finger that pulses gently against the small arteries along the side of the finger and records their blood pressure with every heartbeat. This measurement is non-invasive and painless.
3. The researchers will measure breathing rate and the gases in the air the participants are breathing out with a small nasal cannula (a flexible tube). This small plastic tube will be placed under their nose, on their top lip, and will sample their breathing. Participants will be able to breathe and talk normally while wearing this device.
4. The researchers will measure the blood flow to the brain in an artery called the middle cerebral artery. They will do this using ultrasound (imaging device). They will place a little ultrasound gel on the temple and will position an ultrasound probe overlying the gel. The probe will be held in place with a headband. This means the investigators will not need to touch the participant to hold the probe. Participants can move their head freely when wearing the ultrasound probe. They will not be able to feel the ultrasound.
5. The researchers will also measure blood flow in the arm, in the brachial artery, also with ultrasound. The participant's arm will be placed on a support with a probe positioned over a little ultrasound gel near the elbow. Participants will be asked to keep their arm still during the test.
6. The researchers will place a strap over the participant's knees and a box over their legs that seals against their waist (a bit like a canoe skirt). The strap is to help them stand in a relaxed position without fidgeting their legs too much. The box is placed over their legs so that they can apply lower body negative pressure to their legs later on in the test without disturbing the monitoring.

Once the monitors are in place the researchers will make recordings from them for 15 minutes while they lie on your back and rest. Participants will then be given the water to drink. After a further 15 minutes of resting, they will tilt the table into an upright position (at 60 degrees). This is like standing but leaning backwards slightly. The researchers will make recordings from the monitors for a further 20 minutes. They will ask participants not to move their legs much during the test. After 20 minutes of standing, they will apply lower body negative pressure to the box over their legs. This will feel a little bit draughty, and may be a little noisy, but is not painful or unpleasant. The effect mimics prolonged standing. The researchers will apply the lower body negative pressure at three different levels for 10 minutes each (-20 mmHg, -40 mmHg and -60 mmHg).

The test will be stopped immediately if:

1. Participants complete the whole procedure (30 minutes lying down, 20 minutes standing, and 30 minutes of lower body negative pressure).
2. They experience symptoms of dizziness or lightheadedness and/or their blood pressure or heart rate begin to decrease.
3. They request the test to stop.

Participants will then be returned to the lying position. If they experience dizziness at the end of the test, lying down will quickly resolve this. The monitors will be removed and any residue from the ultrasound gel will be removed. It is common to feel a bit hot and sweaty at the end of the test. There are showers near to the lab, and the researchers can provide clean towels etc for participants to freshen up after the test if they wish.

What are the possible benefits and risks of participating:

There are no direct benefits to participants. It is hoped that the results of this study will ultimately aid in the treatment and management of fainting spells in patients who are prone to these episodes, and so improve quality of life for those affected by recurrent fainting episodes. Participants will receive up to \$75 to compensate you for your participation in the study (\$25 per test). Free parking is available on request.

What are the potential disadvantages to taking part?

The study will take place in a controlled laboratory environment and most participants do not find the assessments unpleasant. Every effort will be made to ensure their safety, privacy and comfort. The following are discomforts or risks that may be associated with the procedures.

1. During the tilt table test participants may experience some dizziness or lightheadedness associated with reduced blood pressure and/or heart rates. Rarely, participants have been known to faint briefly. Actual fainting is unusual and is always very short in duration with rapid return to consciousness.
2. These assessments will take time to perform and participants will be asked to keep still during the assessments. They may find that they become uncomfortable or bored during the course of these investigations. Every effort will be made to maintain their comfort throughout the study. Participants will be provided with pillows, blankets etc as appropriate to ensure their comfort. To compensate them for their time, participants will be paid \$25 for each of the three tests that they complete.
3. Preparing the skin for electrode placement may cause minor irritation or redness. It is possible that participants will experience an allergic reaction to the electrode gel or adhesive.

Where is the study run from?

The study is a collaborative effort between Simon Fraser University (Canada) and King's College London (UK). It will be run from the Department of Biomedical Physiology at Simon Fraser University.

When is the study starting and how long is it expected to run for?

December 2018 to October 2019

Who is funding the study?

National Sciences and Engineering Research Council of Canada (NSERC) (Canada)

Who is the main contact?

Prof. Victoria Claydon
victoria_claydon@sfu.ca

Contact information

Type(s)

Scientific

Contact name

Dr Iain Parsons

ORCID ID

<https://orcid.org/0000-0002-8577-8289>

Contact details

School of Life Sciences and Medicine
St Thomas' Hospital
King's College London
London
United Kingdom

SE17EH
+44 (0)20 7188 7188
iain.parsons@kcl.ac.uk

Type(s)

Public

Contact name

Prof Victoria Claydon

Contact details

Department of Biomedical Physiology
Simon Fraser University
8888 University Drive
Burnaby
Canada
V5A1S6
+1 (0)778 782 8560
victoria_claydon@sfu.ca

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

2019s0224

Study information

Scientific Title

The effect of water temperature on orthostatic tolerance

Study objectives

1. The cold-water condition would attenuate the deterioration of cardiovascular and cerebrovascular stability during orthostatic stress and increase OT (time to presyncope) in comparison to bolus water drinking at room temperature
2. The warm water condition would exacerbate the deterioration of cardiovascular and cerebrovascular stability during orthostatic stress and decrease OT in comparison to bolus water drinking at room temperature.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 16/07/2019, Simon Fraser University Research and Ethics Committee (Office of Research Ethics, Simon Fraser University, 8900 Nelson Way, Burnaby, BC, V5A 4W9, Canada; +1 (0)778 7823 447; dore@sfu.ca), ref: 2019s0224

Study design

Single-blind randomized cross over study

Primary study design

Interventional

Study type(s)

Other

Health condition(s) or problem(s) studied

Prevention of vasovagal syncope in patients with orthostatic intolerance

Interventions

Participants were randomised, in single-blind crossover fashion, to either a 500 ml drink of room temperature water, which was the control condition (20°C) (ROOM), a 500 ml drink of ice-cold water (0-3°C) (COLD), or a 500 ml drink of warm water (45°C) (WARM). Each intervention was applied on a different day, with a single day testing and endpoint (3 x one day).

Intervention Type

Behavioural

Primary outcome(s)

Orthostatic tolerance measured by time to presyncope following the head-up tilt test and subsequent progressive lower body negative pressure. Each intervention was applied on a different day, with a single day testing and endpoint (3 x one day).

Key secondary outcome(s)

Each intervention was applied on a different day, with a single day testing and endpoint (3 x one day):

1. Cardiovascular stability measured by Finometer, echocardiography and forearm ultrasound Doppler following the head-up tilt test and subsequent progressive lower body negative pressure
2. Cerebrovascular stability measured by middle cerebral artery ultrasound Doppler following the head-up tilt test and subsequent progressive lower body negative pressure

Completion date

01/10/2019

Eligibility

Key inclusion criteria

Healthy volunteers 19-50 years of age

Participant type(s)

Healthy volunteer

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Total final enrolment

17

Key exclusion criteria

1. <19 or >50 years old
2. Pregnant
3. Trying to conceive
4. Prior history of cardiovascular or neurological disease

Date of first enrolment

16/07/2019

Date of final enrolment

30/09/2019

Locations

Countries of recruitment

Canada

Study participating centre**Simon Fraser University**

Department of Biomedical Physiology

8888 University Drive

Burnaby

Canada

V5A 1S6

Sponsor information

Organisation

Simon Fraser University

ROR

<https://ror.org/0213rcc28>

Funder(s)

Funder type

Research council

Funder Name

Natural Sciences and Engineering Research Council of Canada (Discovery RGPIN/02982-2021 and Discovery Accelerator Supplement RGPAS/2021-00012-2021).

Alternative Name(s)

Conseil de Recherches en Sciences Naturelles et en Génie du Canada, The Natural Sciences and Engineering Research Council of Canada, nserc_crsng, NSERC, CRSNG

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

Canada

Funder Name

Ministry of Defence

Alternative Name(s)

Ministry of Defence, Government of India, Ministry of Defence, Govt. of India, Indian Ministry of Defence, Raksha Mantralay, MOD

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

India

Results and Publications

Individual participant data (IPD) sharing plan

An anonymised dataset is available from Prof. V Claydon (Victoria_claydon@sfu.ca) upon reasonable request. This will include the raw cardiovascular and cerebrovascular data. This will

be available for 2 years following the peer-reviewed publication of the results. The data will be available for the purposes of academic endeavour only for the purposes of meta-analysis following agreement by participants.

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		01/04/2022	18/07/2022	Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Protocol file	version 1	27/02/2019	05/11/2021	No	No