

AspCOVID-19 study: fungal infection of the airways in severe COVID-19 pneumonia

Submission date 06/04/2020	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 17/11/2020	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
Last Edited 16/04/2021	Condition category Infections and Infestations	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

COVID-19 is a condition caused by the coronavirus (called SARS-CoV-2) that was first identified in late 2019. This virus can infect the respiratory (breathing) system. Some people do not have symptoms but can carry the virus and pass it on to others. People who have developed the condition may develop a fever and/or a continuous cough among other symptoms. This can develop into pneumonia. Pneumonia is a chest infection where the small air pockets of the lungs, called alveoli, fill with liquid and make it more difficult to breathe.

In 2020, the virus has spread to many countries around the world and neither a vaccine against the virus or specific treatment for COVID-19 has yet been developed. As of March 2020, it is advised that people minimize travel and social contact, and regularly wash their hands to reduce the spread of the virus.

Groups who are at a higher risk from infection with the virus, and therefore of developing COVID-19, include people aged over 70 years, people who have long-term health conditions (such as asthma or diabetes), people who have a weakened immune system and people who are pregnant. People in these groups, and people who might come into contact with them, can reduce this risk by following the up-to-date advice to reduce the spread of the virus.

Bacterial or fungal superinfections are well-known complications of viral infections. Especially, the experiences from critically ill patients with severe influenza pneumonia showed that fungal infection (aspergillosis) is a relevant co-infection.

Based on recent observations it seems that the airway system of patients with COVID-19 pneumonia is exposed in the same way as in severe influenza pneumonia which suggests that infection complications could be similar. Invasive pulmonary aspergillosis may be a relevant co-infection in critically ill patients with severe COVID-19 pneumonia.

Aim of the underlying prospective AspCOVID-19 study is the screening for invasive pulmonary aspergillosis in critically ill patients with severe COVID-19 pneumonia.

Who can participate?

Critically ill patients with severe COVID-19 pneumonia, aged 18 years or above.

What does the study involve?

All mechanically ventilated patients with COVID-19 pneumonia will be screened for invasive pulmonary aspergillosis on day 1, 3, 7, 10, 14 and then every three days after ICU admission and intubation/mechanical ventilation.

What are the possible benefits and risks of participating?

Early recognition of co-infection, especially fungal infections, could be a benefit, because early treatment initiation minimize the risk of complications caused by infections.

As the provided procedures are standard examinations and daily routine in critically ill patients that were also done in critically ill without study participation there is no specific risk for study patients.

Where is the study run from?

Munich University Hospital (Germany)

When is the study starting and how long is it expected to run for?

March 2020 to July 2020

Who is funding the study?

Technical University of Munich (Germany)

Who is the main contact?

Dr Tobias Lahmer, tobiaslahmer@me.com

Contact information

Type(s)

Scientific

Contact name

Dr Tobias Lahmer

ORCID ID

<https://orcid.org/0000-0003-1008-5311>

Contact details

Medizinische Klinik und Poliklinik II

Klinikum rechts der Isar

Technische Universität München

Ismaninger Straße 22I

Munich

Germany

81675

+49 89 4140 9345

Tobias.Lahmer@mri.tum.de

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

Nil known

Study information

Scientific Title

Invasive pulmonary aspergillosis in severe COVID-19 pneumonia: AspCOVID-19 study

Acronym

AspCOVID-19 study

Study objectives

Bacterial and/or fungal superinfection are well-known complications of severe viral infections. Driven from experiences with severe influenza pneumonia we suggest that invasive pulmonary aspergillosis might be a relevant complication of severe COVID-19 pneumonia in critically ill patients.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 22/03/2020, Ethikkommission, Fakultät für Medizin, Technische Universität München (Ismaninger Straße 22, 81675 München, Germany; +49 89 4140-7737; ethikkommission@mri.tum.de), ref 149/20

Study design

Prospective observational study

Primary study design

Observational

Study type(s)

Screening

Health condition(s) or problem(s) studied

COVID-19 (SARS-CoV-2 infection) related pneumonia

Interventions

Screening of all mechanical ventilated patients with COVID-19 pneumonia for invasive pulmonary aspergillosis using the galactomannan from serum and broncho-alveolar lavage on day 1, 3, 7, 10, 14 and then every three days after ICU admission and intubation/mechanical ventilation. Combined with standard microbiological testing for bacteria and fungi and standard laboratory parameters (infection parameters: C-reactive protein, procalcitonin, Interleukin 6, leukocyte count, lymphocytes).

Intervention Type

Other

Primary outcome(s)

Pulmonary aspergillosis infection measured using the galactomannan from serum and broncho-alveolar lavage on day 1, 3, 7, 10, 14 and then every three days after ICU admission and intubation/mechanical ventilation

Key secondary outcome(s)

1. Duration of ICU stay measured using patient records
2. Mortality rate measured using patient records
3. Co-Infections measured using patient records
4. Ventilation time measured using patient records
5. Duration of hospital stay measured using patient records

Completion date

31/07/2020

Eligibility**Key inclusion criteria**

Critically ill patients with severe COVID-19 pneumonia

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

1. Younger than 18 years old
2. Pregnancy
3. Missed informed consent

Date of first enrolment

25/03/2020

Date of final enrolment

30/06/2020

Locations**Countries of recruitment**

Germany

Study participating centre

Klinikum rechts der Isar

Medizinische Klinik und Poliklinik II

Technische Universität München

Ismaninger Straße 22

Munich

Germany

81675

Sponsor information

Organisation

Klinikum der Universität München

ROR

<https://ror.org/02jet3w32>

Funder(s)

Funder type

University/education

Funder Name

Technische Universität München

Alternative Name(s)

Technical University of Munich, TUM

Funding Body Type

Government organisation

Funding Body Subtype

Universities (academic only)

Location

Germany

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		17/03/2021	16/04/2021	Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Preprint results	non-peer-reviewed results in preprint	22/07/2020	17/03/2021	No	No