

The regulate your sitting time (RESIT) study in type 2 diabetes

Submission date 23/07/2020	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 06/08/2020	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 17/02/2026	Condition category Nutritional, Metabolic, Endocrine	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Type 2 diabetes is a common condition that causes the level of sugar (glucose) in the blood to become too high. High levels of sitting increase the risk of heart disease and early death in people with type 2 diabetes, regardless of the amount of time they spend exercising. Interventions that help reduce sitting time in this group are therefore needed. Our team has developed an intervention that is designed to reduce the time people with type 2 diabetes spend sitting. The intervention is based on previous work that the research team have carried out over the past few years. The intervention involves patients choosing from a list of tools designed to help reduce their sitting time (so it is tailored to each patient) and patients will also receive support from a health coach and complete an online education session on the risks of sitting and how sitting could be reduced. The aim of this research is to see whether people like and engage with the intervention and how easy it is for us to take some measurements to see how well it worked.

Who can participate?

People with type 2 diabetes aged 18 - 85 years.

What does the study involve?

Participants will be randomly allocated to a control or intervention group for 6 months. The control participants will receive normal diabetes healthcare as usual. Intervention participants will first complete an online interactive education module, which will include information on the health risks of sitting too much, the benefits of reducing sitting, think about their own sitting time and set goals for reducing sitting. This will be followed by access to a list of tools that participants will select from to use during the intervention (e.g. mobile phone and computer apps, and wearable devices that track and provide feedback on sitting). Support will also be given by a health coach at the start and then approximately 2, 6 and 12 weeks into the intervention to help reduce sitting. We will take measures of sitting, physical activity and health (including physical function, body fat, blood sugar and cholesterol, blood pressure, and questionnaires) before the intervention begins and then 3 and 6 months after the first set of measures. We will assess how many people we recruit for the study, how many complete the study and each of the measurements and gather participants' thoughts about the intervention and measurements we take.

What are the possible benefits and risks of participating?

Participants may experience improvements in the participant's health from receiving the intervention; this includes the control group who can have the intervention at the end of the study. We are hoping this study leads to a larger study that may help to change healthcare for people with Type 2 diabetes. Participants will receive £30 of shopping gift vouchers if participants take part in all of the data collection and return the activity monitor each time participants have worn it. Participants will be reimbursed travel expenses for any visits participants make to the university as part of this study.

There is a very small risk of cross-infection when taking blood samples. We will take these samples in line with best practice guidelines to minimise this risk. There is a small chance of skin irritation from the dressing used to attach the activity monitor to the participant's skin. If this happens, the activity monitor can be removed immediately and the problem discussed with the research team. When we measure how well the participant's body functions there may be a risk of injury. During these measures, participants will need to wear suitable footwear to minimise the risk of falling and do the tests without any obvious trip hazards.

Where is the study run from?

The study is being run by Brunel University London (UK)

When is the study starting and how long is it expected to run for?

August 2018 to March 2022

Who is funding the study?

Diabetes UK

Who is the main contact?

Dr Daniel Bailey, daniel.bailey@brunel.ac.uk

Contact information

Type(s)

Scientific

Contact name

Dr Daniel Bailey

ORCID ID

<https://orcid.org/0000-0003-3772-630X>

Contact details

Brunel University London

Kingston Lane

Uxbridge

United Kingdom

UB8 3PH

+44 (0)1895 265363

daniel.bailey@brunel.ac.uk

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Integrated Research Application System (IRAS)

279157

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

22727-NHS-Apr/2020- 252, IRAS 279157

Study information

Scientific Title

A randomised-controlled feasibility study of the REgulate your Sitting Time (RESIT) intervention for reducing sitting time in individuals with Type 2 diabetes

Acronym

RESIT

Study objectives

The RESIT intervention will be acceptable and it will be feasible to deliver and evaluate the intervention

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 29/06/2020, West of Scotland National Health Service Research Ethics Committee 1 (West of Scotland Research Ethics Service, Ward 11, Dykebar Hospital, Grahamston Road, Paisley, PA2 7DE, UK; +44 (0)141 3140213; WoSREC5@ggc.scot.nhs.uk), ref: 20/WS/0080

Study design

Mixed-methods randomized controlled feasibility trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Type 2 diabetes

Interventions

Participants will be randomly allocated to a control or intervention group for 6 months (an independent researcher will use computer generated lists for randomisation of participants in a 1:1 (intervention:control) ratio with a fixed block size of four).

The control participants will receive normal diabetes healthcare as usual.

Intervention participants will first complete an online interactive education module, which will include information on the health risks of sitting too much, the benefits of reducing sitting, think about their own sitting time and set goals for reducing sitting. This will be followed by access to a list of tools that participants will select from to use during the intervention (e.g. mobile phone and computer apps, and wearable devices that track and provide feedback on sitting). Support will also be given by a health coach at the start and then approximately 2, 6 and 12 weeks into the intervention to help reduce sitting.

Measures of sitting, physical activity and health (including physical function, body fat, blood sugar and cholesterol, blood pressure, and questionnaires) will be taken before the intervention begins and then 3 and 6 months after the first set of measures.

Intervention Type

Behavioural

Primary outcome(s)

Feasibility outcome measures calculated at the end of the study:

1. Participant eligibility will be calculated as: $\frac{\text{participants eligible}}{\text{participants assessed for eligibility}} \times 100$
2. Recruitment rate will be calculated as: $\frac{\text{participants randomised}}{\text{number of eligible participants screened}} \times 100$
3. Retention rate will be calculated as: $\frac{\text{participants completing the intervention}}{\text{participants enrolled}} \times 100$
4. Completion rates for the data collection measures will be calculated as: $\frac{\text{participants providing full outcome data}}{\text{participants completing the study}} \times 100$
5. Acceptability of the intervention will be evaluated using self-report questionnaires that include closed and open-ended questions around engagement with the different intervention components. Self-report questionnaires will be completed by all participants to explore if the data collection measurements affected their behaviour. Semi-structured interviews will be completed with a subset of intervention participants to examine acceptability of the intervention. Semi-structured interviews will also assess the suitability of data collection procedures with a subset of control and intervention participants

Key secondary outcome(s)

1. Sitting, standing and stepping: this will be measured using the activPAL4 activity monitor worn on the thigh for 24 hours/day for eight consecutive days
Measured at baseline, 3 months and 6 months:
2. Body composition, anthropometry and blood pressure:
 - 2.1. Height will be measured using a stadiometer
 - 2.2. Weight and body fat % will be measured using a bioelectrical impedance analysis device
 - 2.3. Blood pressure will be measured on the right arm in a seated position using an automatic monitoring device
3. Biochemical measures: A finger prick sample technique will be used to collect fasting capillary blood samples to measure glycated haemoglobin, lipid profile and glucose concentrations
4. Psychological, sleep, musculoskeletal and wellbeing measures:
 - 4.1. Perceived fatigue will be measured using the Chalder Fatigue Scale
 - 4.2. Self-efficacy for reducing sitting will be assessed using an adapted Physical Exercise Self-Efficacy Scale
 - 4.3. A perceived sense of control over one's actions and outcomes will be assessed using the Generalised Self-Efficacy Scale and the Cohen Perceived Stress questionnaire will assess

perceived stress

4.4. The Positive and Negative Affect Scale will measure positive and negative mood

4.5. The World Health Organization Five Well-Being Index will measure psychological wellbeing and quality of life will be measured using the WHOQOL-BREF questionnaire

4.6. The Pittsburgh Sleep Quality Index questionnaire will assess sleep quality and duration and musculoskeletal symptoms will be measured using the Standardised Nordic Questionnaire.

5. Physical function: this will be assessed using the Short Physical Performance Battery (which includes standing balance, walking speed and rising from a chair) and handgrip strength

Completion date

10/03/2022

Eligibility

Key inclusion criteria

1. Aged 18 - 85 years
2. Diagnosed with type 2 diabetes
3. Able to ambulate unassisted (with or without the use of a walking aid)
4. Self-report sitting for ≥ 7 h/day

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Mixed

Lower age limit

18 years

Upper age limit

85 years

Sex

All

Total final enrolment

70

Key exclusion criteria

1. Using insulin medication
2. Unable to communicate in English
3. Pregnant
4. Cognitive or physical conditions interfering with the ability to stand and ambulate

Date of first enrolment

10/09/2020

Date of final enrolment

31/03/2021

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

Brunel University London

Kingston Lane

Uxbridge

England

UB8 3PH

Sponsor information

Organisation

Brunel University London

ROR

<https://ror.org/00dn4t376>

Funder(s)

Funder type

Charity

Funder Name

Diabetes UK

Alternative Name(s)

The British Diabetic Association, DIABETES UK LIMITED, British Diabetic Association

Funding Body Type

Private sector organisation

Funding Body Subtype

Trusts, charities, foundations (both public and private)

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

All data generated or analysed during this study will be included in the subsequent results publication

IPD sharing plan summary

Other

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		24/04/2024	26/04/2024	Yes	No
Protocol article		19/03/2021	08/03/2022	Yes	No
HRA research summary			26/07/2023	No	No
Other publications	Process evaluation	16/02/2026	17/02/2026	Yes	No
Participant information sheet	version v4	26/06/2020	07/08/2020	No	Yes
Participant information sheet	version 6.0		23/01/2023	No	Yes
Study website		11/11/2025	11/11/2025	No	Yes