

Investigating whether advanced MR imaging can be used to demonstrate trapped nerves in healthy volunteers

Submission date 04/08/2020	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered
		<input checked="" type="checkbox"/> Protocol
Registration date 15/10/2020	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
Last Edited 19/10/2021	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Currently standard MRI scans for trapped nerves in the neck are performed using sequential horizontal and vertical cuts through the spine separated by 2 or 3 mm. However, the nerves travel in a canal that is neither in the horizontal or vertical plane and the nerve itself is 2 to 3 mm in diameter. Consequently, nerve root compression can be rather poorly demonstrated on standard MRI sequences. Furthermore, the currently published scoring systems are not well validated and therefore rarely used in clinical practice. This study will use healthy volunteers to develop the technique – the changes are so common that almost all healthy volunteers without symptoms over the age of 40 will have some degenerative neck disease.

Who can participate?

Healthy volunteers aged over 40

What does the study involve?

The researchers will be using standard MRI techniques in a modified plane to image the nerves in the neck at high resolution as they leave the spine. The scans will be angled so that they cut exactly along and across the nerve canal. The researchers will also develop a scoring system based on the modified plane MRI scans and measure its reliability by comparing the scores of six different observers. They will measure these variables for the established, published scoring techniques that use standard MRI sequences and will compare the modified plane MRI scans with standard MRI scans to establish which is better.

What are the possible benefits and risks of participating?

Once the researchers have found the best way to perform and score these MRI scans, they will be able to build on this by applying the newly developed techniques to patients being considered for surgery. They expect that the new technique will permit better selection of patients for surgery and inform decisions on whether to perform surgery from the front or from the back of the neck. There are no risks to the participants expected through the use of a clinical

MRI machine. All images will be reviewed by a consultant neuroradiologist to exclude unexpected findings. Should an unexpected finding be discovered the participant's GP will be informed and the appropriate clinical referral will be made.

Where is the study run from?
Leeds General Infirmary (UK)

When is the study starting and how long is it expected to run for?
September 2018 to August 2022

Who is funding the study?
1. Leeds Neurosurgical Research Fund (UK)
2. Royal College of Surgeons of England (UK)

Who is the main contact?
Dr James Meacock
James.Meacock@nhs.net

Contact information

Type(s)
Scientific

Contact name
Dr James Meacock

ORCID ID
<https://orcid.org/0000-0002-2775-1838>

Contact details
Leeds General Infirmary
Great George Street
Leeds
United Kingdom
LS1 3EX
+44 (0)1133927425
James.Meacock@nhs.net

Type(s)
Public

Contact name
Mr Simon Thomson

ORCID ID
<https://orcid.org/0000-0003-4827-1961>

Contact details
Leeds General Infirmary
Great George Street
Leeds

United Kingdom
LS1 3EX
+44 (0)1133928567
simon.thomson1@nhs.net

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Integrated Research Application System (IRAS)

272604

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

IRAS 272604

Study information

Scientific Title

Assessing cervical foraminal stenosis: volumetric MRI study in healthy volunteers

Study objectives

Modified plane imaging has the same internal consistency as scoring performed on standard axial images.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 26/06/2020, Leeds Teaching Hospitals NHS Trust (R&I Office, c/o Anne Gowing - Leeds General Infirmary, Great George St, Leeds LS1 3EX, UK; +44 (0)113 243 2799; anne.gowing@nhs.net, ltht.researchoffice@nhs.net)

Study design

Non-randomised single institution feasibility/pilot study

Primary study design

Observational

Study type(s)

Diagnostic

Health condition(s) or problem(s) studied

Evaluation of cervical foraminal stenosis in healthy volunteers

Interventions

Three-dimensional volumetric data of the cervical spine will be obtained from an MRI scan. The anatomy of the cervical nerve root canal will be analysed to measure areas of nerve root compression using two-dimensional projections in the plane of the nerve root and with automated mathematical modelling measurements.

Kim System:

Grade 0

Grade 1

Grade 2

Will get number and percentage of each grade (Per cervical level for the right and level sides)

Grade 0, normal – absence of neural foraminal stenosis with narrowest width of neural foramen more than extraforaminal nerve root

Grade 1, non-severe cervical neural foraminal stenosis, including narrowest width of neural foramen same or less than (but more than 50% of) extraforaminal nerve root width.

Grade 2, severe cervical neural foraminal stenosis, including narrowest width of neural foramen same or less than 50% of extraforaminal nerve root width

Inter-rater reliability

Agreement percentage

Kappa value interpretations were poor ($\kappa < 0.1$), slight ($0.1 \leq \kappa \leq 0.2$), fair ($0.2 < \kappa \leq 0.4$), moderate ($0.4 < \kappa \leq 0.6$), substantial ($0.6 < \kappa \leq 0.8$), and nearly perfect ($0.8 < \kappa \leq 1.0$)

Intraclass correlation coefficient (ICC) less than 0.40 indicate poor reproducibility, values of 0.40–0.75 indicate fair to good reproducibility, and values greater than 0.75 indicate excellent reproducibility

Intra-rater reliability

Agreement percentage

Kappa value interpretations were poor ($\kappa < 0.1$), slight ($0.1 \leq \kappa \leq 0.2$), fair ($0.2 < \kappa \leq 0.4$), moderate ($0.4 < \kappa \leq 0.6$), substantial ($0.6 < \kappa \leq 0.8$), and nearly perfect ($0.8 < \kappa \leq 1.0$)

Intraclass correlation coefficient (ICC) less than 0.40 indicate poor reproducibility, values of 0.40–0.75 indicate fair to good reproducibility, and values greater than 0.75 indicate excellent reproducibility

The inter and intra rater reliability will be applied to all of the grading systems used.

Modified Kim System:

Grade 0

Grade 1

Grade 2

Will get number and percentage of each grade (Per cervical level for the right and level sides)

Grade 0 signifies that the narrowest portion of the cervical neural foramen is $>80\%$ of the width of the extraforaminal nerve root (FR ratio $>80\%$).

Grade 1 indicates that the narrowest portion of the cervical neural foramen is $<80\%$ but $>50\%$ of the width of the extraforaminal nerve root ($50\% < \text{FR ratio} \leq 80\%$)

Grade 2 for the mKim system is the same as that for the Kim system

Park System:

Grade 0

Grade 1

Grade 2

Grade 3

Will get number and percentage of each grade (Per cervical level for the right and level sides)

Grade 0, oblique sagittal plane of the cervical neural foramen shows no significant stenosis and no perineural fat obliteration

Grade 1, mild (below 50% of nerve root circumference) perineural fat obliteration. No morphological change of the nerve root is seen

Grade 2, moderate (above 50% of nerve root circumference) perineural fat obliteration. No morphological change of the nerve root is seen.

Grade 3, collapsed nerve root and morphological change of the nerve root. Severe perineural fat obliteration is also combined

Anterior v posterior compression (all views):

a) Measurement of anterior compression from bony margin (Green)

b) Measurement of posterior compression from bony margin (Green)

c) Type of compression is it disc, osteophyte, both or unknown. Based on T2 signal

Laterality of compression (all views):

a) Measurement of the point of maximum compression. This is a measurement from the apex of the ligamentum flavum to the maximum compression (Red)

b) Is the point of maximal compression:

i. Medial to the root canal

ii. Proximal 50% of the nerve root canal

iii. Distal 50% of the nerve root canal

Length of compression (all views):

Length of neuroforamina diameter that is less than uncompressed nerve root diameter (Yellow) or 2.6mm if nerve diameter is unrecordable.

Inter and intra rater reliability as above for Kim

Subjective score (all views):

a) Mild

b) Moderate

c) Severe

Intervention Type

Device

Phase

Not Applicable

Primary outcome(s)

The inter-rater reliability for the Kim grading system on measuring the narrowest diameter of the nerve root canal. This will be applied to the standard axial images and modified plain images. Success will be determined if there is a significant improvement in the inter-observer reliability.

The MRI scan will be graded once to gain the initial reading (which will then be used for inter-rater correlation) and then the same rater will grade the same scan again to determine intra-rater correlation.

Key secondary outcome(s)

1. The inter-rater reliability for subjective score, Kim grade, Modified Kim Grade, Park Grade, length and laterality of compression
2. The intra-rater reliability for narrowest diameter, subjective score, Kim grade, Modified Kim Grade, Park Grade, length and laterality of compression

The MRI scan will be graded once to gain the initial reading (which will then be used for inter-rater correlation) and then the same rater will grade the same scan again to determine intra-rater correlation.

Completion date

31/08/2022

Eligibility

Key inclusion criteria

1. Age over 40 years
2. Able to provide fully informed written consent
3. Able to lie flat for 1 hour in an MRI scanner
4. Females of childbearing age must be using effective contraception
5. Sufficient understanding of English to participate in the trial

Participant type(s)

Healthy volunteer

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Total final enrolment

5

Key exclusion criteria

1. Cervical myelopathy
2. History of cervical trauma
3. Evidence of suspected or histologically proven tumour
4. Previous cervical spine surgery
5. Non-MRI compatible implantable device e.g. pacemaker
6. Unable to have MRI scan due to claustrophobia
7. Female participants must not be pregnant and if of childbearing age must be using adequate contraception

Date of first enrolment

01/03/2021

Date of final enrolment

31/08/2021

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

Leeds Teaching Hospitals NHS Trust

Great George Street

Leeds

United Kingdom

LS1 3EX

Sponsor information

Organisation

Leeds Teaching Hospitals NHS Trust

ROR

<https://ror.org/00v4dac24>

Funder(s)

Funder type

Research organisation

Funder Name

Leeds Neurosurgical Research Fund

Funder Name

Royal College of Surgeons of England

Alternative Name(s)

RCS England, RCS ENG, The Royal College of Surgeons of England, RCS

Funding Body Type

Private sector organisation

Funding Body Subtype

Universities (academic only)

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from Mr Simon Thomson (simon.thomson1@nhs.net) on study publication for 7 years, for researchers wishing to confirm the findings or undertake meta-analysis, by secure data transfer, consent from participants was obtained, data will be anonymised.

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Protocol file	version V7	15/10/2020	15/10/2020	No	No