

How much volume of a local anesthetic is necessary to accomplish a nervus saphenous block

Submission date 22/01/2016	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 25/01/2016	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 25/01/2016	Condition category Other	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

When a person has had surgery on their knee or inside of the lower leg (tibia), effective pain relief is very important part of recovery and rehabilitation. Ideally, pain relief would allow a patient to move their leg and be able to stand or walk (mobilise) straight away after an operation, without complications. An emerging form of providing pain relief for these patients is by using a technique called a saphenous nerve block. This involves injecting a local anaesthetic (numbing agent) into the saphenous nerve. It is the longest branch coming off the femoral nerve (the main nerve that supplies the leg) and is a pure sensory nerve, meaning that it is responsible for detecting sensations (such as temperature, touch or pain) and does not affect movement. The nerve block procedure works by “turning off” (blocking) the pain signals coming from the affected area and preventing excess inflammation (swelling). Many nerve block procedures also affect motor nerves (responsible for controlling movement), and so can prevent patients from moving, possibly slowing down their recovery. It is well known that using a large volume of anaesthetic in a nerve block procedure can lead to unwanted spreading to motor nerves (motor blockade). The aim of this study is to find the minimum effective local anesthetic dose to perform a saphenous nerve block in healthy volunteers.

Who can participate?

Healthy adults aged between 18 and 65.

What does the study involve?

All participants receive an ultrasound-guided saphenous nerve block on their left leg. This involves using an ultrasound probe (which uses high-frequency sound waves) to visualize the nerves in the leg, to guide the injection of mepivacaine (local anaesthetic) along the saphenous nerve. The needle which injects the mepivacaine enters the skin at mid-thigh level, and is guided into the correct place using the ultrasound images of the leg, projected onto a screen. The amount of anaesthetic needed to achieve a nerve block in each patient determines how much is given to the next patient (i.e. increasing the dose by 1ml each time). Thirty minutes after the

block, all participants have the area of skin that is controlled by the saphenous nerve pricked with a pin and are asked whether they can feel it so that the most effective dose can be worked out.

What are the possible benefits and risks of participating?

There are no direct benefits for participants taking part in the study, although they do receive financial compensation for travel costs. There is a very small risk of infection, bleeding or nerve damage, however this is unlikely.

Where is the study run from?

Academic Medical Center (Netherlands)

When is the study starting and how long is it expected to run for?

November 2014 to November 2015

Who is funding the study?

Academic Medical Center (Netherlands)

Who is the main contact?

Dr Werner ten Hoope

w.tenhoope@amc.uva.nl

Contact information

Type(s)

Scientific

Contact name

Dr Werner ten Hoope

ORCID ID

<https://orcid.org/0000-0001-7050-8019>

Contact details

Academisch Medisch Centrum (Academic Medical Center)

Meibergdreef 9

Amsterdam

Netherlands

1105 AZ

+31 20 5669111

w.tenhoope@amc.uva.nl

Additional identifiers

Clinical Trials Information System (CTIS)

2014-004672-38

Protocol serial number

20141125 V2

Study information

Scientific Title

Minimum local anaesthetic volumes for saphenous nerve block: A dose-finding study

Study objectives

The aim of this study is to characterize the minimum volume of local anesthetic volume effective saphenous nerve block in volunteers.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Medical Ethical Committee Academic Medical Centre, 05/01/2015, ref: 2014_348

Study design

Non-randomized single-centre single group dose escalation study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Saphenous nerve block

Interventions

Patients will receive a distal ultrasound-guided saphenous nerve block. Briefly, the ultrasound probe will be used to locate the femoral artery over the medial aspect of the thigh. The artery will be followed distally until it moves posteriorly through the adductor hiatus. Block will be performed 2 cm to this location. At this point, the saphenous nerve is in immediate proximity, medial to the artery. The block is performed using an amount of mepivacaine 2% dictated by the Dixon model.

Each patient's response to the nerve block determines the volume of mepivacaine for the next patient. When successful saphenous nerve block is achieved, the volume for the next patient will be decreased with 1ml mepivacaine. Conversely, when no block can be detected, the volume for the next study patient will be increased with 1 ml mepivacaine. As described by Dixon, the study is finished after three oscillations (up/down) around a given volume. From these data, ED95 and ED50 can be calculated.

After the nerve block, an assessment of block onset will be performed at the apex patellae and, after 20 minutes, assessment of the main study parameter, block yes/no. After this, motor weakness is determined, and the sensory area anesthetized is mapped on the skin surface of each volunteer. Finally, block duration is determined by measuring every 15 minutes until sensory block disappears. The total duration of the intervention takes 4 hours maximum for a complete block offset. When the block wears of the volunteer completed the study.

Intervention Type

Procedure/Surgery

Primary outcome(s)

Effective volume to produce complete neural blockade is determined using a pinprick in the sensory area of the saphenous nerve to provoke a dichotomous yes/no answer, 30 minutes after block.

Key secondary outcome(s)

1. Influence of the volume on duration of sensory blockade is measured using pinpricks at the apex of the patella, medial aspect of the medial facet of the tibial plateau and the skin over the medial ankle at 2, 4, 6, 8, 10, 15, 20, 25, and 30 minutes, and then every 15 minutes until block resolution
2. Extent of potential motor blockade is measured using the MRC-scale 20 minutes after block onset
3. The maximum spread of sensory block on the skin surface is measured using pinpricks to establish the edge of the anaesthetised area 20 minutes after block onset

Completion date

01/11/2015

Eligibility

Key inclusion criteria

1. Aged between 18 and 65 years
2. ASA physical status I-II (healthy)

Participant type(s)

Healthy volunteer

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

1. Allergy against local anesthetics
2. Contraindication for saphenous nerve block (infection at injection site, local pathology)
3. Ingestion of any pain medication within the past 24 hours
4. Pregnancy or breastfeeding

Date of first enrolment

12/02/2015

Date of final enrolment

01/09/2015

Locations

Countries of recruitment

Netherlands

Study participating centre**Academic Medical Center**

Meibergdreef 9

Amsterdam

Netherlands

1105AZ

Sponsor information

Organisation

Academisch Medisch Centrum (Academic Medical Center)

ROR

<https://ror.org/03t4gr691>

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Academisch Medisch Centrum (Academic Medical Center)

Alternative Name(s)

Academic Medical Center, AMC

Funding Body Type

Private sector organisation

Funding Body Subtype

Universities (academic only)

Location

Netherlands

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes