# Defining how antibiotics disrupt lung immune responses in asthma and health

Submission date	<b>Recruitment status</b> Recruiting	[X] Prospectively registered			
13/06/2025		☐ Protocol			
Registration date 18/06/2025	Overall study status Ongoing Condition category	Statistical analysis plan			
		Results			
Last Edited		Individual participant data			
06/10/2025	Other	[X] Record updated in last year			

## Plain English summary of protocol

Background and study aims

This is a trial designed to help us better understand how changes in the bacteria that live in everyone's lungs and gut affect inflammation in the lungs, particularly in asthma. We will find this out by looking at how both things change before and after participants take a common antibiotic medicine called amoxicillin.

We are looking at these changes in people both with and without asthma. We are interested in whether any changes antibiotics cause are bigger and more significant in people who have asthma, and the potential impact for how people with asthma are treated with antibiotics in the future.

When antibiotics are used, they don't just kill the bacteria causing an infection – they will also reduce the number of normal bacteria that usually live in the body, causing no harm. We are interested in whether changing your population of healthy bacteria (with antibiotics), especially in the gut where the vast majority live, changes the type of inflammation in the lungs. If it does, this would be a reason for doctors to be more careful when prescribing them. Especially as we also know many simple infections would have gotten better on their own without antibiotics. We are using amoxicillin in this trial as it is one of the most common antibiotics.

#### Who can participate?

Adults aged 18 to 80 years old, in generally good health, with and without mild asthma.

#### What does the study involve?

The study involves answering health questionnaires and assessment of lung function, followed by two bronchoscopy procedures where samples of lung fluid will be taken. Bronchoscopy involves the use of a camera at the end of a very narrow flexible tube (usually inserted through the nose) to look at the lungs. The narrow tube also has a suction port to take fluid samples. Between the procedures a course of amoxicillin or an inactive placebo tablet is taken. Participants will also provide stool samples and have blood taken at the two procedure visits and at intervals over 1 year afterwards.

What are the possible benefits and risks of participating? There are no expected direct health benefits to participants. There are risks associated with antibiotics (allergy, nausea, diarrhoea). There are risks associated with bronchoscopy, including but not limited to: discomfort during the procedure; coughing, during and immediately afterwards; temporary low oxygen levels; wheeze; nose bleeds; fainting. Sedation used for bronchoscopy may cause nausea, vomiting, or lightheadedness. There are some more serious risks associated with bronchoscopy, such as pneumothorax (air trapped in the chest), bleeding in the lungs and death. All these complications are very rare (risks of 1:1000 or lower). Sedation can rarely cause a depressed breathing rate or low blood pressure. We are only including participants who have no medical conditions or history which would increase their individual risk of serious complications from bronchoscopy or sedation.

Where is the study run from?

The study will run at the Manchester Clinical Research Facility and the North-West Lung Centre at Wythenshawe Hospital (UK)

When is the study starting and how long is it expected to run for? November 2024 to November 2027

Who is the main funder?

The Medical Research Council through an Experimental Medicine Grant (Grant Code: MR /Y008812/1) (UK)

Who is the main study contact? Prof Tim Felton, Tim.felton@manchester.ac.uk

## Contact information

## Type(s)

Public, Scientific, Principal investigator

#### Contact name

Prof Tim Felton

#### **ORCID ID**

https://orcid.org/0000-0001-6868-6633

#### Contact details

Division of Immunology, Immunity to Infection and Respiratory Medicine NIHR Biomedical Research Centre University of Manchester Wythenshawe Hospital Southmoor Road Manchester United Kingdom M23 9LT +44 (0)161 701 2669 tim.felton@manchester.ac.uk

## Additional identifiers

Clinical Trials Information System (CTIS)

## Integrated Research Application System (IRAS)

354874

## ClinicalTrials.gov (NCT)

Nil known

#### Protocol serial number

Grant Code: MR/Y008812/1

## Study information

#### Scientific Title

Defining mechanisms underpinning antibiotic mediated disruption of pulmonary immune responses

## Acronym

**DEMAND** 

## Study objectives

A causal link between disruption of the human microbiome and type two inflammation in the lungs exists - and is of different character and/or magnitude in healthy volunteers and asthmatics.

## Ethics approval required

Ethics approval required

## Ethics approval(s)

approved 24/09/2025, London - Bloomsbury Research Ethics Committee (2 Redman Place, Stratford, London, E20 1JQ, United Kingdom; +44 207 104 8384; bloomsbury.rec@hra.nhs.uk), ref: 25/LO/0649

## Study design

Single-centre interventional double-blinded randomized controlled trial

## Primary study design

Interventional

## Study type(s)

Other

## Health condition(s) or problem(s) studied

Influence of microbiome disruption on expressed immune phenotype in health and in asthma

#### **Interventions**

Two cohorts: healthy volunteers (n = 34), asthmatics (n = 34). Total participants n = 68. Within each cohort, participants are randomised using double-blinded sequential block

randomisation to ensure equal allocations to one of two interventions:

- 1. Amoxicillin 500 mg, three times daily, 7 days
- 2. Visually matched inactive placebo, three times daily, 7 days

The study involves answering health questionnaires and assessment of lung function, followed by two bronchoscopy procedures where samples of lung fluid will be taken. Bronchoscopy involves the use of a camera at the end of a very narrow flexible tube (usually inserted through the nose) to look at the lungs. The narrow tube also has a suction port to take fluid samples. Between the procedures a course of amoxicillin or an inactive placebo tablet is taken. Participants will also provide stool samples and have blood taken at the two procedure visits and at intervals over 1 year afterwards.

## Intervention Type

Drug

#### Phase

Not Applicable

## Drug/device/biological/vaccine name(s)

Amoxicillin

## Primary outcome(s)

- 1. In-depth immune cell characterisation to quantify the degree of expression of interleukins and immune cell surface markers associated with type 2 inflammation in paired broncho-alveolar fluid (BAL) and blood at baseline (day 0) and post-intervention (day 7)
- 2. Matched to fraction of exhaled nitric oxide (FeNO) measurements (in parts per billion [ppb]) at baseline (day 0) and post-intervention (day 7)
- 3. Blood immune cell characterisation to quantify the degree of expression of interleukins and immune cell surface markers associated with type 2 inflammation will be repeated at follow-up at approximately 6, 12, 26 and 52 weeks post-intervention
- 4. FeNO measurements (ppb) will be repeated at follow-up at approximately 6, 12, 26 and 52 weeks post-intervention

## Key secondary outcome(s))

- 1. Characterisation of the gut bacterial microbiome composition pre- and post-intervention (day 0 and day 7) by microbiome sequencing techniques performed on paired BAL and stool samples, including analysis of the change in microbiota attributable to antibiotic administration
- 2. Assessment of the recovery of the gut microbiome over the following year through serial microbiome sequencing on stool samples at approximately 6, 12, 26 and 52 weeks post-intervention
- 3. Quantification and characterisation of stool and blood bacterial metabolites by short chain fatty acid assay at baseline (day 0), post-intervention (day 7) and at approximately 6, 12, 26 and 52 weeks post-intervention

## Completion date

02/11/2027

# Eligibility

Key inclusion criteria

- 1. Males and Females
- 2. Mild asthma patients aged between 18 and 80 years, inclusive
- 3. Healthy volunteers aged between 18 and 80 years, inclusive
- 4. Non-smokers or Ex-smokers with fewer than 10 pack years and >6 months abstinence
- 5. Healthy volunteers: no clinically relevant abnormalities based on history, examination, vital signs, and no history of respiratory disease
- 6. Asthma cohort participants only: confirmed airway responsiveness via methacholine challenge at screening visit
- 7. Asthma cohort participants only: good or partial symptom control according to GINA classification and corticosteroid naïve
- 8. Capacity to consent

## Participant type(s)

Healthy volunteer, Patient

## Healthy volunteers allowed

No

## Age group

Adult

## Lower age limit

18 years

## Upper age limit

80 years

#### Sex

All

## Key exclusion criteria

- 1. Have received any systemic antibiotic medication:
- 1.1. To treat an infection arising during the participant's potential study participation
- 1.2. Within the year prior to potential inclusion
- 2. Have received any topical or systemic treatment for oral thrush in the year prior to inclusion.
- 3. Have received any leukotriene antagonist medication within the year prior to the inclusion.
- 4. History of hypersensitivity or allergy to penicillins or other beta-lactam antibiotics.
- 5. Treatment for asthma beyond intermittent inhaled salbutamol use— i.e. regular inhaled corticosteroids or above or any use of systemic or inhaled corticosteroids during or within one year prior to inclusion in the study
- 6. Baseline FEV1 <70% (asthma cohort) or <80% (healthy volunteers).
- 7. Current smokers or ex-smokers with over 10 pack years or under 6 months abstinence.
- 8. Current regular smokers (over once a month) of cannabis or other illicit drugs.
- 9. Conditions or medications that may affect the safety of bronchoscopy, sedation, amoxicillin administration, or have an independent effect on airway immune responses and reactivity: a. any pulmonary or chest wall disease other than asthma (including but not limited to chronic obstructive pulmonary disease, interstitial lung disease, significant chest wall deformities); b. other disease states that increase the risk of hypoxaemia, or the risk of serious sequalae of hypoxaemia; c. regular use of anti-histamine medications unless participant able to discontinue during bronchoscopy phase of the study; d. regular anticoagulant therapy; e. upper or lower

respiratory tract infection within 4 weeks prior to enrolment; f. history of adverse reactions to sedative or anaesthetic agents beyond expected side effects.

- 10. Conditions or medications that have an independent effect on the host gut or respiratory flora and microbiome (including but not limited to inflammatory bowel disease, irritable bowel syndrome, small intestinal bacterial overgrowth)
- 11. Pregnant
- 12. Other severe, acute or chronic, medical or psychiatric conditions that may increase the risk associated with trial participation or interfere with the interpretation of trial results.
- 13. Inability to collect and provide a stool sample.
- 14. Participation in other interventional research trials within the study period except where appropriate cross-enrolment has been agreed by both Chief Investigators.

**Date of first enrolment** 01/10/2025

Date of final enrolment 03/11/2026

## Locations

**Countries of recruitment**United Kingdom

England

Study participating centre
Wythenshawe Hospital
Southmoor Road
Wythenshawe
Manchester
United Kingdom
M23 9LT

# Sponsor information

## Organisation

University of Manchester

## ROR

https://ror.org/027m9bs27

# Funder(s)

Funder type

## **Funder Name**

Medical Research Council

## Alternative Name(s)

Medical Research Council (United Kingdom), UK Medical Research Council, MRC

## **Funding Body Type**

Government organisation

## Funding Body Subtype

National government

#### Location

**United Kingdom** 

## **Results and Publications**

## Individual participant data (IPD) sharing plan

Anonymised transcriptomic, immunological, metagenomic and metabolomic datasets, alongside anonymised study metadata, will be made publicly available via long-term storage in suitable repositories. This will be made available approximately 12 months after the study conclusion. Access to these public repositories is unrestricted.

Decisions on earlier access of research data to new users will be made by the study leadership team and documented for subsequent independent review. An external independent advisor will be identified to periodically review the outcomes of access requests.

Datasets will be available via the University of Manchester Pure research information repository (http://www.staffnet.manchester.ac.uk/pure/).

Participants will be consented for this data sharing at the outset of their involvement in the study.

## IPD sharing plan summary

Stored in publicly available repository

## **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Study website	Study website	11/11/2025	11/11/2025	No	Yes