

Blood vessel problems in people with prediabetes and diabetes who have healthy heart arteries

Submission date 21/09/2024	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 23/09/2024	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 06/11/2024	Condition category Circulatory System	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Diabetes mellitus (DM) is a global health issue associated with increased heart disease risk. Endothelial dysfunction (narrowed blood vessels) is a known precursor to atherosclerosis (buildup of substances on blood vessel walls) and heart disease, and its role in DM complications is well-documented. There is limited information on endothelial (blood vessel wall) function in prediabetic patients evaluated using flow-mediated dilation (FMD), and studies have not excluded patients with known atherosclerosis or coronary artery disease. This study aimed to evaluate endothelial function using flow-mediated dilation from the brachial (upper arm) artery of DM and prediabetes patients who had normal coronary arteries.

Who can participate?

Patients aged 21-80 years evaluated in the Cardiology outpatient clinic who are found to have normal coronary arteries on coronary angiography, including patients with DM, prediabetes, and non-diabetics

What does the study involve?

The study involves the measurement of flow-mediated dilation of participants according to the established standard methods. The participants are divided into three groups according to their blood sugar status which is determined by measurement of blood glucose levels in the fasting state and after glucose loading.

What are the possible benefits and risks of participating?

The endothelial function of the brachial arteries of the participants will be measured using a noninvasive method. This will enable further determination of heart disease risk and preventive action. The glucose tolerance test and measurement of FMD using ultrasound from the brachial artery carries no risk for the participants.

Where is the study run from?

Ankara Guven Hospital (Türkiye)

When does the study start and how long is it expected to run for?

May 2008 to December 2010

Who is funding the study?

Investigator initiated and funded

Who is the main contact?

Dr İnci Aslı Atar, asliatar@gmail.com, inciasli.atar@saglik.gov.tr

Contact information

Type(s)

Public, Scientific, Principal Investigator

Contact name

Dr İnci Aslı Atar

Contact details

Ankara City Hospital

Ankara

Türkiye

06830

+90 (0)3124572525

inciasli.atar@saglik.gov.tr

Additional identifiers

EudraCT/CTIS number

Nil known

IRAS number

ClinicalTrials.gov number

Nil known

Secondary identifying numbers

GHE 190

Study information

Scientific Title

Evaluation of endothelial dysfunction in prediabetes and diabetes mellitus in patients with normal coronary arteries using flow-mediated dilation of brachial artery

Study objectives

Diabetes mellitus (DM) is an important global pandemic disease and is associated with an increased incidence and mortality of cardiovascular disease. Endothelial dysfunction is considered a precursor to atherosclerosis and cardiovascular disease (CVD) and the role of endothelial dysfunction in the pathogenesis of both microvascular and macrovascular complications is well-documented in patients with DM. Prediabetes is an important diagnosis,

enabling physicians to implement measures to prevent the progression to DM and its related complications or to slow the disease process. Evaluating endothelial function in prediabetic patients may allow for the early detection and treatment of issues before further complications arise. One of the most widely used non-invasive methods for assessing endothelial function in arteries is flow-mediated dilation (FMD) of the brachial artery. The relationship between the presence and severity of coronary artery disease and the degree of endothelial dysfunction is well established. Therefore, assessing endothelial function in a prediabetic population without excluding the presence of coronary artery disease presents limitations in demonstrating a causal relationship between endothelial dysfunction and prediabetes. Thus, this study aimed to evaluate the endothelial functions using FMD from the brachial artery of DM and prediabetes patients who had normal coronary arteries.

Ethics approval required

Ethics approval required

Ethics approval(s)

Approved 23/12/2008, Ankara Guven Hospital Ethics Committee (Ankara Guven Hastanesi, Kavaklıdere, Simsek Sok. No: 29 Cankaya, Ankara, 06640, Türkiye; +90 (0)3124572525; guven@guven.com.tr), ref: 190

Study design

Non-randomized study

Primary study design

Interventional

Secondary study design

Non randomised study

Study setting(s)

Hospital

Study type(s)

Diagnostic

Participant information sheet

Not available in web format, please use contact details to request a participant information sheet.

Health condition(s) or problem(s) studied

Endothelial dysfunction in patients with diabetes and pre-diabetes with normal coronary arteries

Interventions

Patients evaluated in the Cardiology outpatient clinic who are found to have normal coronary arteries on coronary angiography will be included in the study. Diabetes and pre diabetes will be diagnosed according to American Diabetes Association criteria. Participants will be grouped as diabetic, pre diabetic and normal. Flow mediated dilation will be performed following the guidelines outlined in previously published literature by an investigator proficient in vascular ultrasound techniques.

Intervention Type

Procedure/Surgery

Primary outcome measure

Flow-mediated dilation is measured by an ultrasound system equipped with a 5-10 MHz linear array transducer from the brachial artery at baseline and after vasodilation, using an ultrasound, taken once per patient

Secondary outcome measures

There are no secondary outcome measures

Overall study start date

04/05/2008

Completion date

30/12/2010

Eligibility

Key inclusion criteria

1. Patients evaluated in the Cardiology outpatient clinic who were found to have normal coronary arteries on coronary angiography
2. Aged 21-80 years

Participant type(s)

Patient

Age group

Adult

Lower age limit

21 Years

Upper age limit

80 Years

Sex

Both

Target number of participants

75

Total final enrolment

73

Key exclusion criteria

1. A history of previous percutaneous coronary intervention
2. Atrial fibrillation
3. Heart failure
4. More than mild valvular regurgitation
5. Valvular stenosis

6. Uncontrolled hypertension (systolic blood pressure ≥ 160 mmHg and diastolic blood pressure ≥ 90 mmHg in the last 6 weeks)
7. Cardiomyopathy
8. Chronic kidney or liver disease

Date of first enrolment

25/12/2008

Date of final enrolment

30/10/2010

Locations

Countries of recruitment

Türkiye

Study participating centre**Ankara Guven Hastanesi**

Kavaklıdere, Simsek Sok. No: 29 Cankaya

Ankara

Türkiye

06690

Sponsor information

Organisation

Ankara Guven Hospital

Sponsor details

Kavaklıdere, Şimşek Sokak NO: 29

Ankara

Türkiye

06640

+90 (0)3124572525

guven@guven.com.tr

Sponsor type

Hospital/treatment centre

Funder(s)

Funder type

Other

Funder Name

Investigator initiated and funded

Results and Publications

Publication and dissemination plan

Planned publication in peer-reviewed journal upon receipt of registration number

Intention to publish date

10/11/2024

Individual participant data (IPD) sharing plan

The datasets generated during and/or analyzed during the current study will be available upon request from the corresponding author (Dr İnci Aslı Atar, asliatar@gmail.com, inciasli.atar@saglik.gov.tr)

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		06/11/2024	06/11/2024	Yes	No