Comparison of the effectiveness of two different respiratory exercise training methods in prolonged ventilated patients

Submission date	Recruitment status No longer recruiting	Prospectively registered		
24/06/2020		[X] Protocol		
Registration date	Overall study status	Statistical analysis plan		
05/07/2020	Completed	[X] Results		
Last Edited 21/10/2022	Condition category Respiratory	Individual participant data		

Plain English summary of protocol

Background and study aims

Reduced breathing muscle performance contributes to the prolonged duration of mechanical ventilation (external breathing machine support) and makes it difficult to come off from the machine. Reduced breathing muscle function also impacts on non-ventilatory functions of the respiratory system including coughing and swallowing function.

We are planning to conduct a study in prolonged ventilated patients (patients on external breathing machine support) to find out the impact of different types of resistance breathing exercises. These are used in some intensive care units to strengthen breathing muscles worldwide. This study will help us to determine whether this resistance breathing exercises during prolonged ventilation are necessary or not.

Who can participate?

Patients requiring breathing support with a ventilator (external breathing machine) for more than 5 days who are suitable will be included in the study following a discussion with consultant anesthetist and clinical nurse manager.

What does the study involve?

This proposed study plans to monitor and manage breathing muscle dysfunction with multiple tests of strength and endurance and with a dual valve CE marked low-cost device that will be adapted in a novel way to train breathing muscles which help to breathe in and out. Patients will undergo four sets of ten breaths twice a day for two weeks duration with CE marked device. The strength and endurance tests of breathing muscles will be done before and after exercise training.

What are the possible benefits and risks of participating?

Benefits:

The results will help us to determine which exercises help to improve weak respiratory muscle function.

Risks:

There are no anticipated risks with this study. Respiratory assessments are commonly used in

physiotherapy. The tests are not invasive. None of the assessments have the potential to cause harm. Different exercises being compared in this study. They are both in line with the intended use of the CE marked device. We will find out from this study if there are any differences between these different exercises in improving respiratory muscle function.

Where is the study run from? Connolly Hospital Blanchardstown (Ireland)

When is the study starting and how long is it expected to run for? June 2019 to December 2021

Who is funding the study? Anglia Ruskin University (UK)

Who is the main contact?
Mr Atchuta Kishore Kothapalli, kishore.kothapalli@hse.ie

Contact information

Type(s)

Scientific

Contact name

Mr Atchuta Kishore Kothapalli

ORCID ID

https://orcid.org/0000-0002-0728-454X

Contact details

Connolly Hospital
Blanchardstown
Physiotherapy Department
Dublin
Ireland
D15 X40D
+353 16465296
kishore.kothapalli@hse.ie

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

Nil known

Study information

Scientific Title

Comparing effectiveness REspiratory Muscle Dysfunction Interventions IMT versus RMT study in prolonged ventilated patients

Acronym

REMDI

Study objectives

Can respiratory (inspiratory and expiratory) muscle training be feasible and more effective compared to inspiratory muscle training alone in prolonged ventilated patients?

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 18/06/2019, Ethics Committee (Connolly Hospital Blanchardstown, Dublin 15, D15 X40D, Republic of Ireland; +353 18213844; nancy.mcguirk@hse.ie), ref: none provided

Study design

Interventional randomized parallel trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Respiratory muscle dysfunction in prolonged ventilated patients

Interventions

Participants will either receive:

- 1. 2 weeks of inspiratory muscle training, or
- 2. 2 weeks of both expiratory and inspiratory muscle training

The same CE marked device is set at 50% threshold to train either:

- 1. Maximum inspiratory pressure (MIP), or
- 2. Both MIP and Maximum Expiratory Pressure (MEP). Both groups will undergo four sets of ten breaths twice a day for two weeks

Randomisation. To minimise selection bias during randomisation the physiotherapist who is not involved in the study and will have no interaction with any of the trial participants will randomise with concealed randomisation and will have no interaction with any of the trial participants.

There are two data collection points 1) at baseline and 2) at 2 weeks after exercise

Intervention Type

Behavioural

Primary outcome(s)

Maximum Inspiratory Pressure (MIP) measured using CE marked Micro RPM (Respiratory Pressure Monitor) at pre-(baseline) and post-intervention (2 weeks). 3 trials are taken and the highest value will be used.

Key secondary outcome(s))

Measures will be performed pre-(baseline) and post-intervention (2 weeks):

- 1. Maximum Expiratory Pressure (MEP) will be measured using a pressure manometer 3 trials are taken and the highest value will be used
- 2. Cough strength will be measured using a semi-quantitative cough strength score
- 3. Fatigue resistance index will be measured using the ratio of MIP, post-training with Inspiratory Muscle Trainer/MIP pre-training.
- 4. Rectus abdominis muscle thickness assessment will be measured using a diagnostic ultrasound machine
- 5. Hand dynamometry will be measure peripheral muscle function 3 trials are taken on the participants dominant hand and the highest value will be used

Completion date

31/12/2021

Eligibility

Key inclusion criteria

- 1. Adult patients 18 years or above admitted to general critical care unit and required mechanical ventilation support for more than five days
- 2. Patients who can respond to the verbal command and can tolerate breathing exercises such as "deep breath in and out"
- 3. Richmond Agitation-Sedation Scale score 0

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

- 1. Medically unstable during critical care admission and prolonged ventilation period before training
- 2. Use of more than minimal continuous infusion inotropic agents
- 3. Contraindication to disconnect the patient from the mechanical ventilator for RMT treatment

or IMT treatment

- 4. Acute coronary artery disease
- 5. Thoracic contraindications

Date of first enrolment

06/09/2019

Date of final enrolment

31/12/2021

Locations

Countries of recruitment

Ireland

Study participating centre Connolly Hospital Blanchardstown

Mill Rd
Abbotstown
Dublin
Ireland
D15 X40D

Sponsor information

Organisation

Connolly Hospital Blanchardstown

ROR

https://ror.org/03h5v7z82

Organisation

Anglia Ruskin University

ROR

https://ror.org/0009t4v78

Funder(s)

Funder type

University/education

Funder Name

Anglia Ruskin University

Alternative Name(s)

Funding Body Type

Private sector organisation

Funding Body Subtype

Universities (academic only)

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request. Raw data will be available for 5 years from when the results are submitted for publication.

IPD sharing plan summary

Other

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Basic results			21/10/2022	No	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Protocol file			21/10/2022	No	No