Detecting low brain oxygen levels in patients undergoing non-cardiac surgery to assess its association with post-operative delirium

Submission date	Recruitment status	Prospectively registered
29/08/2018	No longer recruiting	[] Protocol
Registration date	Overall study status	[] Statistical analysis plan
06/11/2018	Completed	[X] Results
Last Edited 22/03/2021	Condition category Mental and Behavioural Disorders	Individual participant data

Plain English summary of protocol

Background and study aims (brief description of the disease or area of study, what are the objectives/aim of the study)

The aim is to explore the use of a novel monitor, during general anaesthesia and surgery, that non-invasively measures the levels of oxygen in the brain. We will also look for the uncommon complication of delirium occurring early in the postoperative period.

Who can participate? (what are the age range and gender of the participants, can they only participate if they have a certain condition or if they are healthy volunteers?)

Specific patients having surgery at this hospital will be approached.

What does the study involve? (what interventions will be compared, will all participants receive the same treatment?)

The use of a new monitor before, during and for one hour after surgery.

What are the possible benefits and risks of participating? (what can participants gain from enrolling, are there any side effects of the treatments and if so, what are the symptoms?) No potential benefit. Trivial risk of having allergic to the adhesive on the sensor.

Where is the study run from? (what are the approximate number and names of centres taking part in this trial, if there is a lead centre, which one is it?) Royal Surrey County Hospital NHS Foundation Trust

When is the study starting and how long is it expected to run for? (what is the anticipated start date and the approximate duration of the trial?) September 2018 to January 2019 Who is funding the study? (who will be paying the costs that the trial will incur during its lifecycle?) Medtronic, the manufacturer of the sensors and device.

Who is the main contact? (if this is the same as the contact in the record, please provide the name and email address only, if different to the contact in the record, please provide the name, position they hold at the institution/organisation and their email address) Ben Creagh-Brown, bencb@nhs.net

Contact information

Type(s) Public

Contact name Dr Ben Creagh-Brown

ORCID ID http://orcid.org/0000-0002-4397-1232

Contact details Royal Surrey County Hospital Guildford United Kingdom GU27XX

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers 1.31

Study information

Scientific Title

Using near-infrared spectroscopy to detect cerebral desaturation in non-cardiac surgery and assess its relationship with post-operative delirium– a pilot study

Acronym CereOx

Study objectives

Low baseline cerebral oximetry values, or some metric of peri-operative cerebral desaturation may be more closely associated with post-operative delirium than other variables including systemic blood pressure.

Ethics approval required Old ethics approval format

Old ethics approval format

Ethics approval(s)

London - Bloomsbury Research Ethics Committee reviewed the above application on 21 June 2018. REC reference: 18/LO/1122

Study design

Prospective observational cohort study

Primary study design Observational

Secondary study design Cohort study

Study setting(s) Hospital

Study type(s) Diagnostic

Participant information sheet

Not available in web format, please use the contact details below to request a participant information sheet

Health condition(s) or problem(s) studied

Postoperative delirium

Interventions

Measurement of cerebral oximetry, from entering the anaesthetic room to 1 hour after the end of surgery

Intervention Type

Device

Primary outcome measure

The following will be assessed once the final participant has finished the study (as these measurements require the total number of patients):

1. Proportion of patients enrolled in whom we acquired interpretable regional oxygen saturation (rScO2) data (%) is assessed: 100 X total number recruited with interpretable data / total number recruited

2. Proportion of patients enrolled in whom we acquired delirium data (%) is assessed: 100 X total number recruited with delirium data / total number recruited

Secondary outcome measures

For each cohort of surgical patients, and the cohort as a whole, use descriptive statistics (proportion (%), or mean and standard deviation, or median and interquartile range – according to parametric or non-parametric distribution respectively):

1. Incidence of cerebral desaturation using three previously utilised definitions (as below).

Cerebral saturation is assessed using a cerebral oximetry machine, recorded continuously during surgery and for 1 hour after surgery.

1.1. Any occurrence of a decrease of 10% from baseline

1.2. Any occurrence of a decrease of 20% from baseline

1.3. Any occurrence of an absolute value of regional oxygen saturation (rScO2) <50%

2. Average duration of relative desaturation - time with rScO2 \leq 60%

3. Incidence of systemic hypotension using three common definitions (as below). Systolic hypotension is assessed using blood pressure readings taken during surgery. Baseline values will be obtained from the pre-operative clinic; however, if this is not possible then the pre-induction blood pressure will be used as a baseline.

3.1. Any occurrence of systolic blood pressure (BP) ≤20% of baseline

3.2. Any occurrence of systolic BP ≤30% of baseline

3.3. Any occurrence of systolic BP ≤40% of baseline

4. Average duration of systemic hypotension - time with mean arterial pressure (MAP) ≤20% of baseline

5. Incidence of hypoxaemia using three definitions (as below). Hypoxaemia will be assessed using saturation probe readings during surgery.

5.1. Any occurrence of blood oxygen saturation (SpO2) ≤ 88%

5.2. Any occurrence of SpO2 \leq 90%

5.3. Any occurrence of SpO2 \leq 92%

6. Average duration of hypoxaemia - time with SpO2 \leq 90%

7. Utilisation of vasopressor drugs (α-agonists) - we will record either a vasopressor infusion or bolus is given during surgery, and if any are given during the recovery. Time administered, drug, dose and the Mean Arterial Pressure (obtained from the anaesthetic machine) will be recorded by the clinician

8. Baseline rScO2 values

9. The incidence of delirium during the postoperative in-patient stay (up to day 7), assessed using 4AT (clinic test for delirium)

10. For the cohort as a whole, compare the difference in variables listed above (1 to 8) between those who developed postoperative delirium and those who did not, using appropriate statistical test: Chi-squared for proportions, unpaired test-t for continuous parametricallydistributed variables, Mann-Whitney U test for continuous non-parametrically-distributed variables.

Overall study start date 26/01/2018

Completion date 01/02/2019

Eligibility

Key inclusion criteria Patients: 1. Age ≥ 60 2. Able to provide informed consent

Surgical types: 1. Scheduled orthopaedic surgery (GA/sedation +/- regional)

- 2. Scheduled gastrointestinal surgery (open/closed)
- 3. Scheduled robotic urological or gynaecological surgery
- 4. Emergency laparotomy surgery

Participant type(s)

Patient

Age group

Senior

Sex Both

Target number of participants 60

Total final enrolment 60

Key exclusion criteria

- 1. Acute or chronic cognitive impairment, including delirium or dementia
- 2. Previous neurosurgery
- 3. Insufficient English language to discuss participation

Date of first enrolment 03/09/2018

Date of final enrolment 01/01/2019

Locations

Countries of recruitment England

United Kingdom

Study participating centre

Royal Surrey County Hospital Egerton Road Guildford United Kingdom GU2 7XX

Sponsor information

Organisation Royal Surrey County Hospital

Sponsor details Egerton Road Guildford England United Kingdom GU2 7XX 01483 571122 k.penhaligon@nhs.net

Sponsor type Hospital/treatment centre

Website www.royalsurrey.nhs.uk

ROR https://ror.org/02w7x5c08

Funder(s)

Funder type Industry

Funder Name Medtronic

Alternative Name(s) Medtronic Inc.

Funding Body Type Private sector organisation

Funding Body Subtype For-profit companies (industry)

Location United States of America

Results and Publications

Publication and dissemination plan Publish findings at conferences and as a manuscript in a peer-reviewed journal

Intention to publish date 01/07/2019

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Stored in repository

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<u>Results article</u>		01/06/2021	22/03/2021	Yes	No
HRA research summary			28/06/2023	No	No