# CAMELOT - Continuous rectus sheath Analgesia in eMErgency LaparOTomy

<b>Submission date</b> 04/08/2022	<b>Recruitment status</b> Recruiting	[X] Prospectively registered		
		[X] Protocol		
<b>Registration date</b> 09/08/2022	Overall study status Ongoing	Statistical analysis plan		
		Results		
<b>Last Edited</b> 31/10/2025	Condition category Surgery	Individual participant data		
		[X] Record updated in last year		

## Plain English summary of protocol

Background and study aims

In the UK around 30,000 patients, a year undergo an operation called an emergency laparotomy to treat life-threatening conditions. A large vertical cut is made in the abdomen when patients are asleep under general anesthesia. Good pain relief after surgery will help patients feel better and recover quicker. Because of the high level of pain experienced by many patients, opioid-based painkillers such as morphine are often given, using patient-controlled analgesia (PCA) pumps. However, morphine can cause serious side effects such as breathing problems, nausea and vomiting, and delayed bowel movement, which can slow patient recovery. Rectus sheath catheters (RSCs) are a newer way of providing pain relief, where two thin tubes (catheters) are inserted on either side of the wound during the operation. Local anesthetic is injected slowly into the catheters to numb the nerves and reduce pain for about three days. Small studies suggest that RSCs may provide effective pain relief, reduce the use of morphine, and help patient recovery. Potential disadvantages are that RSCs take time to insert and are expensive. More work is needed to understand whether there are any unwanted effects with RSCs.

This study will find out whether adding a RSCs to standard patient-controlled analgesia provides better pain relief, fewer side effects and complications, and greater satisfaction for patients undergoing emergency laparotomy compared to a control group who will receive a sham catheter with no infusion of analgesia. Our study will also aim to determine whether RSCs are safe and cost-effective.

Who can participate?

Adults who are due to undergo emergency laparotomy surgery.

## What does the study involve?

Participants will be allocated to one of two groups, with an equal chance of being in either group (like tossing a coin). An active Rectus Sheath Catheter (RSC) with a constant infusion of local anesthesia for 72 h from the end of surgery will be given to one group of participants. The other group will receive a sham rectus sheath catheter with an inactive infusion device in place for 72 h from the end of surgery. Participants and researchers will not have a choice in which of the two treatments will be given to each participant and will not know which treatment participants have received during the study.

What are the possible benefits and risks of participating?

The benefits of the CAMELOT study are that we hope that patients will have less pain after their operation and that their recovery/discharge time may be quicker.

The risks for participating in the CAMELOT study are associated with the 'active' rectus sheath catheter and are extremely rare. Participants will be actively monitored throughout the study for any possible side effects. Some of the rare risks are described below:

- 1. Disconnection. Occasionally the RSCs may become accidentally disconnected from the pump. If this happens, we will remove the RSCs to prevent infection, and you will be able to have other forms of pain relief.
- 2. Infection. This will be actively monitored by the participant's clinician who will treat accordingly and remove the RSC.
- 3. Bleeding. This will be actively monitored by the participant's clinician who will treat accordingly and remove the RSC.

Where is the study run from?
University Hospital Southampton NHS Foundation Trust (UK)

When is the study starting and how long is it expected to run for? January 2022 to August 2027

Who is funding the study? National Institute for Health and Care Research (NIHR) Health Technology Assessment (HTA) programme (UK)

Who is the main contact?

Jonathan Evans, camelot-trial@bristol.ac.uk

## Contact information

## Type(s)

Public

#### Contact name

Mr Jonathan Evans

#### Contact details

Trial Coordinator - CAMELOT Bristol Trials Centre (BTC) 1-5 Whiteladies Road Clifton Bristol United Kingdom BS8 1NU +44 (0)117 455 1591 camelot-trial@bristol.ac.uk

## Additional identifiers

Clinical Trials Information System (CTIS)

## Integrated Research Application System (IRAS)

312553

## ClinicalTrials.gov (NCT)

Nil known

#### Protocol serial number

IRAS 312553, Sponsor Number: CRI0420, Funder ID: NIHR133554, CPMS 53589

## Study information

#### Scientific Title

CAMELOT - Continuous rectus sheath Analgesia in eMErgency LaparOTomy. A Multi-centre, randomised sham-controlled trial of rectus sheath catheter-delivered local anaesthetic infusion compared with usual care in patients undergoing emergency bowel surgery.

#### Acronym

**CAMELOT** 

### Study objectives

The use of a rectus sheath catheter (RSC)-delivered local anaesthetic infusion in addition to standard analgesia, is superior to standard analgesia without RSC for postoperative pain control.

### Ethics approval required

Old ethics approval format

## Ethics approval(s)

Approved 24/08/2022, London - Bromley Research Ethics Committee (Meeting held by video-conference via Zoom; +44 (0)207 104 8105; bromley.rec@hra.nhs.uk), ref: 22/LO/0555

## Study design

Multi-centre, double-blind, pragmatic, parallel-group, superiority randomized sham-controlled trial with an internal pilot phase to determine feasibility of recruitment and protocol adherence

## Primary study design

Interventional

## Study type(s)

**Treatment** 

## Health condition(s) or problem(s) studied

Use of rectus sheath catheters (RSCs) in patients undergoing emergency bowel surgery.

#### **Interventions**

CAMELOT is a multi-centre, pragmatic, randomised controlled trial in NHS hospitals, with a 12-month internal pilot phase and active follow-up for 6 months post-surgery.

Patients who are undergoing an emergency laparotomy surgery via a midline incision and eligible for inclusion in the National Emergency

Laparotomy Audit (NELA) will be randomised after their midline incision has been performed and before the start of surgical closure, using a secure internet-based randomisation system ensuring allocation concealment. Participants will be allocated into one of two groups via a 1:1 ratio:

- 1. Insertion of an active RSC with infusion of standard local analgesia for 72 hours from the end of surgery
- 2. A sham RSC, placed on the skin of surgical site by adhesive dressing and not delivering any local anaesthetic for 72 hours, from the end of surgery

### Intervention Type

Procedure/Surgery

## Primary outcome(s)

Postoperative pain control measured using the Overall Benefit of Analgesia Score (OBAS) over the first 5 postoperative days

### Key secondary outcome(s))

- 1. Postoperative complications measured using the incidence of the following postoperative complications with severity of Clavien-Dindo grade II or higher collected in the CRF/patient records within 30-days of randomisation:
- 1.1. Postoperative pulmonary complications (PPC)
- 1.2. Respiratory failure
- 1.3. Paralytic ileus
- 1.4. Incisional surgical site infection
- 1.5. Rectus sheath catheter/infusion-related complications
- 2. Time to tracheal extubation (in days) measured using data collected in the CRF/patient records between randomisation and the day of tracheal extubation
- 3. Time to return of bowel function (in days) measured using data collected in the CRF/patient records between randomisation and the day of the return of bowel function
- 4. Time to first mobilisation (in days) measured using data collected in the CRF/patient records between randomisation and the day of first mobilisation
- 5. Pain intensity at rest and on movement measured using data collected in the CRF/patient records on postoperative days 1, 2, 3, 4, and 5
- 6. Postoperative opioid use measured using data collected in the CRF/patient records in the first five days from the end of surgery
- 7. Mortality measured using the incidence of death collected in the CRF/patient records at 30 and 90 days from the date of randomisation
- 8. Chronic postoperative pain measured using the Brief Pain Inventory at 3 and 6 months from the date of randomisation
- 9. Health-related quality of life measured using the EQ5D-5L at 3 days, and 3 and 6 months from the date of randomisation
- 10. Return to work and activity measured using the Work Productivity and Activity Impairment Questionnaire: General Health (WPAI:GH) at 3 and 6 months from the date of randomisation

## Completion date

31/08/2027

## **Eligibility**

### Key inclusion criteria

- 1. Aged ≥18 years
- 2. Undergoing emergency laparotomy surgery via a midline incision
- 3. Eligible for inclusion in the National Emergency Laparotomy Audit (NELA)

### Participant type(s)

Patient

### Healthy volunteers allowed

No

### Age group

Adult

### Lower age limit

18 years

#### Sex

All

### Key exclusion criteria

Current participant exclusion criteria:

- 1. Planned epidural anaesthesia
- 2. Clinician refusal
- 3. Lack of mental capacity to consent to trial participation
- 4. Contraindications to RSC including allergy to local anaesthetic (LA), anatomical factors making RSC insertion impossible
- 5. Existing co-enrolment in another clinical study if:
- 5.1. The intervention in the other study is expected to influence the primary outcome (this will be considered by a senior clinician on a case-by-case basis)
- 5.2. It is considered too burdensome for the patient; or
- 5.3. It is not permitted by the other study
- 6. Previous enrolment in the CAMELOT trial

#### Previous participant exclusion criteria:

- 1. Clinician or patient refusal to participate
- 2. Planned epidural anaesthesia
- 3. Contraindications to Rectus Sheath Catheter (RSC) including allergy to local anaesthetic (LA)
- 4. Anatomical factors making RSC insertion impossible

#### Date of first enrolment

25/01/2023

#### Date of final enrolment

31/07/2026

## Locations

#### Countries of recruitment

United Kingdom

## England

#### Scotland

## Study participating centre North Bristol NHS Trust

Southmead Hospital Southmead Road Westbury-on-trym Bristol United Kingdom BS10 5NB

## Study participating centre Leeds Teaching Hospitals NHS Trust

St. James's University Hospital Beckett Street Leeds United Kingdom LS9 7TF

## Study participating centre Medway NHS Foundation Trust

Medway Maritime Hospital Windmill Road Gillingham United Kingdom ME7 5NY

## Study participating centre Musgrove Park Hospital

Musgrove Park Taunton United Kingdom TA1 5DA

## Study participating centre Royal Alexandra Hospital

Corsebar Road

Paisley United Kingdom PA2 9PN

## Study participating centre Barts Health NHS Trust

The Royal London Hospital 80 Newark Street London United Kingdom E1 2ES

## Study participating centre The Dudley Group NHS Foundation Trust

Russells Hall Hospital Pensnett Road Dudley United Kingdom DY1 2HQ

## Study participating centre Blackpool Teaching Hospitals NHS Foundation Trust

Victoria Hospital Whinney Heys Road Blackpool United Kingdom FY3 8NR

## Study participating centre City Hospitals Sunderland NHS Foundation Trust

Sunderland Royal Hospital Kayll Road Sunderland United Kingdom SR4 7TP

## Study participating centre Great Western Hospitals NHS Foundation Trust

Great Western Hospital Marlborough Road Swindon United Kingdom SN3 6BB

## Study participating centre University Hospitals Birmingham NHS Foundation Trust

Queen Elizabeth Hospital Mindelsohn Way Edgbaston Birmingham United Kingdom B15 2GW

## Study participating centre The Royal Wolverhampton NHS Trust

New Cross Hospital Wolverhampton Road Heath Town Wolverhampton United Kingdom WV10 0QP

## Study participating centre Royal Derby Hospital

Uttoxeter Road Derby United Kingdom DE22 3NE

## Study participating centre Royal Liverpool University Hospital NHS Trust

Royal Liverpool University Hospital Prescot Street Liverpool United Kingdom L7 8XP

## Study participating centre Croydon University Hospital

London Road

Croydon United Kingdom CR7 7YE

## Study participating centre York Hospitals NHS Trust Hq

York Hospital Wigginton Road York United Kingdom YO31 8HE

## Study participating centre Manchester University NHS Foundation Trust

Cobbett House Oxford Road Manchester United Kingdom M13 9WL

## Study participating centre Ayrshire Central Hospital

Kilwinning Road Irvine United Kingdom KA12 8SS

## Study participating centre NHS Greater Glasgow and Clyde

J B Russell House Gartnavel Royal Hospital 1055 Great Western Road Glasgow Glasgow United Kingdom G12 0XH

## Study participating centre South Tees Hospitals NHS Trust Middlesbrough General Hospital

Middlesbrough General Hospital Ayresome Green Lane Middlesbrough

## Study participating centre Heartlands Hospital

Bordesley Green East Bordesley Green Birmingham United Kingdom B9 5ST

## Study participating centre Medway NHS Foundation Trust

Medway Maritime Hospital Windmill Road Gillingham United Kingdom ME7 5NY

## Sponsor information

## Organisation

University Hospital Southampton NHS Foundation Trust

#### **ROR**

https://ror.org/0485axj58

## Funder(s)

## Funder type

Government

#### **Funder Name**

National Institute for Health and Care Research

#### Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

## **Funding Body Type**

#### Government organisation

## **Funding Body Subtype**

National government

#### Location

**United Kingdom** 

#### **Funder Name**

Health Technology Assessment Programme

### Alternative Name(s)

NIHR Health Technology Assessment Programme, Health Technology Assessment (HTA), HTA

#### **Funding Body Type**

Government organisation

## **Funding Body Subtype**

National government

#### Location

United Kingdom

## **Results and Publications**

## Individual participant data (IPD) sharing plan

Data will not be made available for sharing until after publication of the main results of the study unless agreed by the Chief Investigator/Trial Management Group on a case-by-case basis. Thereafter, anonymised individual patient data will be made available for secondary research, conditional on assurance from the secondary researcher that the proposed use of the data is compliant with the MRC Policy on Data Sharing regarding scientific quality, ethical requirements, and value for money. A minimum requirement with respect to scientific quality will be a publicly available pre-specified protocol describing the purpose, methods, and analysis of the secondary research, e.g. a protocol for a Cochrane systematic review. Please contact Mark Edwards using the following email: camelot-trial@bristol.ac.uk.

## IPD sharing plan summary

Available on request

## Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
HRA research summary			28/06/2023	No	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Protocol file	version 1.0	01/07/2022	22/12/2022	No	No
Protocol file	version 2.0	12/04/2023	03/05/2023	No	No