# A clinical trial comparing the effectiveness of a plaster cast to a functional brace in the treatment of adults with ankle fractures

Submission date	Recruitment status  No longer recruiting	[X] Prospectively registered		
17/07/2017		[X] Protocol		
Registration date	Overall study status	Statistical analysis plan		
24/07/2017	Completed	[X] Results		
Last Edited	Condition category	Individual participant data		
06/09/2023	Injury, Occupational Diseases, Poisoning			

## Plain English summary of protocol

Background and study aims

Ankle fractures represent 9% of the trauma workload and demand is increasing. This number is expected to grow due to an increase in older adults who are remaining active. The frequency of this injury is an increasing burden on the NHS year on year. The short-term impact of this injury results in physical impairments of pain, stiffness, weakness and swelling. The longer-term impact results in prolonged time off work, development of posttraumatic arthritis and psychological consequences of depression and anxiety. Ankle fractures that occur below the level of the syndesmosis (the movable joint that connects the two lower leg bones) are considered 'stable' and are usually treated with functional braces. All other ankle fractures are less stable, with some requiring a surgery called open reduction and internal fixation (ORIF) with the aim of restoring stability. Regardless of the decision to operate or not, the immediate management has traditionally been plaster cast for several weeks, while the bone heals. A cast provides maximum support; however, there are potential problems. Firstly, there is the immediate impact on mobility (ability to move) for a period of around six weeks. Secondly, there are the risks associated with not moving the ankle for a long time such as muscle atrophy (wasting), deep vein thrombosis (blood clots forming in a deep vein) and joint stiffness. Finally, there are the long-term consequences, which include prolonged gait (way of walking) abnormalities, persistent calf muscle weakness and an inability to return to previous activity levels. Alternative functional bracing may potentially address these issues. However, it does not provide the same degree of support to the healing bones. The aim of this study is to evaluate if there is a different in ankle function and symptoms between a plaster cast and a functional brace.

## Who can participate?

Adults aged 18 and older who have an ankle fracture that requires a cast and is within three weeks of surgery or injury.

## What does the study involve?

Participants are allocated to one of two groups. Those in the first group receive the functional brace which is worn for three to eight weeks depending on their clinician's advice. They receive an exercise sheet that outlines two simple exercises to help maintain and improve the

movement in their ankle joint. This is done as often as the pain allows. Those in the second group receive a plaster cast which is worn for three to eight weeks depending on their clinician's advice. Participants are followed up for two years and complete surveys via the post and receive their routine follow up by their treating clinician.

What are the possible benefits and risks of participating? There are no notable benefits or risks with participating.

Where is the study run from?
University Hospitals Coventry and Warwickshire (UK)

When is the study starting and how long is it expected to run for? January 2017 to December 2021

Who is funding the study? National Institute for Health Research (UK)

Who is the main contact? Dr Rebecca Mckeown air@warwick.ac.uk

## Contact information

## Type(s)

Public

#### Contact name

Miss Rebecca McKeown

#### Contact details

Warwick Clinical Trials Unit Warwick Medical School Gibbett Hill Road Coventry United Kingdom CV4 7AL +44 2476 575848 air@warwick.ac.uk

## Type(s)

Scientific

#### Contact name

Dr Rebecca Kearney

#### **ORCID ID**

https://orcid.org/0000-0002-8010-164X

#### Contact details

Clinical Trials Unit Gibbett Hill Road Coventry United Kingdom CV7 4AL +44 2476 573156 r.s.kearney@warwick.ac.uk

## Additional identifiers

Protocol serial number 35014

# Study information

#### Scientific Title

Ankle Injury Rehabilitation - A multi-centre randomised controlled trial to assess the difference between plaster cast and functional bracing in the management of ankle fractures

## Acronym

AIR

## Study objectives

The aim of this study is to see if there is a difference in the Olerud Molander Ankle Score at 4 months post randomisation between plaster cast and functional bracing for the treatment of ankle fractures.

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

NRES Committee – West Midlands, Edgbaston, 04/07/2017, ref: 17/LO/1015

## Study design

Randomised; Interventional; Design type: Treatment, Complex Intervention, Physical, Rehabilitation

## Primary study design

Interventional

## Study type(s)

Treatment

## Health condition(s) or problem(s) studied

Ankle fracture

#### Interventions

All adults with a fractured ankle under the care of a clinician at any of the named recruiting sites are potentially eligible. New patients with an ankle fracture are reviewed each day by the trauma team. Participants are randomised to either receiving either the functional brace or the plaster cast. This process of randomisation is done via a computer generated sequence to determine the treatment to be given.

The plaster cast or functional brace is usually worn from three to eight weeks, which is decided by the treating clinician. The clinician also decides how much weight can be put on the healing ankle.

Participants who receive the functional brace are also be given an exercise sheet, outlining two very simple exercises to maintain and improve the movement in the ankle joint. It is recommended these are completed little and often, as pain allows.

Participants are followed up in this study for a duration of two years and all trial follow up is completed by post. Participants will receive routine follow up by their treating clinician.

#### Intervention Type

Other

#### **Phase**

Phase III

## Primary outcome(s)

Symptom evaluation is measured using Olerud Molander Ankle Score questionnaire (this will be collected using patient questionnaires sent via post) at 4 months

## Key secondary outcome(s))

- 1. Symptom evaluation is measured using Olerud Molander Ankle Score questionnaire at six weeks, 10 weeks, six months and 24 months
- 2. Surgical outcomes are measured using the Manchester-Oxford Foot and Ankle Questionnaire at four months
- 3. Health status is measured using EuroQol five dimensions questionnaire (EQ-5D) at "6 weeks, 10 weeks, 16 weeks, 24 weeks, 12 months, 18 months and 24 months.
- 4. Disability is measured using the disability rating index at "6 weeks, 10 weeks, 16 weeks, 24 weeks, 12 months, 18 months and 24 months
- 5. Cost is measured using a patient reported questionnaire at "6 weeks, 10 weeks, 16 weeks, 24 weeks, 12 months, 18 months and 24 months
- 6. Complication rate is measured using a patient reported questionnaire at "6 weeks, 10 weeks, 16 weeks, 24 weeks, 12 months, 18 months and 24 months

## Completion date

31/12/2021

# **Eligibility**

## Key inclusion criteria

- 1. Provision of written informed consent
- 2. Aged 18 years or over
- 3. A closed ankle fracture for which the treating clinician would consider plaster cast a reasonable management option
- 4. Within 3 weeks of operative treatment or injury if non-operatively managed

## Participant type(s)

Patient

## Healthy volunteers allowed

No

## Age group

Adult

#### Lower age limit

18 years

#### Sex

All

#### Total final enrolment

671

#### Key exclusion criteria

- 1. Ankle fracture secondary to known metastatic disease
- 2. Complex intra-articular fracture (e.g. Pilon fracture)
- 3. In the opinion of the surgeon the patient would require manipulation and close contact casting
- 4. In the opinion of the surgeon the patient would require manipulation and moulded cast
- 5. Wound complications following surgical management contraindicating Functional brace intervention
- 6. Previous ankle fracture randomised in the present trial
- 7. Evidence that the patient would be unable to adhere to trial procedures or complete postal questionnaires
- 8. Known pre-existing neuropathic joint disease contraindicating functional brace intervention

#### Date of first enrolment

01/10/2017

#### Date of final enrolment

30/09/2019

## Locations

#### Countries of recruitment

**United Kingdom** 

England

## Study participating centre University Hospitals Coventry and Warwickshire

Clifford Bridge Road Coventry United Kingdom CV2 2DX

# Sponsor information

#### Organisation

University of Warwick

#### **ROR**

https://ror.org/01a77tt86

# Funder(s)

## Funder type

Government

#### **Funder Name**

National Institute for Health Research

## Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

## **Funding Body Type**

Government organisation

## Funding Body Subtype

National government

#### Location

United Kingdom

## **Results and Publications**

## Individual participant data (IPD) sharing plan

The data sharing plans for the current study are unknown and will be made available at a later date

#### IPD sharing plan summary

Data sharing statement to be made available at a later date

## **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		05/07/2021	07/07/2021	Yes	No
Results article		06/07/2021	30/07/2021	Yes	No
Protocol article	protocol	18/12/2018	04/11/2019	Yes	No

Other publications	Qualitative pre-results	04/02/2020	06/09/2023 Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025 No	Yes
Study website	Study website	11/11/2025	11/11/2025 No	Yes