

# A clinical trial comparing the effectiveness of a plaster cast to a functional brace in the treatment of adults with ankle fractures

<b>Submission date</b> 17/07/2017	<b>Recruitment status</b> No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
<b>Registration date</b> 24/07/2017	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 06/09/2023	<b>Condition category</b> Injury, Occupational Diseases, Poisoning	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

Ankle fractures represent 9% of the trauma workload and demand is increasing. This number is expected to grow due to an increase in older adults who are remaining active. The frequency of this injury is an increasing burden on the NHS year on year. The short-term impact of this injury results in physical impairments of pain, stiffness, weakness and swelling. The longer-term impact results in prolonged time off work, development of posttraumatic arthritis and psychological consequences of depression and anxiety. Ankle fractures that occur below the level of the syndesmosis (the movable joint that connects the two lower leg bones) are considered 'stable' and are usually treated with functional braces. All other ankle fractures are less stable, with some requiring a surgery called open reduction and internal fixation (ORIF) with the aim of restoring stability. Regardless of the decision to operate or not, the immediate management has traditionally been plaster cast for several weeks, while the bone heals. A cast provides maximum support; however, there are potential problems. Firstly, there is the immediate impact on mobility (ability to move) for a period of around six weeks. Secondly, there are the risks associated with not moving the ankle for a long time such as muscle atrophy (wasting), deep vein thrombosis (blood clots forming in a deep vein) and joint stiffness. Finally, there are the long-term consequences, which include prolonged gait (way of walking) abnormalities, persistent calf muscle weakness and an inability to return to previous activity levels. Alternative functional bracing may potentially address these issues. However, it does not provide the same degree of support to the healing bones. The aim of this study is to evaluate if there is a different in ankle function and symptoms between a plaster cast and a functional brace.

### Who can participate?

Adults aged 18 and older who have an ankle fracture that requires a cast and is within three weeks of surgery or injury.

### What does the study involve?

Participants are allocated to one of two groups. Those in the first group receive the functional brace which is worn for three to eight weeks depending on their clinician's advice. They receive an exercise sheet that outlines two simple exercises to help maintain and improve the

movement in their ankle joint. This is done as often as the pain allows. Those in the second group receive a plaster cast which is worn for three to eight weeks depending on their clinician's advice. Participants are followed up for two years and complete surveys via the post and receive their routine follow up by their treating clinician.

What are the possible benefits and risks of participating?  
There are no notable benefits or risks with participating.

Where is the study run from?  
University Hospitals Coventry and Warwickshire (UK)

When is the study starting and how long is it expected to run for?  
January 2017 to December 2021

Who is funding the study?  
National Institute for Health Research (UK)

Who is the main contact?  
Dr Rebecca Mckeown  
air@warwick.ac.uk

**Study website**  
<http://www2.warwick.ac.uk/fac/med/research/hscience/ctu/trials/airmain>

## Contact information

**Type(s)**  
Public

**Contact name**  
Miss Rebecca McKeown

**Contact details**  
Warwick Clinical Trials Unit  
Warwick Medical School  
Gibbett Hill Road  
Coventry  
United Kingdom  
CV4 7AL  
+44 2476 575848  
air@warwick.ac.uk

**Type(s)**  
Scientific

**Contact name**  
Dr Rebecca Kearney

**ORCID ID**  
<https://orcid.org/0000-0002-8010-164X>

**Contact details**

Clinical Trials Unit  
Gibbett Hill Road  
Coventry  
United Kingdom  
CV7 4AL  
+44 2476 573156  
r.s.kearney@warwick.ac.uk

**Additional identifiers**

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

35014

**Study information****Scientific Title**

Ankle Injury Rehabilitation - A multi-centre randomised controlled trial to assess the difference between plaster cast and functional bracing in the management of ankle fractures

**Acronym**

AIR

**Study objectives**

The aim of this study is to see if there is a difference in the Olerud Molander Ankle Score at 4 months post randomisation between plaster cast and functional bracing for the treatment of ankle fractures.

**Ethics approval required**

Old ethics approval format

**Ethics approval(s)**

NRES Committee – West Midlands, Edgbaston, 04/07/2017, ref: 17/LO/1015

**Study design**

Randomised; Interventional; Design type: Treatment, Complex Intervention, Physical, Rehabilitation

**Primary study design**

Interventional

**Secondary study design**

Randomised controlled trial

**Study setting(s)**

Hospital

**Study type(s)**

Treatment

**Participant information sheet**

Not available in web format, please use the contact details below to request a patient information sheet

**Health condition(s) or problem(s) studied**

Ankle fracture

**Interventions**

All adults with a fractured ankle under the care of a clinician at any of the named recruiting sites are potentially eligible. New patients with an ankle fracture are reviewed each day by the trauma team. Participants are randomised to either receiving either the functional brace or the plaster cast. This process of randomisation is done via a computer generated sequence to determine the treatment to be given.

The plaster cast or functional brace is usually worn from three to eight weeks, which is decided by the treating clinician. The clinician also decides how much weight can be put on the healing ankle.

Participants who receive the functional brace are also be given an exercise sheet, outlining two very simple exercises to maintain and improve the movement in the ankle joint. It is recommended these are completed little and often, as pain allows.

Participants are followed up in this study for a duration of two years and all trial follow up is completed by post. Participants will receive routine follow up by their treating clinician.

**Intervention Type**

Other

**Phase**

Phase III

**Primary outcome measure**

Symptom evaluation is measured using Olerud Molander Ankle Score questionnaire (this will be collected using patient questionnaires sent via post) at 4 months

**Secondary outcome measures**

1. Symptom evaluation is measured using Olerud Molander Ankle Score questionnaire at six weeks, 10 weeks, six months and 24 months
2. Surgical outcomes are measured using the Manchester-Oxford Foot and Ankle Questionnaire at four months
3. Health status is measured using EuroQol five dimensions questionnaire (EQ-5D) at "6 weeks, 10 weeks, 16 weeks, 24 weeks, 12 months, 18 months and 24 months.
4. Disability is measured using the disability rating index at "6 weeks, 10 weeks, 16 weeks, 24 weeks, 12 months, 18 months and 24 months
5. Cost is measured using a patient reported questionnaire at "6 weeks, 10 weeks, 16 weeks, 24

weeks, 12 months, 18 months and 24 months

6. Complication rate is measured using a patient reported questionnaire at "6 weeks, 10 weeks, 16 weeks, 24 weeks, 12 months, 18 months and 24 months

**Overall study start date**

01/01/2017

**Completion date**

31/12/2021

## **Eligibility**

**Key inclusion criteria**

1. Provision of written informed consent
2. Aged 18 years or over
3. A closed ankle fracture for which the treating clinician would consider plaster cast a reasonable management option
4. Within 3 weeks of operative treatment or injury if non-operatively managed

**Participant type(s)**

Patient

**Age group**

Adult

**Lower age limit**

18 Years

**Sex**

Both

**Target number of participants**

Planned Sample Size: a minimum of 478; UK Sample Size: a minimum of 478

**Total final enrolment**

671

**Key exclusion criteria**

1. Ankle fracture secondary to known metastatic disease
2. Complex intra-articular fracture (e.g. Pilon fracture)
3. In the opinion of the surgeon the patient would require manipulation and close contact casting
4. In the opinion of the surgeon the patient would require manipulation and moulded cast
5. Wound complications following surgical management contraindicating Functional brace intervention
6. Previous ankle fracture randomised in the present trial
7. Evidence that the patient would be unable to adhere to trial procedures or complete postal questionnaires
8. Known pre-existing neuropathic joint disease contraindicating functional brace intervention

**Date of first enrolment**

01/10/2017

**Date of final enrolment**

30/09/2019

## **Locations**

**Countries of recruitment**

England

United Kingdom

**Study participating centre**

**University Hospitals Coventry and Warwickshire**

Clifford Bridge Road

Coventry

United Kingdom

CV2 2DX

## **Sponsor information**

**Organisation**

University of Warwick

**Sponsor details**

Sponsorship Office

Gibbett Hill Road

Coventry

England

United Kingdom

CV4 7AL

**Sponsor type**

University/education

**ROR**

<https://ror.org/01a77tt86>

## **Funder(s)**

**Funder type**

Government

**Funder Name**

National Institute for Health Research

**Alternative Name(s)**

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

**Funding Body Type**

Government organisation

**Funding Body Subtype**

National government

**Location**

United Kingdom

## Results and Publications

**Publication and dissemination plan**

Planned publication in a high-impact peer reviewed journal no later than 12 months after the end of the follow up period.

**Intention to publish date**

31/12/2022

**Individual participant data (IPD) sharing plan**

The data sharing plans for the current study are unknown and will be made available at a later date

**IPD sharing plan summary**

Data sharing statement to be made available at a later date

**Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Protocol article</a>	protocol	18/12/2018	04/11/2019	Yes	No
<a href="#">Results article</a>	Qualitative pre-results	05/07/2021	07/07/2021	Yes	No
<a href="#">Results article</a>		06/07/2021	30/07/2021	Yes	No
<a href="#">Other publications</a>		04/02/2020	06/09/2023	Yes	No