Can Clovinol (clove extract) lower blood sugar levels after meals in healthy volunteers?

Submission date	Recruitment status No longer recruiting	Prospectively registered			
21/09/2018		☐ Protocol			
Registration date	Overall study status Completed	Statistical analysis plan			
28/09/2018		[X] Results			
Last Edited	Condition category	Individual participant data			
09/05/2019	Nutritional, Metabolic, Endocrine				

Plain English summary of protocol

Background and study aims

The main energy source the body uses is sugar (i.e. glucose), which is derived from the food we eat. The amount of sugar in our blood (i.e. blood glucose levels) is an important indicator of overall health. Both too much sugar (hyperglycemia) and too little sugar (hypoglycemia) in the blood is linked with negative effects on health. Fasting blood sugar is measured when someone has not eaten or drunk anything (except water) for at least 8 hours. Normal fasting blood sugar levels should be below 100 mg/dL. Prediabetes is characterized by fasting blood sugar levels between 100 to 125 mg/dL, and diabetes is when fasting blood sugar is greater than 125 mg/dL. When we eat, our blood sugar increases, and then within 2 to 3 hours blood sugar should return to the level it was before a meal. The amount of sugar in our blood after we eat is called the postprandial blood sugar level. This measurement of blood sugar is also an important indicator of overall health and can be an early sign of abnormal blood sugar control. Previous studies have identified that clove extract can play a role in helping to manage blood glucose levels. The main objective of this study is to test whether a specialized extract of clove bud, Clovinol®, can help control postprandial blood sugar.

Who can participate?

Individuals who are between 25 and 35 years old, not taking any medication, not obese, and have preprandial (before eating) blood sugar levels below 125 mg/dL.

What does the study involve?

Participants will need to maintain their typical South Indian diet and complete their breakfast by 8 am, eat lunch at 12 pm, and take a Clovinol® capsule immediately after lunch throughout this 30-day study. Between breakfast and lunch, participants should have no snacks or drinks (except water). On specified days of the study, blood sugar will be measured using a digital glucometer immediately before lunch and 2 hours after the start of their lunch.

What are the possible benefits and risks of participating?

Potential benefits include reductions in blood glucose after a meal, which can possibly reduce post-meal fatigue. Potential side effects are limited, but may include mild stomach discomfort.

Where is the study run, and who is funding the study? This study is funded by and will be conducted at the R&D Center of Akay Flavours & Aromatics Ltd. in Kerala, India.

When is the study starting and how long is it expected to run for? The study is expected to start in September of 2014 and continue for 30 days.

Who is the main contact?

The main contact is Dr. Krishnakumar IM, Krishnakumar.im@akay-group.com.

Contact information

Type(s)

Public

Contact name

Dr IM Krishnakumar

ORCID ID

https://orcid.org/0000-0003-0594-7650

Contact details

Malaidamthuruthu P. O. Ernakulam Cochin India 683561

Additional identifiers

Protocol serial number

AKCN-CLS_01/14

Study information

Scientific Title

Effects of Clovinol on random blood sugar in healthy and pre-diabetic adults - an open-label pilot study

Study objectives

Type 2 diabetes (T2D) is a global pandemic, and contributes significantly to the increasing incidence of other conditions such as cardiovascular disease. While diabetes screening typically relies on assessment of fasting blood glucose, this may not detect subtle early manifestations of abnormal glucose metabolism. One such early manifestation is impaired glucose tolerance which can precede T2D by many years and is a risk factor for all-cause mortality. Postprandial plasma glucose measured 2 hours after the start of a meal is a good indicator of the overall status of glucose metabolism. Clove (Syzygium aromaticum L.) and its essential oils (eugenol and acetyl eugenol) have been shown in preclinical studies to modulate pathways involved in glucose metabolism. The objective of this pilot study is to test whether Clovinol could influence glucose metabolism. We will evaluate the effect of Clovinol® supplementation on preprandial glucose

levels and 2-h PPG levels in otherwise healthy volunteers who have preprandial glucose levels less than 125 mg/dL. This study will test the hypothesis that daily supplementation with Clovinol will reduce pre- and post-prandial blood glucose by 5 and 40 mg/dL, respectively.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Sri Rama Hospital ethics committee, 08/08/2014, ECR/184/Indt/KA/2014

Study design

Single-center open-label interventional pilot study

Primary study design

Interventional

Study type(s)

Prevention

Health condition(s) or problem(s) studied

Prediabetes

Interventions

250 mg Clovinol, a hydroethanolic water-soluble exract of unopened clove bud (Syzygium aromaticum L.), administered in one gelatin capsule once daily immediately after lunch for 30 days

Intervention Type

Supplement

Primary outcome(s)

Blood glucose measured with a digital glucometer 2 hours after lunch (post-prandial) measured on days 1, 12, 24 and 30. Day 1 is without Clovinol.

Key secondary outcome(s))

Blood glucose measured with a digital glucometer immediately before lunch (pre-prandial) measured on days 1, 12, 24 and 30. Day 1 is without Clovinol.

Completion date

23/10/2014

Eligibility

Key inclusion criteria

- 1. Healthy
- 2. Non-obese
- 3. Aged 25 35 years
- 4. Blood glucose below 125 mg/dL

Participant type(s)

Healthy volunteer

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Total final enrolment

13

Key exclusion criteria

- 1. Take any medication
- 2. Obese
- 3. Have diabetes

Date of first enrolment

05/09/2014

Date of final enrolment

20/09/2014

Locations

Countries of recruitment

India

Study participating centre

R&D Center, Akay Flavours & Aromatics Pvt Ltd

Ambunadu, Malaidamthirith P O Ernakulum India 683561

Sponsor information

Organisation

Akay Flavours & Aromatics Pvt. Ltd.

Funder(s)

Funder type

Not defined

Funder Name

Investigator initiated and funded

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated and/or analysed during the current study during this study will be included in the subsequent results publication

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	07/05/2019	09/05/2019	Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes