Retraining automatic action tendencies for smoking using mobile phone-based approachavoidance bias training

Submission date 20/11/2018	Recruitment status No longer recruiting	[X] Prospectively registered	
		[X] Protocol	
Registration date	Overall study status	[] Statistical analysis plan	
24/01/2019 Last Edited	Completed Condition category	[_] Results	
		Individual participant data	
09/12/2020	Mental and Behavioural Disorders	[_] Record updated in last year	

Plain English summary of protocol

Background and study aims

Tobacco smoking remains a major public health problem. The World Health Organization (WHO) states that smoking cigarettes is the largest preventable risk factor for morbidity and mortality worldwide, with smoking-related deaths being estimated at over five million people per year. Even following regular, gold-standard smoking cessation interventions, only about 25 percent will achieve sustained abstinence. This might be the case because contemporary interventions often fail to target automatic processes associated with smoking. One such process involves automatic tendencies to approach smoking-related stimuli that have been implicated in smoking intensity. The Approach-Avoidance-Task (AAT) has proven valuable in both measuring and modifying smoking-related approach tendencies. During the training version of the task, smokers are instructed to make an avoidance movement in response to all smoking-related images using a joystick attached to the computer. Although recent literature shows that the AAT-training has merit in modifying smoking-related approach biases or reducing nicotine consumption, accessibility of the training is impaired by the fact that trainings are carried out in the lab on desktop PCs. To facilitate access to the training and to increase training efficiency, the present study provides mobile phone-based AAT training (app-AAT), which can be completed outside the laboratory context (i.e., at home). The main goal is to investigate the effectiveness of the app-AAT in reducing smoking behavior or facilitating abstinence. The results of this study can inform future research in the optimization and advancement of treatment for nicotine addiction.

Who can participate?

People who have smoked at least six cigarettes per day for at least six months, are active smokers at study entry and are motivated to quit smoking

What does the study involve?

Participants take part in a brief behavioral support for smoking. Afterwards, smokers randomly allocated to the app-AAT group are given access to the mobile-phone based training. Participants in this group are instructed to perform the training on a daily basis. Training effects on automatic approach tendencies and smoking behavior are measured immediately after

training and at a four-week follow-up. Participants who do not receive the training (waitlist control group) are then given access to the training app.

What are the possible benefits and risks of participating? A possible benefit is that the app-AAT may help smokers to reduce or stop smoking. There are no significant risks of participating in the study.

Where is the study run from? University of Siegen (Germany)

When is the study starting and how long is it expected to run for? July 2018 to May 2020

Who is funding the study? Ministerium für Kultur und Wissenschaft, NRW (Germany)

Who is the main contact? Dr Alla Machulska

Contact information

Type(s) Scientific

Contact name Dr Alla Machulska

Contact details Adolf-Reichwein-Str. 2a Siegen Germany 57076

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers N/A

Study information

Scientific Title

Retraining automatic action tendencies for smoking using mobile phone-based approachavoidance bias training: a randomized controlled study

Study objectives

It is expected that participants in the intervention condition will show reduced smoking behavior, increased abstinence rates and diminished cognitive biases than participants in the control condition. Close (diminished approach bias for untrained pictures in the assessment version of the AAT) and broad (diminished attentional and association biases) generalization of training is explored, as well therapy effects on clinical outcomes such as smoking behavior, attitudes towards smoking and motivation to quit smoking.

It is expected that the training intervention will decrease or reverse cognitive biases and that these changes can possibly mediate the effects on the clinical outcome.

Ethics approval required

Old ethics approval format

Ethics approval(s) Ethics Committee of the University of Siegen

Study design Single-centre randomised trial, 2(condition)x3(time) factorial design

Primary study design Interventional

Secondary study design Randomised controlled trial

Study setting(s) Other

Study type(s) Treatment

Participant information sheet

Not available in web format, please use contact details to request a patient information sheet

Health condition(s) or problem(s) studied

Nicotine addiction

Interventions

Participants will be randomly assigned to the experimental or control group, according to an externally constructed randomization plan.

1. TAU (brief behavioral support for smoking) + app-based approach bias training 2. TAU

Research Flow for the treatment and control group:

1. Recruitment

2. Brief behavioral support for smoking (TAU) + Baseline assessment (First lab session): After the first session, participants are randomly allocated to one of two conditions

3. Training: app-AAT training at home for 14 consecutive days

4. Post-test (Second lab session): post-test two weeks after the baseline assessment.

5. Follow-up (Third lab session): after 4 weeks.

Prior to randomization, participants receive psychoeducation containing information on nicotine addiction and maintenance and short and long-term effects associated with cigarette smoking (TAU; about 60 minutes). Afterwards, smokers will be handed over a self-help book ("The Easy Way to Stop Smoking" by Allen Carr) to aid smoking cessation. Participants allocated to the experimental condition are then given access to the AAT-training app and are instructed to train at least once per day for 14 consecutive days. Participants perform a practice training with the researchers to ensure that the training handling and concepts have been fully understood. To ensure compliance, participants will receive daily reminders via short message service (SMS) to complete the training. After competition of the study (at the end of the four-week follow-up session), the waitlist-control condition will receive access to the training app.

Intervention Type

Behavioural

Primary outcome measure

Measured at baseline, post-training and at a 4-week follow-up:

1. Approach bias measured by the assessment version of the Approach-Avoidance Task

2. Self-reported nicotine consumption measured via a questionnaire

follow-up. In addition, participants will be instructed to log smoked cigarettes when smoking by using a smartphone-based cigarette-tracking app. The logging period will cover 6 weeks, which is the study duration for each participant

3. Abstinence from smoking determined through self-report and biochemically-verified expired CO (Smokerlyzer)

Secondary outcome measures

Measured at baseline, post-training and at a 4-week follow-up:

1. Cognitive biases measured by a visual dot-probe task and the Implicit Association Task 2. Smoking behavior assessed using Fagerström Test for Nicotine Dependence (FTND; Heatherton et al., 1991)

3. Motivation to stop smoking assessed using Stages of Change Scale; Prochaska et al., 1991; Thoughts About Abstinence Scale; Hall et al., 1990

4. Cigarette craving assessed using single item craving rating ranging from: 0 = not at all, to 5 = very much

5. Explicit attitudes toward smoking assessed using semantic differential based on Swanson et al., 2001

Overall study start date

01/07/2018

Completion date

31/05/2020

Eligibility

Key inclusion criteria

Smokers are included if they had smoked for at least six months at least six cigarettes per day, are active smokers at study entry and are motivated to quit smoking.

Participant type(s)

Healthy volunteer

Age group

Adult

Sex Both

Target number of participants 100

Key exclusion criteria

Exclusion criteria are current alcohol or drug misuse or dependency, psychiatric illness, insufficient German language skills or uncorrected visual or auditory impairment.

Date of first enrolment 01/03/2019

Date of final enrolment 01/03/2020

Locations

Countries of recruitment Germany

Study participating centre University of Siegen Adolf-Reichwein-Str. 2a Siegen Germany 57076

Sponsor information

Organisation University of Siegen

Sponsor details Department of Clinical Psychology Adolf-Reichwein-Str. 2a Siegen Germany 57076

Sponsor type

University/education

Website https://www.uni-siegen.de/start/

ROR https://ror.org/02azyry73

Funder(s)

Funder type Government

Funder Name Ministerium für Kultur und Wissenschaft, NRW, Germany

Results and Publications

Publication and dissemination plan

Planned publication of the study protocol in Trials, intention to publish date: 01/02/2019 Planned publication of the study results in a high-impact peer reviewed journal, intention to publish date: 01/03/2021

Intention to publish date

01/03/2021

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from Dr Alla Machulska (Alla.machulska@uni-siegen.de).

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Protocol article	protocol	12/12/2019	09/12/2020	Yes	No