Evaluation of home rehabilitation for hip fracture patients

| Submission date | Recruitment status No longer recruiting | Prospectively registered | | |
|-------------------------------------|---|-----------------------------|--|--|
| 16/06/2008 | | ☐ Protocol | | |
| Registration date 09/10/2008 | Overall study status Completed | Statistical analysis plan | | |
| | | [X] Results | | |
| Last Edited 10/03/2023 | Condition category Injury, Occupational Diseases, Poisoning | Individual participant data | | |

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Prof Yngve Gustafson

Contact details

Umeå University Geriatric Medicine Umeå Sweden SE-901 87

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers N/A

Study information

Scientific Title

Home rehabilitation for older people with hip fracture - a randomised controlled trial

Study objectives

Multidisciplinary home rehabilitation is applicable and effective as conventional rehabilitation at hospital.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics Committee of the Faculty of Medicine at Umea University. Date of approval 01/04/2008 (ref: Dnr 08-053M)

Study design

Randomised controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Not specified

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Rehabilitation for hip fracture patients

Interventions

Multidisciplinary home rehabilitation compared to conventional multidisciplinary rehabilitation at hospital with follow-ups at 12 weeks after randomisation and 12 months after surgery.

Those randomised to control group will be at hospital as long as necessary and those randomised to intervention/experimental group have home rehabilitation as long as necessary but not longer than 10 weeks.

Both groups will have comprehensive geriatric assessments, management and rehabilitation with active prevention, detection and treatment of postoperative complications such as falls, delirium, pain and infections, for example. Those randomised to home rehabilitation will receive some of these assessments and rehabilitation at home.

Intervention Type

Other

Phase

Not Specified

Primary outcome measure

- 1. In-patient days and re-admissions, followed-up until 12 months post-surgery
- 2. Complications, for example falls and injuries, followed-up until 12 months post-surgery
- 3. Depression, assessed by the Geriatric Depression Scale (GDS) followed-up until 12 months post-surgery
- 4. Delirium, assessed by the Organic Brain Syndrome Scale followed-up until 12 months postsurgery
- 5. Functional performance regarding activities of daily living (ADL), assessed by the Katz ADL index and Barthel ADL index, at 12 weeks post-randomisation and 12 months post-surgery 6. Walking ability and use of walking aids, assessed by the Swedish version of the Clinical Outcome Variable Scale (S-COVS) at 12 weeks post-randomisation and 12 months post-surgery

Secondary outcome measures

The following will be assessed at 12 weeks post-randomisation and 12 months post-surgery:

- 1. Balance, assessed by the Bergs Balance Scale and Fear of Falling
- 2. Quality of life, assessed by EQ-5D
- 3. Nutritional status, assessed by the mini nutritional assessment (MNA)
- 4. Costs
- 5. Self-perceived health, assessed by the 36-item Short Form (SF-36) health survey
- 6. Pain, assessed by a visual analogue scale (VAS)
- 7. Subjective well-being, assessed by the Philadelphia Geriatric Morale Scale (PGCM)

Overall study start date

01/08/2008

Completion date

01/08/2010

Eligibility

Kev inclusion criteria

- 1. Both males and females, 70 years and above
- 2. Hip fracture patients after surgery
- 3. Those who live in the municipality of Umeå (both independent living and from institutions)

Participant type(s)

Patient

Age group

Senior

Sex

Both

Target number of participants

200

Total final enrolment 205

Key exclusion criteria

No exclusion criteria

Date of first enrolment

01/08/2008

Date of final enrolment

01/08/2010

Locations

Countries of recruitment

Sweden

Study participating centre Umeå University

Umeå Sweden SE-901 87

Sponsor information

Organisation

Umeå University (Sweden)

Sponsor details

_

Umeå Sweden SE-901 87

Sponsor type

University/education

Website

http://www.umu.se/umu/index_eng.html

ROR

https://ror.org/05kb8h459

Funder(s)

Funder type

University/education

Funder Name

University of Umeå (Sweden)

Funder Name

County Council of Västerbotten (Sweden)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

Not provided at time of registration

IPD sharing plan summary

Not provided at time of registration

Study outputs

| Output type | Details | Date created | Date added | Peer reviewed? | Patient-facing? |
|-----------------|---------|--------------|------------|----------------|-----------------|
| Results article | | 04/09/2020 | 10/03/2023 | Yes | No |
| Results article | | 09/03/2023 | 10/03/2023 | Yes | No |