

Is a new gum replacement material from a pig as effective as a person's own gum taken from the palate placed around dental implants?

Submission date 25/09/2019	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 22/01/2020	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 22/01/2020	Condition category Oral Health	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Dental implants offer a reliable treatment in the replacement of missing teeth. A well-integrated titanium implant serves as an artificial root into which a prosthetic crown can then be fitted. The successful integration of an implant depends, among other things, on healthy surrounding soft tissues or mucosa. The oral mucosa, lining the mouth can be classed as either keratinised or non-keratinised. Keratinised mucosa is more resilient and provides a stable cuff of tissue around the dental implant. This helps the patient to maintain good oral hygiene and creates an aesthetic profile that mirrors the natural tooth-tissue relationship. The soft tissue surrounding the implant may become inflamed as a result of poor and/or painful brushing and plaque accumulation. This condition is known as peri-implant mucositis. If left untreated, it could cause inflammatory bone loss around the implants called peri-implantitis and may ultimately lead to the premature failure of the implant.

Studies have shown that surgical reconstruction of the peri-implant keratinised mucosa (PIKM) can prevent disease progression. In order to reconstruct the missing PIKM, various materials can be applied such as patient's own gum tissue from palate (autograft) or graft material from another species (xenograft). To date autograft has yielded the most reliable results and serves as "gold standard". However, the harvesting of a patient's own tissue can cause significant pain and discomfort. Further drawbacks include the limited amount of tissue available for harvesting and a compromised colour match at the recipient site as compared to the neighbouring soft tissue. Such disadvantages have led to an increased need for xenografts.

A recently developed xenogenic material derived from a pig tissue called mucoderm® has shown promising results in increasing PIKM width. The aim of this study is to test whether the mucoderm® works as successful as the own gum tissue to improve the width of PIKM. In addition, we compare the treatment time, pain, discomfort after the surgeries.

Who can participate?

Dental implant patients with insufficient gum width at the implant area

What does the study involve?

Participants will be randomly assigned to receive treatment as usual or treatment using the mucoderm® material.

What are the possible benefits and risks of participating?

Subjects would benefit from the increased PIKM as described to be a prerequisite for implant longevity. Side effects to include general post-operative surgical site effects such as pain, discomfort, minor bleeding and swelling (as usual side effects in such “out of study” interventions). In addition, the main aim of the study is to test a material and technique which may reduce these side effects.

Where is the study run from?

Department of Periodontology, Faculty of Dentistry, Semmelweis University, Hungary

When is the study starting and how long is it expected to run for?

October 2018 to December 2020

Who is funding the study?

Department of Periodontology, Faculty of Dentistry, Semmelweis University, Hungary

Who is the main contact?

Dr Attila Horvath

attila.horvath.dr@hotmail.com

Contact information

Type(s)

Scientific

Contact name

Dr Attila Horvath

ORCID ID

<http://orcid.org/0000-0002-0472-4060>

Contact details

Szentkiralyi u. 47.

Budapest

Hungary

1088

+36 1 4591500 ext. 59188

horvath.attila@dent.semmelweis-univ.hu

Additional identifiers

EudraCT/CTIS number

Nil known

IRAS number

ClinicalTrials.gov number

Nil known

Secondary identifying numbers

P-2/2017

Study information

Scientific Title

Comparison of epithelized connective tissue graft and porcine dermal matrix (mucoderm®) to increase the amount of peri-implant soft tissue. Randomised controlled clinical trial

Acronym

PIKM increase

Study objectives

New porcine-derived xenograft (mucoderm®) may be equally effective as the gold standard epithelized connective tissue autograft to increase peri-implant keratinised mucosa (PIKM) with the reduction of the inherent risks and side effects of autograft harvest

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 13/11/2019, Semmelweis University Regional and Institutional Committee of Science and Research Ethics (Üllői út 93, Budapest 1091, Hungary; +36 2157300 ext. 53513; sotonyi.peter@med.semmelweis-univ.hu), ref: 223/2017

Study design

Single centre interventional randomised single masked controlled clinical trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use contact details to request a participant information sheet

Health condition(s) or problem(s) studied

Peri-implant mucositis, peri-implantitis

Interventions

Subjects were divided into control and test groups for both the upper and lower jaws. The total duration of the study is one year.

Subject Groups:

1. Upper jaw test group (Group 1): PIKM augmented with mucoderm®
2. Upper jaw control (Group 2) group: PIKM augmented with ECTG
3. Lower jaw test group (Group 3): PIKM augmented with a combination of mucoderm® and ECTG strip
4. Lower jaw control group (Group 4): PIKM augmented with ECTG

Treatment Summary:

Group1:

A split-thickness flap is elevated apically at the mucogingival junction (MGJ) and anchored to the periosteum with continuous sutures. The recipient site is exposed and the mucoderm® is trimmed and sutured to the underlying periosteum using two different suture techniques (a mixture of internal and external horizontal mattress sutures) to ensure tight contact and avoid blood clot formation between the recipient site and the mucoderm®.

Group 2:

Same as in group 1 but instead of xenograft, an ECTG autograft is harvested from the hard palate (donor site) which is restored using a collagen matrix (lyostypt®) and stabilized by continuous sutures.

Group 3:

Same as in Group 1, but apically to the mucoderm® a narrow ECTG strip (2mm width) is sutured to the periosteum.

Group 4:

Same as in group 2

Details of randomization process:

Block randomization technique

The control group and test group are assigned the letters T (21 test) and C (12 control), respectively. These 24 letters (papers) will be placed in opaque envelopes and sealed right after. All these sealed envelopes will be placed in an opaque box. On the day of each surgery a third party (independent of study) pulls blindly and opens one sealed opaque envelop. Therefore, the random assignment will be revealed.

The same randomization method will be used in both jaw bones, respectively

Intervention Type

Procedure/Surgery

Primary outcome measure

PIKM width (PIKM-W) in mm, measured at baseline, 1, 3, 6, 9, 12 months

Secondary outcome measures

1. PIKM thickness (PIKM-T) in mm, measured at baseline, 1, 3, 6, 9, 12 months
2. Pocket Probing Depths (PPD) measured at baseline, 3, 6, 9, 12 months
3. Plaque Index (PI) measured at baseline, 1, 3, 6, 9, 12 months

4. Bleeding on Probing (BoP) measured at baseline, 1, 6, 9, 12 months
5. Full mouth plaque score (FMPS) measured at baseline
6. Patient-centred outcomes:
 - 6.1 Postoperative pain measured by VAS at 2 weeks
 - 6.2 Cumulative painkillers taken post-operatively at 2 weeks
 - 6.3 Surgical time

Overall study start date

25/10/2017

Completion date

31/12/2020

Eligibility

Key inclusion criteria

1. Patients who underwent implant surgery restored with dental implant-supported prosthesis, but PIKM is either absent or PIKM-W is less than 2mm
2. Patients prior to implant surgical procedure, but keratinised mucosa was insufficient at the edentulous area

Participant type(s)

Patient

Age group

Adult

Sex

Both

Target number of participants

48

Key exclusion criteria

1. Elective oral surgical intervention is contraindicated. In particular patients with uncontrolled or poorly controlled diabetes should be excluded.
2. Uncontrolled or untreated periodontal disease
3. Smoker
4. Infections or recent surgical procedures within 30 days of study initiation
5. Pregnant or lactating
6. Chronic treatment with any medication known to affect oral status (e.g., phenytoin, dihydropyridine, calcium antagonists, cyclosporine)
7. Radiotherapy or chemotherapy in the past 12 months prior to surgery
8. HIV or hepatitis
9. Physical handicaps that would interfere with the ability to perform adequate oral hygiene
10. Any investigational drug within 30 days of study initiation
11. Alcoholism or chronically drug abuse causing systemic compromise
12. Severe bruxism or clenching habits

Date of first enrolment

10/01/2018

Date of final enrolment

31/12/2019

Locations

Countries of recruitment

Hungary

Study participating centre

Department of Periodontology, Faculty of Dentistry, Semmelweis University

Szentkiralyi utca 47

Budapest

Hungary

1088

Sponsor information

Organisation

Department of Periodontology, Faculty of Dentistry, Semmelweis University

Sponsor details

Szentkiralyi utca 47

Budapest

Hungary

1088

+36 1 4591500/59188

attila.horvath.dr@hotmail.com

Sponsor type

University/education

Website

<http://semmelweis.hu/parodontologia/>

Funder(s)

Funder type

University/education

Funder Name

Results and Publications

Publication and dissemination plan

Publication in international peer-reviewed journals

Intention to publish date

01/02/2021

Individual participant data (IPD) sharing plan

All data generated or analysed during this study will be included in the subsequent results publication

IPD sharing plan summary

Other