

# The feasibility and acceptability of acceptance and commitment therapy for depression and/or anxiety during pregnancy

<b>Submission date</b>	<b>Recruitment status</b>	<input checked="" type="checkbox"/> Prospectively registered
02/02/2026	Recruiting	<input type="checkbox"/> Protocol
<b>Registration date</b>	<b>Overall study status</b>	<input type="checkbox"/> Statistical analysis plan
03/02/2026	Ongoing	<input type="checkbox"/> Results
<b>Last Edited</b>	<b>Condition category</b>	<input type="checkbox"/> Individual participant data
03/02/2026	Mental and Behavioural Disorders	<input checked="" type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Mental health conditions are highly prevalent in low to middle-income countries, particularly during the perinatal period. In Pakistan, rates of perinatal depression and anxiety have been reported at around 37% and 39%. There is a need for psychological interventions to support mothers and families impacted by perinatal depression and anxiety. Acceptance and Commitment Therapy for Perinatal Mental Health (ACT-PNMH) has been developed to meet this need. ACT-PNMH is an evidence-based intervention designed to improve symptoms of depression and anxiety during pregnancy and the postnatal period. We are testing ACT-PNMH in the NUR study to understand if the intervention and the methods that we use for the evaluation are helpful and acceptable for pregnant women in Pakistan. We also hope to explore if ACT influences mothers' mental health in the postnatal period and their relationship with their baby. To do this, we will compare the outcomes of women who received ACT-PNMH with those who received treatment as usual (TAU). TAU is the standard antenatal care delivered in the area where you live.

### Who can participate?

Pregnant women aged 18 years and above with psychological distress, anxiety or depression

### What does the study involve?

We will deliver ACT-PNMH over eight consecutive weeks to groups of 8-12 women in community venues. ACT-PNMH will be delivered by healthcare professionals in Pakistan who have been trained in the intervention by the team that developed ACT-PNMH from Cardiff University, Wales, UK.

When meeting with your healthcare provider, you will be invited to take part in the study, and you will be provided with information to make an informed decision about participating or not. If you are interested in taking part, you will be asked to complete a consent to contact form and a screening questionnaire about your mental health and pregnancy. If you are eligible to take part in the NUR study, you will then be contacted by a researcher from the study team, who can answer any further questions you may have about the study. If you are happy to take part, you will then complete a consent form and the baseline questionnaire about your mental health and

pregnancy. You will then be placed, at random, into either the 8-week ACT-PNMH intervention or the treatment-as-usual group (TAU).

If you are allocated to the ACT-PNMH intervention or the TAU group, you will be asked to complete questionnaires 9 weeks later and again when your baby is 3 months old. You will also be asked to complete a brief video recording of you playing with your baby at 3 months postpartum, and we will ask you some brief questions about your baby's development. If you are allocated to the ACT-PNMH intervention, you will be asked to complete an additional brief questionnaire at session three and at session five. We will also invite you to participate in semi-structured interviews about your experience of taking part in the NUR study and of the ACT-PNMH intervention. If you are allocated to the treatment-as-usual group, you will be offered the ACT-PNMH intervention after your baby is 3 months old.

**What are the possible benefits and risks of participating?**

Currently, there is a lack of information about the helpfulness and long-term benefits of psychological interventions for the treatment of anxiety and depression in Pakistan, compared to usual care. Regardless of the group you will be randomised into, your input will help us learn about how best to support perinatal women who experience depression and anxiety and how best to evaluate psychological interventions in Pakistan. As a small token of our appreciation for taking part, we can offer you 2000 Rupees for completing the baseline questionnaire, 2000 Rupees for completing the 9-week follow-up questionnaire, and 2000 Rupees for completing the 3-month postpartum assessment.

We are not expecting there to be any disadvantages or risks to you or your baby from taking part in the NUR study. You may feel anxious before sessions or tired after taking part in sessions. This may also be the case when completing the questionnaires. We will do everything we can to minimise or prevent this, and you may notice that the anxiety lessens as you get to know the ACT-PNMH facilitators over the 8 weeks.

**Where is the study run from?**

Health Services Academy (Pakistan)

**When is the study starting and how long is it expected to run for?**

February 2026 to October 2026

**Who is funding the study?**

Higher Education Funding Council for Wales, MEDR, through Official Development Assistance funding

**Who is the main contact?**

Dr Samina Naeem Khalid ([drsamina@hsa.edu.pk](mailto:drsamina@hsa.edu.pk)) is the main contact for the study in Pakistan

Dr Cerith Waters ([waterscs@cardiff.ac.uk](mailto:waterscs@cardiff.ac.uk)) is the main contact for the study in Wales, UK

## Contact information

### Type(s)

Principal investigator

### Contact name

Dr Cerith Waters

### Contact details

School of Psychology  
Cardiff University  
Tower Building  
Park Place  
Cardiff  
United Kingdom  
CF10 3AT  
+44 (0)292087026  
waterscs@cardiff.ac.uk

#### Type(s)

Scientific

#### Contact name

Dr Jennifer Berrett

#### Contact details

School of Psychology  
Cardiff University  
Tower Building  
Park Place  
Cardiff  
United Kingdom  
CF10 3AT  
+44 (0)292087026  
ThomasJL4@cardiff.ac.uk

#### Type(s)

Public

#### Contact name

Prof Samina Naeem Khalid

#### Contact details

Department of Maternal, Neonatal and Child Health  
Health Services Academy  
Ministry of National Health Services - Regulations and Coordination  
Prime Minister's National Health Complex,  
Government of Pakistan; Chak Shahzad, Park Road  
Islamabad  
Pakistan  
44000  
+92 (0)51-9255590-94  
drsamina@hsa.edu.pk

## Additional identifiers

## Study information

#### Scientific Title

# The NUR study: a pilot randomised controlled trial and Nested process evaluation stUdy of acceptance and commitment theRapy for pregnant women with antenatal depression and/or anxiety in Pakistan

## Acronym

NUR

## Study objectives

### Primary objectives (feasibility and acceptability):

The primary objective is to evaluate the feasibility and acceptability of the pilot RCT design and of the culturally adapted Acceptance and Commitment Therapy for Perinatal Mental Health (ACT-for-PNMH) intervention.

1. Feasibility of recruitment and retention: determine whether eligible pregnant women can be successfully identified, approached, recruited, randomised, and retained throughout the study.
2. Feasibility of intervention delivery: assess whether participants can attend the ACT-for-PNMH sessions as planned, including session attendance, completion, and reasons for non-attendance.
3. Data feasibility: evaluate whether outcome measures and process measures can be completed reliably, with acceptable levels of data completeness and minimal missing data.
4. Intervention fidelity and adherence: assess whether the ACT-for-PNMH intervention can be delivered as intended, including adherence to the intervention manual and facilitator fidelity.
5. Acceptability to participants and facilitators: explore the acceptability of the intervention, study procedures, and research processes from the perspectives of participants and facilitators.

### Secondary objectives (preliminary efficacy):

To explore potential signals of clinical improvement and estimate variability to inform future sample size calculations.

1. Psychological symptom change: explore initial trends and variability in psychological distress, depression, and anxiety symptoms, e.g., psychological distress (CORE-10), depression (PHQ-9), and anxiety (GAD-7).
2. Functional outcomes: explore trends and variability in maternal functioning, e.g., CORE-10, quality of life (EUROQoL).
3. Intervention mechanisms: explore trends and variability in ACT-PNMH intervention mechanisms, including psychological flexibility (Multidimensional Psychological Flexibility Inventory [MPFI]) and the therapeutic alliance (Working Alliance Inventory – Short Form [WAI-SF]).
4. Impact on childbirth, infant and mother-infant relationship: explore trends and variability in outcomes on childbirth (Birth Experiences Questionnaire [BEQ]), infant (WHO-Global Scales for Early Development [GSED]) and mother-infant relationship outcomes (Postpartum Bonding Questionnaire [PBQ]; audio-visual recording of mother-infant interaction).

### Process evaluation objectives:

To understand how the intervention and the RCT procedures operate within real-world antenatal care settings and identify factors that may support or hinder scalability.

1. Implementation in routine care: examine how the ACT-for-PNMH intervention is delivered within existing healthcare systems and identify organisational, structural, and contextual factors affecting implementation.
2. Mechanisms of impact: explore how participants engage with ACT-for-PNMH components and how these may contribute to changes in psychological or functional outcomes.
3. Contextual influences: identify barriers and facilitators to delivery, including cultural, logistical, and system-level factors that may influence future scalability and sustainability.

**Ethics approval required**

Ethics approval required

**Ethics approval(s)**

approved 16/01/2026, Health Services Academy, Institutional Ethical Review Committee (PM National Health Complex, Park Road, Chak, Shahzad, Islamabad,, 44000, Pakistan; (92-51) 9255590.94; academy@hsa.edu.pk), ref: No.7.82/IERC-HSA/2022-241

**Primary study design**

Interventional

**Allocation**

Randomized controlled trial

**Masking**

Open (masking not used)

**Control**

Active

**Assignment**

Parallel

**Purpose**

Health services research, Treatment

**Study type(s)****Health condition(s) or problem(s) studied**

Clinically significant antenatal depression and/or anxiety symptoms in the second trimester of pregnancy

**Interventions**

Computer-generated, 1:1 allocation to ACT-PNMH plus usual care or usual care alone, stratified by site and baseline severity.

**Intervention:**

Eight weekly sessions (90-minute duration) of group-delivered ACT-PNMH by trained therapists, plus usual care. The original ACT-PNMH manual (Waters et al., 2020) has been translated and culturally adapted for the Urdu-speaking population of Pakistan (Maqsood et al., 2025).

Intervention sessions will be delivered in person or remotely (e.g. via videoconferencing) if a participant cannot attend a given group session.

**Comparator:**

Usual care provided by standard antenatal mental health or primary care services. There is no standard model of care for antenatal depression and anxiety in Pakistan, but could include support provided by general practitioners, midwives and/or referrals to mental health services provided by psychologists and/or psychiatrists.

**Intervention Type**

Behavioural

## **Primary outcome(s)**

1. Recruitment rates measured using number approached, number declined, number consented at baseline
2. Reasons for declining RCT and/or intervention measured using qualitative methods at baseline
3. Intervention completion rates, defined as attending 4+ sessions, measured using descriptive statistics (%) at weekly and post-treatment
4. Mean, median and modal number of sessions attended measured using mean, median and mode at weekly sessions during intervention delivery and collated at post-treatment
5. Reasons for missing a session measured using a checklist at weekly sessions during intervention delivery and post-treatment
6. Reasons for dropping out of intervention before completion measured using a checklist at weekly sessions during intervention delivery and post-treatment
7. The completeness of outcome measures and data collection measured using % at baseline, post-treatment and 3-month follow-up
8. ACT-for-PNMH adherence measured using % of sessions that are adherent to the manual assessed at weekly sessions during intervention delivery
9. ACT fidelity ratings measured using the ACT Fidelity Measure (ACT-FM) at weekly sessions during intervention delivery
10. The acceptability of the ACT-for-PNMH intervention measured using Likert rating scales (range 0-5) at sessions 3, 5 and 8
11. Acceptability of the RCT design measured using Likert rating scales (range 0-5) at baseline and post-treatment
12. Adverse events measured using a checklist at baseline, weekly during intervention delivery, at post-treatment, and at 3 months postpartum

## **Key secondary outcome(s)**

### **Completion date**

02/10/2026

## **Eligibility**

### **Key inclusion criteria**

1. Age: 18 years or older
2. Gestation: 16–28 weeks at the start of the intervention
3. Consent: able to provide informed consent
4. Psychological distress: clinical range on the Clinical Outcomes in Routine Evaluation 10 (CORE-10)

10) score  $\geq 11$  and either the Patient Health Questionnaire-9 (PHQ-9) score  $\geq 10$  (depression) and /or General Anxiety Disorder-7 (GAD-7) score  $\geq 8$  (anxiety)

5. Intervention attendance: able to attend the intervention sessions

### **Healthy volunteers allowed**

No

### **Age group**

Mixed

### **Lower age limit**

18 years

### **Upper age limit**

100 years

### **Sex**

Female

### **Total final enrolment**

0

### **Key exclusion criteria**

1. Severe psychiatric conditions: current psychosis or bipolar disorder
2. Substance use: current substance dependence
3. Concurrent psychological treatment: receiving another structured psychological intervention at baseline
4. Medication changes: change in psychotropic medication within one month prior to randomisation
5. Suicidal risk: suicidal intent or high risk at baseline assessment
6. Other conditions: any medical or obstetric condition that would prevent participation in the intervention or compromise safety (e.g., severe pregnancy complications)

### **Date of first enrolment**

06/02/2026

### **Date of final enrolment**

26/04/2026

## **Locations**

### **Countries of recruitment**

Pakistan

### **Study participating centre**

**Health Services Academy, Ministry of National Health Services**  
PM National Health Complex, Park Road, Chak, Shahzad

Islamabad  
Pakistan  
44000

## Sponsor information

### Organisation

Health Services Academy Islamabad, Ministry of National Health Services

## Funder(s)

### Funder type

### Funder Name

Medr

## Results and Publications

### Individual participant data (IPD) sharing plan

### IPD sharing plan summary

Not expected to be made available