

# Adherence to Mediterranean diet can reduce the risk of obesity in people with genetic predisposition to obesity

<b>Submission date</b> 23/07/2017	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
<b>Registration date</b> 02/08/2017	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
<b>Last Edited</b> 20/09/2023	<b>Condition category</b> Nutritional, Metabolic, Endocrine	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

The obesity epidemic worldwide is fast increasing and affecting individuals of all ages, races and both genders. It is related with a series of other issues including diabetes, hypertension, cancer and cardiovascular diseases. Obesity is caused by genetic and environmental factors such as over-consumption of energy and sedentary lifestyle. Of genetic variants, fat mass and the obesity associated gene (FTO) have been found to be consistently associated with obesity traits in several populations. However there is increasing interest in finding out whether lifestyle factors modify the association of FTO variants and obesity as this could better provide insight into the role of diet/environmental factors in obesity. The Mediterranean diet is a diet that is high in vegetables, olive oil and fish (lean sources of protein). A higher adherence to the Mediterranean dietary pattern (Med Diet) using Mediterranean diet scores (MDS) has been associated with a decrease in obesity, regardless of genetic risk. This study uses participants in the Tehran Lipid and Glucose Study (TLGS). The TLGS is a large-scale, community-based, prospective study being performed on a sample of residents of District 13 of Tehran, capital of Iran. The first phase of the TLGS was conducted from 1999 to 2001 and follow-up examinations have been conducted every 3 years (2002–2005; 2006–2008; 2008–2011, and 2011–2014) to identify newly developed diseases. The aim of this study is to investigate whether Med Diet could interact with FTO in relation to obesity among adult participants in order to identify environmental interactions to create preventive approaches in individuals with greater genetic susceptibility to obesity.

### Who can participate?

Adults aged 18 years and older.

### What does the study involve?

Participants undergo a baseline survey. Those who are evaluated as obese are excluded. The other participants are followed and randomly selected for follow-up surveys. The surveys include questions about food frequency and types of food ingested in order to provide a MDS score to participants. Other measures such as physical activity, smoking, and other demographics are also assessed. DNA samples are provided by participants in order to identify their genetic risk.

What are the possible benefits and risks of participating?

Participants may help prevent obesity in those with FTO risk alleles. There are no risks with participating.

Where is the study run from?

This study is run by the Shahid Beheshti University of Medical Sciences (Iran) and the TLGS is performed on a sample of residents of District 13 of Tehran (Iran).

When is the study starting and how long is it expected to run for?

February 1998 to February 2017

Who is funding the study?

Research Institute for Endocrine Sciences, Shahid Beheshti University of Medical Sciences (Iran)

Who is the main contact?

Parvin Mirmiran, Prof.

mirmiran@endocrine.ac.ir

## Contact information

### Type(s)

Scientific

### Contact name

Dr Firoozeh Hosseini-Esfahani

### ORCID ID

<http://orcid.org/0000-0002-9697-0081>

### Contact details

Nutrition and Endocrine Research Centre  
Research Institute for Endocrine Sciences  
Shahid Beheshti University of Medical Sciences  
No.24, Arabi Street  
Yemen Avenue  
Chamran Highway  
Tehran  
Iran  
1985717413  
+98 212 243 2500  
f.hosseini@sbmu.ac.ir

### Type(s)

Scientific

### Contact name

Prof Parvin Mirmiran

### ORCID ID

<http://orcid.org/0000-0003-2391-4924>

## Contact details

Nutrition and Endocrine Research Centre  
Research Institute for Endocrine Sciences  
Shahid Beheshti University of Medical Sciences  
No.24, Arabi Street  
Yemen Avenue  
Chamran Highway  
Tehran  
Iran  
1985717413  
+98 212 243 2500  
mirmiran@endocrine.ac.ir

## Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

### Secondary identifying numbers

Research Institute for Endocrine Sciences (Grant No. 840), Shahid Beheshti University of Medical Sciences

## Study information

### Scientific Title

Mediterranean dietary pattern adherence modify the association between FTO genetic variations and obesity phenotypes

### Study objectives

Can Mediterranean Diet interact with FTO gene polymorphisms (rs1121980, rs1421085, rs9939973, rs8050136, rs17817449 and rs3751812) in isolation or combined form (GRS) in relation to obesity phenotypes among participants of the Tehran Lipid and Glucose Study (TLGS)?

### Ethics approval required

Old ethics approval format

### Ethics approval(s)

Ethics committee of the Research Institute for Endocrine Sciences, Shahid Beheshti University of Medical Sciences, 10/01/1999, ref: 840

### Study design

Observational-nested case-control study

### Primary study design

Observational

### Secondary study design

Nested case-control study

**Study setting(s)**

Community

**Study type(s)**

Other

**Participant information sheet**

**Health condition(s) or problem(s) studied**

Obesity

**Interventions**

Participants undergo a baseline survey. Those who are evaluated as obese are excluded. The other participants are followed and randomly selected for follow-up surveys between 2006-2008, 2008-2011 and 2011-2014. Each of these cases are individually paired and matched by age (plus or minus five years), sex and duration of follow-up with a random control from a population with normal body mass index. The duration of follow-up is from baseline to the follow up survey which marks the end of their participation in the study.

The survey that participants fill out assessed participants dietary intakes using a valid and reliable 168-item semi-quantitative food frequency questionnaire (FFQ) at baseline and follow up surveys. The Mediterranean dietary score is computed according to richopoulou et al. based on the following eight components: Legumes, vegetables, nuts and fruits, fish, and cereals intake; dietary ratio of monounsaturated fatty acid (MUFA) to saturated fatty acid (SFA); intake of dairy products, mostly in the form of cheese or yogurt and intakes of meat and poultry. The anthropometric, physical activity, smoking and the other demographic variables are assessed at baseline and follow-up surveys, DNA products of participants were genotyped and genetic risk score is computed.

The risk of obesity (odds ratio) of individuals with different genetic risk scores assessed across the Mediterranean dietary score quartiles from baseline to follow-up survey.

**Intervention Type**

Other

**Primary outcome measure**

1. Weight is measured using the digital scales at baseline and each follow-up examination (every three years)
2. Height is measured using a tape measure at baseline and each follow-up examination (every three years)
3. Waist circumference is measured using the upstretched tape meter at baseline and each follow-up examination (every three years)
4. Hip circumference is measured using the upstretched tape meter at baseline and each follow-up examination (every three years)

**Secondary outcome measures**

1. BMI (body mass index) is calculated by dividing weight in kilograms on height in meters squared baseline ( $\text{kg/m}^2$ ) at baseline and each follow-up examination (every three years)
2. WHR (waist to hip ratio) is calculated by dividing waist on hip circumferences at baseline and each follow-up examination (every three years)

3. Obesity was defined as a BMI  $\geq 30$  kg/m<sup>2</sup>
4. BMI between 18.5 and 24.9 classified a person as having a normal weight
5. Waist circumference  $>95$  cm for both genders was considered as indicators of abdominal obesity
6. WHR  $>0.8$  in men and  $>0.9$  in women, was considered as indicators of abdominal obesity

**Overall study start date**

01/02/1998

**Completion date**

15/02/2017

## Eligibility

**Key inclusion criteria**

1. Age  $\geq 18$  years
2. Both genders
3. Participating in baseline and follow-up examinations of Tehran Lipid and Glucose Study
4. BMI  $\geq 30$  for cases, BMI  $\geq 18.5$  and  $<25$  for controls.
5. Not pregnant or lactating women
6. Not history of weight loss or gain  $>5$  kg in the last 6 months
7. Not taking drugs that affect weight
8. Not under or over-reporters of energy intake
9. DNA purification in the range of  $1.7 < A_{260}/A_{280} < 2$

**Participant type(s)**

Mixed

**Age group**

Adult

**Lower age limit**

18 Years

**Sex**

Both

**Target number of participants**

627 pairs of persons with obesity and their matched controls (1254 persons)

**Total final enrolment**

1000

**Key exclusion criteria**

1. Individuals with a history of weight loss or gain  $>5$  kg in the last 6 months
2. Pregnant and lactating
3. Taken drugs that affect weight
4. Cases/controls lacking DNA purification in the range of  $1.7 < A_{260}/A_{280} < 2$
5. Reported energy intakes divided by the predicted energy intake did not qualify for the  $\pm 3SD$  range

**Date of first enrolment**

31/01/1999

**Date of final enrolment**

20/01/2015

## **Locations**

**Countries of recruitment**

Iran

**Study participating centre****Shahid Beheshti University of Medical Sciences**

Nutrition and Endocrine Research Center

Research Institute for Endocrine Sciences

No. 24, Arabi Street

Yemen Avenue

Chamran highway

Tehran

Iran

1985717413

## **Sponsor information**

**Organisation**

Shahid Beheshti University of Medical Sciences

**Sponsor details**

Nutrition and Endocrine Research Centre

Research Institute for Endocrine Sciences

No. 24, Arabi Street

Yemen Avenue

Chamran Highway

Tehran

Iran

1985717413

+98 212 243 2500

mirmiran@endocrine.ac.ir

**Sponsor type**

Research organisation

**ROR**

<https://ror.org/034m2b326>

# Funder(s)

## Funder type

Not defined

## Funder Name

Shahid Beheshti University of Medical Sciences

## Alternative Name(s)

Shahid Beheshti University of Medical Sciences (SBMU), Iran, Shahid Beheshti University of Medical Sciences (Iran), Shahid Beheshti University of Medical Sciences - Iran, , Université des Sciences Médicales de Shahid Beheshti, , Shahid Beheshti University of Medical Sciences Tehran Iran, Shahid Beheshti University of Medical Sciences Tehran, Université des Sciences Médicales de, , , SBUMS

## Funding Body Type

Government organisation

## Funding Body Subtype

Universities (academic only)

## Location

Iran

# Results and Publications

## Publication and dissemination plan

Phase one and two have been published. The manuscript is planned to submit in American Journal of Clinical Nutrition as soon as we receive Clinical Trial Registration Number.

## Intention to publish date

15/08/2017

## Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from Dr. Amir-Abbas Momenan at [momenan@gmail.com](mailto:momenan@gmail.com) or [momenan@endocrine.ac.ir](mailto:momenan@endocrine.ac.ir)

## IPD sharing plan summary

Available on request

## Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results (phase one)	01/05/2002		Yes	No
<a href="#">Results article</a>	results (phase two) in	25/01/2009		Yes	No

[Results article](#)

26/09/2017      20/09/2023      Yes      No