Cognitive behavioral therapy for psychological distress in young people diagnosed with cancer during adolescence

Submission date	Recruitment status No longer recruiting	Prospectively registered		
07/08/2017		[X] Protocol		
Registration date	Overall study status Completed Condition category Mental and Behavioural Disorders	Statistical analysis plan		
22/08/2017		Results		
Last Edited		Individual participant data		
21/09/2023		Record updated in last year		

Plain English summary of protocol

Background and study aims

Young people who are diagnosed with cancer during adolescence may suffer from psychological distress after the end of cancer treatment. Cognitive behavior therapy (CBT) is a talking therapy that is effective for a range of psychological disorders and has been tested for the treatment of depression, anxiety and post-traumatic stress disorder. There are currently few psychological treatments developed specifically to fit the needs of young survivors of cancer and no study has tested individualized CBT specifically tailored to reduce psychological distress in young survivors of cancer diagnosed during adolescence. The aim of this study is therefore to develop and test a CBT-based treatment targeting cancer-related psychological distress experienced by young people who have completed treatment for cancer diagnosed during adolescence.

Who can participate?

Young people aged between 15-25 who have been diagnosed with cancer during adolescence, have completed successful cancer treatment and experience cancer-related suffering

What does the study involve?

Participants are offered up to 15 weekly 45-minute sessions of individual CBT provided by three resident psychologists. Psychological distress is assessed before treatment, at the end of treatment, and at 3 months after the end of treatment.

What are the possible benefits and risks of participating?

By enrolling in this study participants can receive psychological support. There are no anticipated side effects of the treatment. Working with one's mental health problems can involve some distress. Participants who do not experience improvement or who deteriorate during the treatment might have less confidence in psychological treatments in the future. Participants in need of immediate/more intensive/other psychological treatment than can be offered within this study are guided to appropriate healthcare.

Where is the study run from?
Uppsala University Children's Hospital (Sweden)

When is the study starting and how long is it expected to run for? May 2014 to October 2016

Who is funding the study?

- 1. Swedish Research Council
- 2. Swedish Childhood Cancer Foundation
- 3. Swedish Cancer Society

Who is the main contact? Prof. Louise von Essen louise-von.essen@kbh.uu.se

Contact information

Type(s)

Scientific

Contact name

Prof Louise von Essen

ORCID ID

http://orcid.org/0000-0001-5816-7231

Contact details

Uppsala universitet
Uppsala biomedicinska centrum
Institutionen för kvinnors- och barns hälsa
Klinisk psykologi i hälso- och sjukvård
Box 572
Uppsala
Sweden
75123
+46 (0)184 713 484
louise-von.essen@kbh.uu.se

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

U-CARE: MAYA

Study information

Scientific Title

Development of psychological treatment for cancer-related suffering experienced by young people diagnosed with cancer during adolescence

Study objectives

Since there are no evidence-based interventions for young survivors of cancer during adolescence who experience cancer-related suffering, the purpose of the current study was to develop and preliminary evaluate if individual CBT can reduce distress among these survivors.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Regional Ethical Review Board in Uppsala, Sweden, 22/12/2014, Dnr: 2014/443

Study design

Single-group non-randomised open trial

Primary study design

Interventional

Secondary study design

Non randomised study

Study setting(s)

Other

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use the contact details to request a patient information sheet

Health condition(s) or problem(s) studied

Cancer-related psychological distress

Interventions

All participants are offered up to 15 weekly 45-minute sessions of individual CBT provided by three resident psychologists under supervision. The CBT interventions are based on a clinical behavior analysis. Psychological distress is assessed before treatment (baseline), at the end of treatment (post-treatment), and 3-months after end of treatment.

Intervention Type

Behavioural

Primary outcome measure

- 1. Anxiety, assessed with the Beck Anxiety Inventory (BAI) at baseline, post-treatment and at 3 months after end of treatment
- 2. Depression, assessed with the Montgomery Åsberg Depression Rating Scale Self-assessment (MADRS-S) at baseline, post-treatment and at 3 months after end of treatment

- 3. Post-traumatic stress symptoms, assessed with the The PTSD Checklist Civilian (PCL-C) at baseline, post-treatment and at 3 months after end of treatment
- 4. Worry, assessed with the Penn State Worry Questionnaire (PSWQ) at baseline, post-treatment and at 3 months after end of treatment

Secondary outcome measures

- 1. Body dissatisfaction, assessed with the Body Image Scale (BIS) at baseline, post-treatment and at 3 months after end of treatment
- 2. Experiential avoidance, assessed with the Acceptance and Action Questionnaire (AAQ-II) at baseline, post-treatment and at 3 months after end of treatment
- 3. Fatigue, assessed with the Fatigue Assessment Scale (FAS) at baseline, post-treatment and at 3 months after end of treatment
- 4. Functional impairment, assessed with the Sheehan Disability Scale (SDS) at baseline, post-treatment and at 3 months after end of treatment
- 5. Health anxiety, assessed with the Short Health Anxiety Inventory (SHAI) at baseline, post-treatment and at 3 months after end of treatment
- 6. Somatic symptoms, assessed with the Patient Health Questionnaire-15 (PHQ-15) at baseline, post-treatment and at 3 months after end of treatment
- 7. Rumination, assessed with the Rumination Scale of the Response Style Questionnaire (RSQ): at baseline, post-treatment and at 3 months after end of treatment
- 8. The presence of psychiatric disorders, assessed with the Mini-International Neuropsychiatric Interview (M.I.N.I), a structured diagnostic psychiatric interview for DSM-IV and ICD-10, at baseline, post-treatment and at 3 months after end of treatment

Overall study start date

01/05/2014

Completion date

10/10/2016

Eligibility

Key inclusion criteria

- 1. Aged 15-25 years at study start
- 2. Diagnosed with cancer when aged 13-18 years
- 3. Treated at a paediatric oncology unit in Sweden
- 4. Completed successful cancer treatment
- 5. Experience cancer-related suffering

Participant type(s)

Patient

Age group

Mixed

Sex

Both

Target number of participants

20

Key exclusion criteria

- 1. Currently receive psychological treatment
- 2. Display psychiatric symptoms in immediate need of treatment

Date of first enrolment

27/02/2015

Date of final enrolment

03/11/2015

Locations

Countries of recruitment

Sweden

Study participating centre Uppsala University Children's Hospital

Akademiska barnsjukhuset, ing. 95 nbv Uppsala Sweden 75185

Sponsor information

Organisation

Uppsala University

Sponsor details

Box 256 Uppsala Sweden 75105

Sponsor type

University/education

Website

www.uu.se

ROR

https://ror.org/048a87296

Funder(s)

Funder type

Government

Funder Name

Vetenskapsrådet

Alternative Name(s)

Swedish Research Council, VR

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

Sweden

Funder Name

Barncancerfonden

Alternative Name(s)

Swedish Childhood Cancer Foundation

Funding Body Type

Private sector organisation

Funding Body Subtype

Trusts, charities, foundations (both public and private)

Location

Sweden

Funder Name

Cancerfonden

Alternative Name(s)

Swedish Cancer Society

Funding Body Type

Private sector organisation

Funding Body Subtype

Trusts, charities, foundations (both public and private)

Results and Publications

Publication and dissemination plan

The results from the trial will be published in a doctoral dissertation at Uppsala University and will be submitted for publication in a scientific journal approximately in November 2017.

Intention to publish date

01/11/2017

Individual participant data (IPD) sharing plan

Ethical restrictions prohibit making the dataset publicly available. Data are available on request from the research group Clinical Psychology in Healthcare at the Department of Women's and Children's Health, Uppsala University for researchers who meet the criteria for access to confidential data. Requests should be sent to PI Prof. Louise von Essen (louise-von.essen@kbh. uu.se).

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Other publications	Qualitative feasibility results	17/04/2018	21/09/2023	Yes	No
<u>Protocol file</u>			21/09/2023	No	No