

Nasal airway obstruction study - NAIROS

Submission date 06/03/2017	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 24/03/2017	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 09/06/2025	Condition category Ear, Nose and Throat	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

A deviated septum is a condition where the thin wall separating the nostrils (nasal septum) is off-center or crooked (deviation). Most people only have a minor deviation, but it can lead to difficulty breathing, snoring and a blocked nose. A septoplasty is a type of operation that corrects a deviated septum by removing excess bone and cartilage to straighten the septum and nasal passages. Like any operation, there is a risk of complications. Most patients need to take at least 5 days off work or usual activities after the operation. Some patients seem not much better after the surgery. Practice varies around the country, and there is no good evidence about this operation or its alternatives, or about who might benefit most from treatment, to inform help patients and doctors decide when it should be carried out. The aim of this study is to compare the effectiveness septoplasty or non-surgical care (medical management), which consists of nasal sprays in the management of adults with deviated septums.

Who can participate?

Adults with have a deviated septum.

What does the study involve?

Participants are randomly allocated to one of two groups. Those in the first group are treated surgically using a septoplasty procedure. Those in the second group are given a six month supply of two different nasal sprays, each to use twice daily. At the start of the study and then again after six and 12 months, participants in both groups complete a number of questionnaire in order to assess their quality of life, nasal symptoms and how much they are using health care services as well as a test to measure how well they are breathing through their noses.

What are the possible benefits and risks of participating?

Participants may benefit from improvement in symptoms of nasal blockage, sleep disturbance and congestion/headache if these symptoms are related to blocked nose due to a deviated septum. For participants who receive the nasal spray, there is a risk of bleeding or irritation in the nose. For those who receive surgery, there is a risk of general complications from surgery, including minor bleeding, discomfort (common), heavy bleeding, temporary numbness of central upper teeth, hole in the septum (perforation) and cosmetic change in the appearance of nose (rare).

Where is the study run from?

Freeman Hospital (lead centre) and 15 other NHS hospitals in England, Scotland and Wales (UK)

When is the study starting and how long is it expected to run for?

March 2017 to June 2021

Who is funding the study?

National Institute for Health Research, Health Technology Assessment Programme (UK)

Who is the main contact?

1. Dr Sean Carrie (scientific)
 2. Dr Katherine Rennie (public)
- nairos.trial@newcastle.ac.uk

Contact information

Type(s)

Scientific

Contact name

Dr Sean Carrie

Contact details

Freeman Hospital
High Heaton
Newcastle Upon Tyne
United Kingdom
NE7 7DN

Type(s)

Public

Contact name

Dr Katherine Rennie

ORCID ID

<https://orcid.org/0000-0003-1703-3768>

Contact details

Newcastle University Clinical Trials Unit
Newcastle University
1-4 Claremont Terrace
Newcastle Upon Tyne
United Kingdom
NE2 4AE
+44 (0)191 208 2519
nairos.trial@newcastle.ac.uk

Additional identifiers

Clinical Trials Information System (CTIS)

2017-000893-12

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

8302

Study information

Scientific Title

Multicentre RCT to determine the clinical and cost effectiveness of septoplasty compared with non-surgical strategy in the management of nasal septal deviation in adult patients with nasal airway obstruction in the presence of a deviated nasal septum, what is effectiveness of septal surgery with turbinate reduction compared with 6 months' topical nasal treatment in improving nasal symptoms

Acronym

NAIROS

Study objectives

The aim of this study is to determine whether and to what extent septoplasty is superior to medical treatment in the management of adult patients with nasal obstruction in the presence of a septal deviation.

Ethics approval required

Old ethics approval format

Ethics approval(s)

North East – Newcastle & North Tyneside 2 Research Ethics Committee, 31/08/2017, ref: 17/NE/0239

Study design

Multi-centre randomized controlled trial with qualitative process and economic evaluation

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Deviated septum

Interventions

Patients will be randomised to one of two groups in a 1:1 ratio using a variable block stratified design stratified by severity (baseline self-report NOSE category – moderate 30-50; Severe 55-75; extreme 80-100) and gender.

Surgery: Participants will undergo septoplasty and turbinate reduction

Medical management: Participants receive medical management, which will comprise of regular use of a nasal saline spray (sterimar) followed by a fluorinated steroid spray Mometasone for 6 months.

All patients will be followed up at 6 and 12 months post randomisation.

Intervention Type

Mixed

Primary outcome(s)

Patient reported assessment of nasal and general symptoms is assessed using the SNOT 22 questionnaire (Sinonasal Outcome Test) at baseline, 6 and 12 months.

Key secondary outcome(s)

1. Objective assessment nasal airflow is measured using rhinosprometry (nasal partitioning ratio and peak nasal airflow measurements) at baseline, 6 and 12 months
2. Quality of life is measured using the Short Form 36 Quality of Life Questionnaire at baseline, 6 and 12 months
3. Health utilisation is measured using a health utilisation questionnaire at baseline, 6 and 12 months

Completion date

04/03/2021

Eligibility

Key inclusion criteria

1. Adults aged ≥ 18 years with a baseline NOSE score ≥ 30
2. Septal Deflection visible at naseondoscopy
3. Capacity to provide informed written consent and complete the study questionnaires.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

378

Key exclusion criteria

1. Patients <18 years
2. Patients with a baseline NOSE score <30
3. Any prior septal surgery
4. Systemic inflammatory disease
5. Granulomatosis with poly angiitis
6. Naseoscopic evidence of unrelated associated pathology eg. adenoid pad, septal perforation, chronic rhinosinusitis indicated by the prescence of polyposis or pus
7. Intranasal recreational drug use
8. Breastfeeding, pregnancy or intended pregnancy
9. Bleeding diathesis
10. Therapeutic anticoagulation
11. Known adverse reactions to general anaesthesia
12. Immunocompromised patients

Date of first enrolment

01/09/2017

Date of final enrolment

31/01/2020

Locations

Countries of recruitment

United Kingdom

England

Scotland

Wales

Study participating centre

Freeman Hospital

High Heaton

Newcastle Upon Tyne

United Kingdom

NE7 7DN

Study participating centre

Aberdeen Royal Infirmary

Foresterhill

Aberdeen

United Kingdom

AB25 2ZN

Study participating centre
Queen Elizabeth Hospital Birmingham
Mindelsohn Way
Edgbaston
Birmingham
United Kingdom
B15 2TH

Study participating centre
Bradford Royal Infirmary
Duckworth Lane
Bradford
United Kingdom
BD9 6RJ

Study participating centre
Darlington Memorial Hospital
Hollyhurst Road
Darlington
United Kingdom
DL3 6HX

Study participating centre
Ninewells Hospital
ENT Department (Ward 26)
Dundee
United Kingdom
DD1 9SY

Study participating centre
Monklands Hospital
Monkscourt Avenue
Airdrie
United Kingdom
ML6 OJS

Study participating centre
Leeds General Infirmary
Great George Street
Leeds

United Kingdom
LS1 3EX

Study participating centre

Aintree Hosptial

Lower Lane
Liverpool
United Kingdom
L9 7AL

Study participating centre

Royal Gwent Hospital

Cardiff Road
Newport
United Kingdom
NP20 2UB

Study participating centre

Cumberland Infirmary

Newtown Road
Carlisle
United Kingdom
CA2 7HY

Study participating centre

Derriford Hospital

Derriford Road
Plymouth
United Kingdom
PL6 8DH

Study participating centre

Salisbury District Hospital

Odstock Road
Salisbury
United Kingdom
SP2 8BJ

Study participating centre

Wrightington Hospital

Hall Lane
Appley Bridge
Wigan
United Kingdom
WN6 9EP

Study participating centre**Guy's Hospital**

Great Maze Pond
London
United Kingdom
SE1 9RT

Study participating centre**Norfolk and Norwich University Hospital**

Colney Lane
Norwich
United Kingdom
NR4 7UY

Sponsor information**Organisation**

Newcastle Upon Tyne NHS Foundation Trust

ROR

<https://ror.org/05p40t847>

Funder(s)**Funder type**

Government

Funder Name

National Institute for Health Research

Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

The current data sharing plans for the current study are unknown and will be made available at a later date.

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		18/10/2023	19/10/2023	Yes	No
Results article		13/03/2024	13/03/2024	Yes	No
Results article	Healthcare professionals' and patients' views and experiences of surgical and medical treatment for nasal obstruction: a qualitative interview study for a Nasal Airway Obstruction Study (NAIROS)	08/06/2025	09/06/2025	Yes	No
Protocol article	protocol	13/02/2020	17/02/2020	Yes	No
Basic results		17/12/2021	16/06/2022	No	No
HRA research summary			28/06/2023	No	No
Study website	Study website	11/11/2025	11/11/2025	No	Yes