

The effect of dry needling in gluteus medius on hip strength and stability in active populations

Submission date 20/06/2018	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 24/07/2018	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
Last Edited 22/06/2018	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Trigger points are muscular "knots" which can often be felt as tightness or aches in muscles. They are commonly found in both sportspeople and the general population alike. One common treatment technique which has recently gained popularity is that of dry needling. Dry needling uses acupuncture type needles to directly stimulate the trigger point and in doing so, release the tightness (contraction) and pain associated with the trigger point. There are two categories of trigger points: active (trigger points which generate pain spontaneously) and latent (trigger points which only generate pain if stimulated). Because dry needling relaxes the trigger point within muscles, some temporary weaknesses have been reported previously. If these weaknesses were to occur in a muscle which is crucial in stability, athletes may be at an increased risk of injury. The gluteus medius is one such muscle which plays a pivotal role in maintaining hip stability during movements and single leg tasks. It is located on the outermost part of the pelvis and controls the side to side movement of the leg (frontal plane), providing what is known as lumbo-pelvic-hip stability. Decreases in either hip strength or hip stability have been linked to a host of lower limb injuries to the foot, ankle, knee, thigh, hip and even lower back. Trigger points are commonplace in the gluteus medius of multi-directional athletes because of the demands placed on maintaining hip stability during sprinting, cutting, jumping and landing. As such, dry needling is a technique frequently performed on the gluteus medius yet it is still unclear as to the appropriate time before performance which dry needling should be performed, or more importantly, avoided due to possible temporary decreases in hip strength and stability. The aim of this study is to measure hip strength and single leg stability before and after dry needling.

Who can participate?

Male, third level student of the Institute of Technology Carlow, aged 18 to 45, participating in a multi-directional jump-sport (e.g. gaelic football, hurling, soccer, volleyball, basketball, handball) more than 3 times per week, with 2 to 5 latent myofascial trigger points present in their self-reported preferred jumping leg

What does the study involve?

Participants are randomly allocated to be treated with dry needling or dummy dry-needling (placebo) where the needle does not break the skin. Participants only receive the treatment once and it is performed on between 2 and 5 trigger points. There is a quick informal

conversation about how the participant feels about the treatment received, their soreness and their confidence to perform in their given sport. Hip strength and single leg stability are measured before, immediately after and 3, 7, 10 and 14 days after the treatment.

What are the possible benefits and risks of participating?

Participants should they volunteer, may benefit from dry needling as it may reduce tension in their gluteus medius and allow their muscle to perform better in the long term. Some soreness is expected both during the dry needling treatment and in the 24 hours afterwards with slight weaknesses possibly being felt by participants. The participants may also have a slight bruise as a result of the needle piercing the skin.

Where is the study run from?

Institute of Technology Carlow (Ireland)

When is the study starting and how long is it expected to run for?

September 2017 to August 2019

Who is funding the study?

Institute of Technology Carlow (Ireland)

Who is the main contact?

Mr Aaron Byrne

Aaron.Byrne@itcarlow.ie

Contact information

Type(s)

Scientific

Contact name

Mr Aaron Byrne

Contact details

K221 Dargan Centre

Institute of Technology Carlow

Kilkenny Road

Carlow

Ireland

0000

+353 (0)862331849

aaron.byrne@itcarlow.ie

Additional identifiers

Protocol serial number

182

Study information

Scientific Title

The longitudinal effects of dry needling of latent myofascial trigger points in gluteus medius on hip strength and single-leg time to stabilisation in multi directional and jump-sport athletes

Study objectives

1. Dry needling will cause a short term decrease in hip strength
2. Dry needling will cause a short term increase in time to stabilisation

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics in Research Committee in the Departments of Science and Health of the Institute of Technology Carlow, Ireland, 26/10/2017, ref: 182

Study design

Randomised placebo-controlled intervention study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Latent myofascial trigger points in gluteus medius

Interventions

All participants will be randomly assigned one of the treatments and will not know which treatment they are receiving (single-blinded):

1. Dry needling (intra-muscular stimulation)
2. Placebo dry needling (blunted needle, zero skin penetration)

Participants will only receive the single treatment once and it will be performed on between 2 and 5 trigger points. There will also be a quick informal conversation about how the participant feels about the treatment received, their soreness and their confidence to perform in their given sport.

Hip strength and single leg stability are measured before, immediately after and 3, 7, 10 and 14 days after the treatment. Hip strength will be measured using a portable strain gauge (hand-held dynamometer) which will be fitted to the end of a stabilisation device. Single leg stability will be recorded on a force plate in the form of "time to stabilisation" which records the amount of time needed to stabilise the body to within a certain threshold of bodyweight (5%), a lower time to stabilisation corresponds to better stability as they are able to stabilise faster.

Intervention Type

Other

Primary outcome(s)

Measured before, immediately after and 3, 7, 10 and 14 days after the treatment:

1. Hip abduction strength (maximal voluntary isometric contraction), measured using a portable strain gauge (hand-held dynamometer)

2. Hip internal rotation strength (maximal voluntary isometric contraction), measured using a portable strain gauge (hand-held dynamometer)
3. Hip external rotation strength (maximal voluntary isometric contraction), measured using a portable strain gauge (hand-held dynamometer)
4. Single-leg time to stabilisation, recorded on a force plate

Key secondary outcome(s)

The secondary outcome measures will be qualitative by nature and obtained through semi-structured interviews upon the completion of each of the follow-up measurements of stability (immediately post treatment, 3, 7, 10 and 14 days post treatment). All interviews will be conducted in private by a single interviewer and will be audio recorded for thematic analysis at a later date in order to give greater depth as to the effects dry needling may have on sporting performance.

1. Perceived stability
2. Perceived ability to perform in their given sport
3. Confidence in ability to perform
4. Local soreness (VAS)

Completion date

01/08/2019

Eligibility

Key inclusion criteria

Participants meet inclusion criteria if they:

1. Are a male, third level student of the Institute of Technology Carlow
2. Are aged between 18 and 45 years
3. At the time of testing, are participating in a multi-directional jump-sport including but not limited to; gaelic football, hurling, soccer, volleyball, basketball, handball more than 3 times per week
3. Have, as identified in a formal screening session; between 2 and 5 latent myofascial trigger points present in their self-reported preferred jumping leg

Participant type(s)

Healthy volunteer

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

Male

Key exclusion criteria

Participants will be excluded if they:

1. Have, at the time of testing, any injury to the lower limb or low back. Injury is defined as "any injury that prevents a player from taking a full part in all training and match play activities" (Brooks et al. 2005)
2. Present with signs of or report during screening process, any neurological or bleeding disorders
3. Have a needle phobia or aicmophobia

Date of first enrolment

01/09/2018

Date of final enrolment

22/12/2018

Locations

Countries of recruitment

Ireland

Study participating centre

Institute of Technology Carlow

Kilkenny Road

Carlow

Ireland

0000

Sponsor information

Organisation

Institute of Technology Carlow

Funder(s)

Funder type

University/education

Funder Name

Institute of Technology Carlow

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from Mr Aaron Byrne (aaron.byrne@itcarlow.ie). Upon the completion of the trial, all anonymised raw datasets will be made available upon request and will also be present in appendices of the final dissertation submission.

IPD sharing plan summary

Available on request