# Is internet-delivered therapy more effective than internet-delivered brief support in reducing depressive symptoms in adolescents?

Submission date	Recruitment status  No longer recruiting	<ul><li>Prospectively registered</li></ul>			
20/12/2018		☐ Protocol			
Registration date	Overall study status	Statistical analysis plan			
25/01/2019	Completed	[X] Results			
Last Edited	Condition category	Individual participant data			
28/02/2024	Mental and Behavioural Disorders				

## Plain English summary of protocol

Background and study aims

The need of development of early, short and focused interventions for depressed adolescents is urgent. Internet-based interventions are designed to increase access to psychological treatments and cost-effectiveness. Empirical support is already established for Internet-based cognitive behavioural therapy (I-CBT) with response rate of about 60 %. In the present study, the aim is to test the feasibility and efficacy of a recently developed internet-based treatment for depressed adolescents – internet-delivered psychodynamic therapy (I-PDT).

## Who can participate?

Adolescents aged 15-18 years with mild to moderate depression will be included.

## What does the study involve?

The study is a randomized controlled trial for 72 participants, in which half of the participants will begin treatment immediately after inclusion and the other half will receive a brief supportive contact until they receive the treatment 10 weeks after inclusion. The interventions consist of 8 therapist-supported self-help modules delivered over 8 weeks with added chat sessions. Primary outcome will be severity of depressive symptoms. Secondary outcomes will be anxiety symptoms and psychiatric diagnoses. The aim of I-PDT is to decrease emotional avoidance and increase awareness and experience of emotions. Participants are encouraged to gradually approach previously warded off feelings. They will also be taught how to link their emotions to their symptoms. Another treatment goal is to acquire a greater capacity for anxiety regulation.

## What are the possible benefits and risks of participating?

The expected benefit of the treatment is that it will reduce depressive symptoms among the participants. Internet-based treatments bear the possibility to reach depressed adolescents who do not have access to adequate psychological treatment, for example due to geographical reasons, or are reluctant to seek face-to-face treatment within the health care system. Psychological treatments are based on the fact that participants share very personal information and there is a risk that some participants might experience this as a threat to their integrity. The

project will use established strategies to manage other types of risks, for example that some participants turns out having more serious medical, psychological or social problems than was discovered at inclusion.

Where is the study run from?

Department of Psychology, Stockholm University, Sweden.

When is the study starting and how long is it expected to run for?

This trial is a pilot study that will start in 3 January, 2019, and run until 30 June, 2019. The pilot study is the first part of a larger project and the next step will be a large non-inferiority randomized controlled trial investigating comparative effects between I-PDT and an already established internet based treatment (I-CBT) for adolescents.

Who is funding the study?

The project has received funding from the Kavli Trust for the years 2019-2023.

Who is the main contact?

Main contact is principal investigator Björn Philips, Associative Professor, at the Department of Psychology, Stockholm University, Sweden.

# **Contact information**

## Type(s)

Scientific

#### Contact name

Dr Björn Philips

#### Contact details

Department of Psychology Stockholm University Stockholm Sweden SE-106 91 +46 8 162010 bjorn.philips@psychology.su.se

# Additional identifiers

Protocol serial number

2018/2268-31/5

# Study information

#### Scientific Title

ERICA I-PDT for depressed adolescents – a pilot randomized controlled trial

#### Acronym

**ERICA** 

## **Study objectives**

Internet-delivered psychodynamic therapy (I-PDT) is more effective than symptom monitoring and brief support via internet with regard to reducing depressive symptoms.

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

The Regional Ethical Board in Stockholm, 13/12/2018, ref. 2018/2268-31/5.

## Study design

Interventional, randomized controlled trial, pilot study

## Primary study design

Interventional

## Study type(s)

Treatment

## Health condition(s) or problem(s) studied

Mild to moderate depression in adolescents

### **Interventions**

Directly after inclusion, participants will be randomized to one of two arms with 1:1 ratio: intervention or control condition. An independent researcher, not involved in the study, will conduct the randomisation procedure by means of a computerised random number service.

Intervention: Internet-delivered psychodynamic treatment (I-PDT) that consists of 8 therapist-supported self-help modules delivered over 8 weeks in a secure online platform. Modules consist of texts followed by assignments which they send to their therapist and receive feedback within a few days, complemented with 30 minutes chat support weekly. The aim of I-PDT is to decrease emotional avoidance and increase awareness and experience of emotions. Participants are encouraged to gradually approach previously warded off feelings. They will also be taught how to link their emotions to their symptoms. Another treatment goal is to acquire a greater capacity for anxiety regulation. Treatment duration is 8 weeks and measurements are made prior to treatment, weekly during treatment, and at termination.

Control condition: Brief supportive contact over internet over 8 weeks, with monitoring of symptoms and well-being as well as exchange of short messages between participant and therapist. Duration is 8 weeks and measurements are made prior to the supportive contact, weekly during the supportive contact, and at termination. After 10 weeks, participants in the control condition will be offered the I-PDT-program with email-support. New measurements will be made weekly during the treatment and at termination.

# Intervention Type

Behavioural

## Primary outcome(s)

1. Depressive symptoms will be measured using the Quick Inventory of Depressive Symptomatology in Adolescents (Bernstein et al., 2010) via internet delivered self-rating forms pre-treatment, weekly during treatment and post-treatment.

- 1.1. In order to fully explore trajectories of change and possible moderating and mediating factors over time, a parallel process latent growth curve strategy will be employed.
- 2. Differences in efficacy between conditions will be investigated by modelling interaction effects of group and time.

## Key secondary outcome(s))

- 1. Anxiety symptoms will be measured weekly using the Generalised Anxiety Disorder 7- item scale (GAD-7; Kroenke et al., 2010).
- 2. The severity of depression symptoms will be measured pre- and post-treatment using the Montgomery Asberg Depression Rating Scale (Svanborg & Asberg, 1994).

## Completion date

20/10/2019

# **Eligibility**

## Key inclusion criteria

- 1. Adolescents 15-18 years of age
- 2. Suffering from mild to moderate symptoms of depression according to a diagnostic interview (MINI 7.0).
- 3. Have access to a computer/smartphone/tablet with internet connection
- 4. Able to read, write and speak Swedish without the aid of an interpreter.

## Participant type(s)

Patient

# Healthy volunteers allowed

No

## Age group

Other

#### Sex

All

## Total final enrolment

76

## Key exclusion criteria

- 1. Risk of suicidality and/or earlier suicide attempts,
- 2. Partaking in other psychological treatment,
- 3. Psychotropic medication not stable since at least three months,
- 4. Other primary diagnoses in need of other treatment
- 5. Current fulfillment of any of the following diagnoses: any psychotic disorder, bipolar I/II disorder, antisocial personality disorder, and autism-spectrum disorder.
- 6. Comorbid drug or alcohol abuse.

## Date of first enrolment

03/01/2019

## Date of final enrolment

01/02/2019

# Locations

## Countries of recruitment

Sweden

SE-106 91

## Study participating centre

Department of Psychology, Stockholm University

Department of Psychology, Stockholm University Stockholm Sweden

Sponsor information

## Organisation

Department of Psychology, Stockholm University

## **ROR**

https://ror.org/05f0yaq80

# Funder(s)

# Funder type

Charity

## **Funder Name**

Kavli Trust

# **Results and Publications**

# Individual participant data (IPD) sharing plan

The data sharing plans for the current study are unknown and will be made available at a later date.

# IPD sharing plan summary

Data sharing statement to be made available at a later date

# Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient- facing?
Results article	results	30/03 /2020	01/04 /2020	Yes	No
Results article	results	14/07 /2020	10/08 /2020	Yes	No
Results article	findings from interviews	06/12 /2022	08/12 /2022	Yes	No
Results article	Sudden gains and large intersession improvements	17/08 /2020	28/02 /2024	Yes	No
Participant information sheet	Participant information sheet	11/11 /2025	11/11 /2025	No	Yes
Study website	Study website	11/11 /2025	11/11 /2025	No	Yes