

# Correlation between peripheral muscle strength and breathing tube removal outcome

<b>Submission date</b> 01/05/2019	<b>Recruitment status</b> No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 30/05/2019	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 10/08/2021	<b>Condition category</b> Other	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

Peripheral muscle weakness is common in critically ill patients with mechanical ventilator. Evidence from previous studies showed that peripheral muscle strength is associated with weaning outcome. However, the predictive value of peripheral muscle strength on extubation outcome from mechanical ventilation has not been investigated. The purpose of this study was to evaluate the relationship between peripheral muscle strength and extubation failure among patients in an intensive care unit (ICU).

### Who can participate?

ICU patients who are mechanically ventilated for more than 48 hrs and planning to wean according to standard protocol can participate.

### What does the study involve?

Evaluation of the patient's biceps and quadriceps muscle strength

### What are the possible benefits and risks of participating?

**Benefits:** We use peripheral muscle strength to predict patient extubation outcome. It might help the patient to decrease the extubation failure rate.

**Risks:** Maybe when the patient performs maximum isometric contraction, the blood pressure will increase. However, if the patient can not tolerate the test, we will stop the test and keep follow the patient's vital sign until stable

### Where is the study run from?

Landseed International Hospital, Taoyuan City, Taiwan

### When is the study starting and how long is it expected to run for?

July 2019 to June 2020

### Who is funding the study?

Landseed International Hospital, Taiwan

Who is the main contact?  
Dr Tsung-Hsien Wang  
wth75529@gmail.com

## Contact information

### Type(s)

Public

### Contact name

Dr Tsung-Hsien Wang

### ORCID ID

<http://orcid.org/0000-0001-5096-1131>

### Contact details

No.452, Huanqiu Rd  
Luzhu District  
Kaohsiung City  
Taiwan  
82144  
+886 912-956-122  
wth75529@gmail.com

## Additional identifiers

### EudraCT/CTIS number

Nil known

### IRAS number

### ClinicalTrials.gov number

Nil known

### Secondary identifying numbers

Nil known

## Study information

### Scientific Title

Peripheral muscle strength at the time of extubation may be a valuable predictor for extubation outcome

### Study objectives

Peripheral muscle strength at the time of extubation may be a valuable predictor for extubation outcome

### Ethics approval required

Old ethics approval format

### **Ethics approval(s)**

Approved 02/05/2019, Landseed International Hospital ethics institutional research committee (Pingzhen District Taoyuan City Guangtai Road 77, Taiwan), ref: IRB-19-017

### **Study design**

Prospective observational study

### **Primary study design**

Observational

### **Secondary study design**

Longitudinal study

### **Study setting(s)**

Hospital

### **Study type(s)**

Screening

### **Participant information sheet**

No participant information sheet available

### **Health condition(s) or problem(s) studied**

Mechanical ventilation

### **Interventions**

ICU patients who were mechanically ventilated for more than 48 hrs and were planning to wean according to standard protocol were enrolled in this study. Limb muscle strength was assessed using the hand-held dynamometer (MicroFET) on the day of planned extubation.

The MicroFET (Fet stands for Force Evaluating & Testing) is an electronic hand-held dynamometer which fits perfectly in the palm of the hand. It is the most cost effective and ergonomically designed hand-held dynamometer which is available on the market. It was designed to be a standalone gauge for capturing individual force measurements for any muscle test.

For testing isometric quadriceps femoris muscle force, the patient was placed in semi-Fowler position with knee extension and the transducer was placed on the anterior surface of the lower leg proximal to the ankle.

For testing isometric biceps muscle force, the patient was placed in semi-Fowler position with elbow slight flexion and the transducer was placed on the anterior surface of the wrist. Examiners demonstrated and verbally explained the task before testing. Instruction and encouragement were given to have the patient gradually apply maximum force against the transducer pad of the microFET2 over three seconds. At least three repetitions were performed until results were reproducible.

Follow-up continued until the patient was transferred to the general ward.

### **Intervention Type**

Other

**Primary outcome measure**

1. Limb muscle strength assessed using the hand-held dynamometer (MicroFET) on the day of planned extubation.
2. Extubation failure rate, defined as the need for reintubation within 72 hours after extubation (determined by the attending physician observation)
3. Extubation rate (calculated as extubation failure participants divided by all participants)

**Secondary outcome measures**

1. In-hospital mortality defined as death occurring during the hospital stay

**Overall study start date**

01/05/2019

**Completion date**

30/06/2020

## Eligibility

**Key inclusion criteria**

ICU patients who were mechanically ventilated for more than 48 hrs and were planning to wean according to standard protocol

**Participant type(s)**

Patient

**Age group**

Adult

**Sex**

Both

**Target number of participants**

60

**Total final enrolment**

52

**Key exclusion criteria**

1. Brain death
2. Ventilator dependent
3. Tracheostomy
4. Patients unable to perform the test were excluded (any rheumatologic conditions, previously known abnormal limitations of strength, amputations, muscular disease)

**Date of first enrolment**

01/07/2019

**Date of final enrolment**

30/06/2020

## Locations

### Countries of recruitment

Taiwan

### Study participating centre

#### Landseed International Hospital

Departments of Critical Care Medicine

No. 77 Guangtai Road

Pingzhen District

Taoyuan City

Taiwan

32449

## Sponsor information

### Organisation

Landseed International Hospital

### Sponsor details

No.77

Guangtai Road

Pingzhen District

Taoyuan City

Taiwan

23449

+886 (0)3 4941234

wth75529@gmail.com

### Sponsor type

Not defined

### Website

<https://ssl.landseed.com.tw/lishin/index.php>

### ROR

<https://ror.org/006arvw77>

## Funder(s)

### Funder type

Hospital/treatment centre

### Funder Name

Landseed International Hospital

## Results and Publications

### Publication and dissemination plan

Planned publication in a high-impact peer-reviewed journal

### Intention to publish date

30/12/2022

### Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request

### IPD sharing plan summary

Available on request

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>		09/08/2021	10/08/2021	Yes	No