Proinsulin in gestational diabetes mellitus

Submission date	Recruitment status No longer recruiting	[X] Prospectively registered		
03/10/2017		[X] Protocol		
Registration date	Overall study status Completed	Statistical analysis plan		
11/10/2017		[X] Results		
Last Edited	Condition category	[] Individual participant data		
09/09/2025	Pregnancy and Childbirth			

Plain English summary of protocol

Background and study aims

In pregnancy, some women are unable to process glucose (sugar) properly - this is referred to as gestational diabetes mellitus. It is a condition that is present only in pregnancy and the glucose levels usually return to normal after the birth of the baby. Gestational diabetes mellitus is linked with an increased risk of complications to both the baby (large baby, shoulder getting stuck in the birth canal and stillbirth) and the mother (prolonged labour). It is therefore important to screen and diagnose the condition, and early diagnosis and management has been shown to reduce the complications by up to 60%. Currently, at 24-28 weeks women with a risk factor (body mass index greater than 30 kg/m2, family history of diabetes in a first degree relative, previous history of diabetes or previous large baby i.e. birth weight >4.5 kg) are offered an oral glucose tolerance test (a glucose drink) with blood samples taken before and 2 hours after the drink. In this study the same test (oral glucose tolerance test) is done but at an earlier stage of pregnancy (at 16-18 weeks) in addition to the recommended test at 24-28 weeks to see if this new test is able to diagnose the condition at an earlier stage in pregnancy. Additional blood samples are also taken for measurement of beta cell function. The beta cells of the pancreas are responsible for controlling blood sugar by secretion of insulin. They also secrete very small amounts of proinsulin (which is normally converted to insulin inside the beta cells). Proinsulin is normally released in small quantities but concentrations increase with the onset of diabetes before glucose levels rise. The aim of this study is to find out whether measurement of the concentration of proinsulin provides an earlier and accurate means of identifying gestational diabetes mellitus.

Who can participate?

Pregnant women at least 18 years of age at 16-18 weeks of pregnancy with any one of the following: body mass index greater than 30 kg/m2, family history of diabetes in a first degree relative, previous history of diabetes or previous large baby i.e. birth weight >4.5 kg

What does the study involve?

This study involves three visits for participants at different stages of pregnancy. At visit 1 (16-18 weeks of pregnancy) participants fast overnight and an oral glucose tolerance test is conducted, with blood samples taken at 0, 30, 60 and 120 minutes after consumption of a glucose drink. Visit 2 (24 - 28 weeks of pregnancy) is the same as visit 1 and includes an oral glucose tolerance test after an overnight fast. At visit 3 (6 weeks after delivery) participants are asked to attend after fasting overnight and have a fasting blood sample taken. Additional data are collected

including birth weight of the baby, Apgar score at birth and medications including insulin doses before birth.

What are the possible benefits and risks of participating?

If this test does prove useful, in future pregnant women can potentially be diagnosed with gestational diabetes mellitus at an earlier stage, reducing the risk of complications. It is not expected that participating in this study will cause any problems, but as blood samples are taken participants may experience bleeding, discomfort, bruising, infection or inflammation where the blood is taken or dizziness and feeling faint.

Where is the study run from? Swansea University (UK)

When is the study starting and how long is it expected to run for? October 2017 to August 2020

Who is funding the study?
Abertawe Bro Morgannwg University Health Board (UK)

Who is the main contact?
Sarah Dowrick
sarah.c.dowrick@swansea.ac.uk

Contact information

Type(s)

Public

Contact name

Miss Sarah Dowrick

Contact details

Diabetes Research Unit Cymru Grove Building Swansea University Swansea United Kingdom SA2 8PP +44 (0)1792 602590 sarah.c.dowrick@swansea.ac.uk

Additional identifiers

Protocol serial number

1.0

Study information

Scientific Title

Proinsulin in the diagnosis and risk stratification of gestational diabetes mellitus

Study objectives

The primary objective of this study is to test the hypothesis that fasting proinsulin measurements at 16-18 weeks gestation will discriminate/risk stratify gestational diabetes from women with normal glucose tolerance.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Wales REC 6, 06/07/2017, ref: 17/WA/0194

Study design

Observational cross-sectional pilot study

Primary study design

Observational

Study type(s)

Diagnostic

Health condition(s) or problem(s) studied

Gestational diabetes mellitus

Interventions

This study will involve three visits for participants, at different stages of gestation.

Visit 1 (16-18 weeks gestation): participants will be asked to attend at 16-18 weeks gestation, having fasted overnight. An oral glucose tolerance test will be conducted, with blood samples taken at 0, 30, 60 and 120 mins following consumption of a glucose drink.

Visit 2 (24 - 28 weeks gestation): this will be the same as visit one and include an oral glucose tolerance test after an overnight fast.

Visit 3 (6 weeks post delivery): participants will be asked to attend having fasted overnight and will have a fasting blood sample taken. Additional data including birth weight of the baby, Apgar score at birth and medications including insulin doses prior to birth will be collected.

Intervention Type

Other

Primary outcome(s)

Circulating concentration of fasting total and intact proinsulin at 16-18 weeks gestation

Key secondary outcome(s))

- 1. Diagnosis of gestational diabetes mellitus as established by glucose measurements following an oral glucose tolerance test at 24-28 weeks
- 2. Need for insulin during the pregnancy obtained from clinical notes
- 3. Relationship between individual risk factors and proinsulin for the development of gestational diabetes mellitus, measured using logistic regression at 16-18 weeks

Completion date

01/08/2020

Eligibility

Key inclusion criteria

Pregnant women at least 18 years of age at 16-18 weeks gestation with any one of the following:

- 1. BMI >30kg/m2
- 2. Previous macrosomic baby (>4.5kg)
- 3. Previous gestational diabetes mellitus
- 4. Family history of type 2 diabetes mellitus (first degree relative with diabetes)

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

Female

Total final enrolment

83

Key exclusion criteria

- 1. Subjects unable or unwilling to sign informed consent
- 2. Known previous diabetes mellitus or treatment with metformin
- 3. Known chronic infection like Hepatitis or HIV or chronic kidney, liver or heart disease
- 4. Previous bariatric surgery

Date of first enrolment

16/10/2017

Date of final enrolment

01/08/2020

Locations

Countries of recruitment

United Kingdom

Wales

Study participating centre

Joint Clinical Research Facility

Institute of Life Sciences 2 Swansea University Swansea United Kingdom SA2 8PP

Sponsor information

Organisation

Abertawe Bro Morgannwg University Health Board

ROR

https://ror.org/04zet5t12

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Abertawe Bro Morgannwg University Health Board

Results and Publications

Individual participant data (IPD) sharing plan

The data sharing plans for the current study are unknown and will be made available at a later date.

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		05/09/2025	09/09/2025	Yes	No
Protocol article	protocol	29/08/2018	21/10/2019	Yes	No
HRA research summary			28/06/2023		No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes