# Translating evidence for early intervention in psychosis (TRANSLATE) in low and lower-middle countries (LMIC): implementation and evaluation

| Submission date   | <b>Recruitment status</b> Recruiting | <ul><li>Prospectively registered</li></ul> |  |
|-------------------|--------------------------------------|--|--|
| 09/09/2025        |                                      | ☐ Protocol                                 |  |
| Registration date | Overall study status                 | Statistical analysis plan                  |  |
| 22/10/2025        | Ongoing                              | Results                                    |  |
| Last Edited       | Condition category                   | Individual participant data                |  |
| 22/10/2025        | Mental and Behavioural Disorders     | [X] Record updated in last year            |  |

## Plain English summary of protocol

Background and study aims

This study is about helping young people in Pakistan and Sri Lanka who are experiencing psychosis for the first time. Psychosis is a mental health condition that affects how people think, feel, and behave. In many low- and middle-income countries, people often wait a long time before getting treatment—sometimes over two years. This delay can seriously affect their quality of life, education, and ability to work.

In countries like the UK, early intervention services help people get treatment quickly, which leads to better outcomes. This study aims to set up similar early intervention services in Pakistan and Sri Lanka. It also wants to understand why some people don't respond well to treatment (a condition called treatment resistant schizophrenia) and whether we can predict who might be at risk.

#### Who can participate?

People in Pakistan and Sri Lanka who are experiencing psychosis for the first time may be invited to take part in the study. The research team will work with local mental health services to identify potential participants.

#### What does the study involve?

Participants will receive care through the new early intervention services. The team will collect information about their health, treatment, and progress over time. Some participants may also be asked to provide additional information to help researchers understand why some people don't respond to treatment.

What are the possible benefits and risks of participating?

The main benefit is receiving care earlier than usual, which can lead to better recovery and

improved quality of life. There may be some risks, such as feeling uncomfortable when answering personal questions or sharing health information, but the research team will take steps to protect participants' privacy and wellbeing.

Where is the study run from? Keele University (UK)

When is the study starting and how long is it expected to run for? April 2024 to February 2029

Who is funding the study? National Institute for Health and Care Research (NIHR) (UK).

Who is the main contact? s.farooq@keele.ac.uk n.wellappuli@keele.ac.uk h.n.a.fonseka@keele.ac.uk

## Contact information

#### Type(s)

Principal investigator

#### Contact name

Prof Saeed Faroog

#### **ORCID ID**

https://orcid.org/0000-0002-6910-3003

#### Contact details

School of Medicine, University of Keele, Keele Newcastle under Lyme United Kingdom ST5 5GB +44 7958012102 s.farooq@keele.ac.uk

#### Type(s)

Scientific

#### Contact name

Dr Nalinda Wellappuli

#### **ORCID ID**

https://orcid.org/0000-0001-6934-2226

#### Contact details

School of Medicine, University of Keele, Keele Newcastle under Lyme United Kingdom ST5 5GB +44 7405862955 n.wellappuli@keele.ac.uk

## Type(s)

**Public** 

#### Contact name

Dr Nishani Fonseka

#### **ORCID ID**

https://orcid.org/0000-0001-5955-2211

#### Contact details

School of Medicine, University of Keele, Keele Newcastle under Lyme United Kingdom ST5 5GB +44 7902549258 h.n.afonseka@keele.ac.uk

## Additional identifiers

#### Clinical Trials Information System (CTIS)

Nil known

## ClinicalTrials.gov (NCT)

Nil known

#### Protocol serial number

Nil known

# Study information

#### Scientific Title

Translating evidence for Early Intervention in Psychosis (TRANSLATE) in Low and Lower-Middle Countries (LMIC): Implementation and Evaluation

## **Acronym**

**TRANSLATE** 

## Study objectives

- 1. To evaluate the implementation of Early Intervention in Psychosis (EIP) services in maintaining engagement with the services, achieving remission in First Episode of Psychosis, and other relevant implementation outcomes.
- 2. To assemble a cohort of FEP within the EIP service and identify potential predicting factors of Treatment Resistant Schizophrenia at one-year follow-up.
- 3. To develop a prognostic model for estimating an individual's risk of treatment resistance at one year and to undertake the validation of the model's predictive performance.

#### Ethics approval required

#### Ethics approval required

#### Ethics approval(s)

1. approved 11/06/2025, Keele University's Research Ethics Committee (Keele University, Keele, Newcastle Under Lyme, ST5 5GB, United Kingdom; +44 1782 733937; health.ethics@keele.ac.uk), ref: 1028

2. approved 28/03/2025, Kyber Medical University - Institute of Public Health and Social Sciences (KMU, Phase-5, Hayatabad, -, Pakistan; +91-5892867; drshaista.iph@kmu.edu.pk), ref: KMU/IPHSS /Ethics/2025/TE/260

#### Study design

Observational cross sectional cohort study

#### Primary study design

Observational

## Study type(s)

Other

#### Health condition(s) or problem(s) studied

First episode psychosis and treatment resistant schizophrenia

#### **Interventions**

Patients with a diagnosis of First episode Psychosis (FEP) will be recruited. The participants will be assessed at the baseline and follow up in 1, 3,6, and 12-month period with pre identified tools. All participants will receive care following the clinical guidelines developed for the management of FEP in the study settings. The package of care will include evidence-based pharmacotherapy, cognitive behaviour therapy and other psychosocial interventions. The treatment will be provided by trained psychiatrists and psychologists. The supervising psychiatrist will decide the choice of medication, dosage and other interventions after involving patients and families in the treatment according to the clinical practice guidelines adopted for management of First Episode Psychosis in Pakistan and Sri Lanka. Each participant will be allocated a care coordinator to coordinate the services to ensure the receipt of services that prescribed to given for each patient. The patients will be followed up for one year to estimate the rates of remission, engagement with the service and other clinical outcomes during the study.

The duration of observation and total duration of follow-up will be 12 months for each participant. This study is not aimed at testing the efficacy related to the individual drugs or other interventions, these are well established. Our aim is to assess the implantation effectiveness of the Early Intervention in Psychosis services in resource poor settings.

#### Intervention Type

Other

## Primary outcome(s)

- 1. Disengagement from service is measured using clinic records and care coordinator notes at 1,
- 3, 6, and 12 months
- 2. Remission is measured using PANSS (score  $\leq$ 3) or YMRS (score  $\leq$ 12) at 1, 3, 6, and 12 months

## Key secondary outcome(s))

- 1. Change in occupational and social functioning is measured using WHODAS 2.0 at 1, 3, 6, and 12 months
- 2. Psychopathology symptom severity is measured using PANSS, YMRS, and HDRS at 1, 3, 6, and 12 months
- 3. Family burden is measured using the Family Burden Scale at 3 and 12 months
- 4. Perceived stigma is measured using the Modified Internalized Stigma of Mental Illness tool (ISMI) at 12 months
- 5. Blood pressure is measured using clinic records at 1, 3, 6, and 12 months
- 6. BMI is measured using clinic records at 1, 3, 6, and 12 months
- 7. Cardiovascular disease risk is measured using WHO-STEPS at 12 months
- 8. Appropriateness is measured using the Intervention Appropriateness Measure tool (IAM) at 12 months
- 9. Feasibility is measured using the Feasibility Intervention Measure at 12 months
- 10. Acceptability is measured using the Applied Mental Health Research Group tool at 12 months
- 11. Fidelity of CBT delivery is measured using the Revised Cognitive Therapy Scale (CTS-R) at 12 months
- 12. Implementation cost is measured using administrative data logs collected over the one-year period

#### Completion date

01/02/2029

# Eligibility

## Key inclusion criteria

- 1. Patients aged 18-35 years and
- 2. Residing within the study districts
- 3. Diagnosed with First-ever psychotic episode who have not received antipsychotic medication previously, or if they already have used antipsychotic medications, it was for no longer than six weeks

## Participant type(s)

**Patient** 

## Healthy volunteers allowed

No

## Age group

Adult

## Lower age limit

18 years

## Upper age limit

35 years

#### Sex

All

#### Key exclusion criteria

- 1. Those with an overt learning disability
- 2. Those with severe substance abuse (except nicotine dependence)
- 3. Those with organic illness associated with psychotic symptoms

#### Date of first enrolment

01/09/2025

#### Date of final enrolment

31/01/2029

## Locations

#### Countries of recruitment

Pakistan

Sri Lanka

#### Study participating centre

Deaprtment of Psychiatry, Ayub Medical Teaching Institution

Main Mansehra Road Abbottabad Pakistan Kyber Pakhtunkhwa Province

## Study participating centre

## Deapartment of Psychiatry and Behavioural Sciences

Faisalabad Medical University, Allied/ District Head Quarters Hospital -Faisalabad, Mall Road Faisalabad
Pakistan
38000

#### Study participating centre

Department of Psychiatry, Liaquat University of Medical and Health Sciences, Jamshoro, Hyderabad

Administration Block, Deh Soun Valhar Jamshoro, Hyderbad Pakistan 76090

## Study participating centre

Department of Psychiatry, Balochistan Institute of Psychiatry and Behavioural Sciences (BIPBS) , Quetta

Balochistan Institute of Psychiatry and Behavioral Sciences (BIPBS), Behind BMC Hospital, Berwery Road Quetta Pakistan 00

Study participating centre
Department of Psychiatry, Medical Teaching Institute, Lady Reading Hospital, Peshawar
Soekarno Rd, Pipal Mandi
Peshawar
Pakistan
25000

## Study participating centre

Institute of Public Mental Health and Behavioural Sciences, Khyber Medical University, Peshawar KMU Main Campus, Phase 5, Hayatabad Peshawar Pakistan Kyber Pakhtunkhwa Province

Study participating centre National Hospital Galle Karapitiya Galle Sri Lanka 00

# Sponsor information

## Organisation

**Keele University** 

#### **ROR**

https://ror.org/00340yn33

# Funder(s)

## Funder type

Government

#### Funder Name

National Institute for Health and Care Research

#### Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

#### **Funding Body Type**

Government organisation

## Funding Body Subtype

National government

#### Location

**United Kingdom** 

## **Results and Publications**

#### Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be available upon request from Professor Saeed Farooq - s.farooq@keele.ac.uk

## IPD sharing plan summary

Available on request

## **Study outputs**

| Output type                   | Details                       | Date created | Date added Peer reviewed? | Patient-facing? |
|-------------------------------|-------------------------------|--------------|---------------------------|-----------------|
| Participant information sheet | Participant information sheet | 11/11/2025   | 11/11/2025 No             | Yes             |
| Study website                 | Study website                 | 11/11/2025   | 11/11/2025 No             | Yes             |