

# A study comparing treatments for severe toothache in adult teeth: Full Pulpotomy or Root Canal Treatment

<b>Submission date</b> 20/03/2023	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 12/07/2023	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 12/07/2023	<b>Condition category</b> Oral Health	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Tooth decay is very common; almost all adults have some decay in their back teeth. Treating decay costs the NHS over £3.4 billion each year. Where the tooth decay has spread deep into the nerve, and caused irreversible damage, there are different treatments available. The most common treatments are to take out the tooth or use Root Canal Treatment. Root Canal Treatment removes all the nerve and is very expensive, time consuming and technically demanding. A possible alternative is to remove only the infected part of the nerve, which is an easier and quicker treatment. This technique is called Pulpotomy.

Research has already been conducted in children's teeth that supports using pulpotomy instead of Root Canal Treatment. However, there are gaps in our knowledge about using Pulpotomy in adult's teeth.

The aim of this study is to compare Pulpotomy with root canal treatment in adult patients. We will consider the pain experienced by patients, the impact of treatment on their lives, whether further treatment is needed due to infection and the time and financial costs incurred.

### Who can participate?

NHS adult (16 years or older) dental patients who have irreversible damage of the nerve of a tooth in adult back teeth and who agree to take part.

### What does the study involve?

Participants will be randomly allocated to receive one of the two treatments (Root Canal Treatment or Pulpotomy).

We will measure how well the treatments work over one year. We will look for evidence of treatment success and compare symptoms such as swelling. We will measure if the tooth needs further treatment, for example, a new or repeat Root Canal Treatment or if the tooth needs to be extracted. We will also measure patients' pain and anxiety, how their oral health affects their quality of life, treatment costs and satisfaction with their dental treatment.

We will use patient questionnaire surveys, X-rays and data from dentists to evaluate the success of the two treatments. We will also interview dental staff and patients about their views on the two treatments.

What are the possible benefits and risks of participating?

Root Canal Treatment is the standard procedure used by dentists to treat Irreversible Pulpitis, but in some cases, the Full Pulpotomy Treatment may be quicker and less invasive than Root Canal Treatment. Both treatments are an alternative to having an extraction. As with all treatments, whichever treatment participants have, there is a small risk that it may not work, and further treatment may be required. Further treatment for Full Pulpotomy could be a Root Canal Treatment or having the tooth removed. Both treatments include having X-rays taken. This is part of routine care. Dental radiographs need very low doses of radiation and are equivalent to less than a day of background radiation. We do not think that there are any possible disadvantages for those that this treatment is suitable for, as they will require some form of treatment for the condition.

Where is the study run from?

University of Dundee (UK)

When is the study starting and how long is it expected to run for?

June 2020 to December 2025

Who is funding the study?

The National Institute for Health Research (NIHR) Health Technology Assessment (HTA) Programme (UK)

Who is the main contact?

Lori Brown, Study Administrator  
l.v.brown@dundee.ac.uk

## Contact information

### Type(s)

Public

### Contact name

Mrs Lorna Macpherson

### ORCID ID

<https://orcid.org/0000-0003-1897-923X>

### Contact details

Dundee Dental Hospital & Research School  
Level 9 DHSRU Park Place  
Dundee  
United Kingdom  
DD1 4HN  
+44 7595428209  
l.e.macpherson@dundee.ac.uk

## Additional identifiers

### Clinical Trials Information System (CTIS)

Nil known

## **Integrated Research Application System (IRAS)**

323138

## **ClinicalTrials.gov (NCT)**

Nil known

## **Protocol serial number**

IRAS 323138, Sponsor ID 2-054-22, CPMS 56091

# **Study information**

## **Scientific Title**

Pulpotomy or root canal treatment for the management of irreversible pulpitis in mature teeth

## **Acronym**

PIP Study

## **Study objectives**

Tooth decay is very common; almost all adults have some decay in their back teeth. This costs the NHS over £3.4 billion each year. Where tooth decay has spread deep into the nerve, and caused serious and irreversible damage, there are different treatments available. The most common treatments are to take out the tooth or have root canal treatment (RCTx). RCTx removes all the nerve, is expensive, time consuming and technically demanding. A possible alternative is to remove only the infected part of the nerve, which could be an easier and quicker treatment. This is called pulpotomy.

Research in children's teeth supports using pulpotomy instead of root canal treatment.

However, there are gaps in our knowledge about using pulpotomy in adult teeth. Following a feasibility study, the PIP Study aims to provide this knowledge.

The PIP Study compares pulpotomy with root canal treatment in adult patients.

## **Ethics approval required**

Ethics approval required

## **Ethics approval(s)**

approved 13/06/2023, West Midlands - Coventry & Warwickshire Research Ethics Committee (The Old Chapel, Royal Standard Place, Nottingham, NG1 6FS, United Kingdom; +44 207 104 8013; coventryandwarwick.rec@hra.nhs.uk), ref: 23/WM/0094

## **Study design**

Interventional randomized controlled trial

## **Primary study design**

Interventional

## **Study type(s)**

Treatment

## **Health condition(s) or problem(s) studied**

Pulpotomy or root canal treatment for the management of irreversible pulpitis in mature teeth

## **Interventions**

We will recruit 530 NHS adult dental patients who have irreversible damage of the nerve in a suitable tooth who agree to take part. They will receive one of the two treatments (Root Canal Treatment or Full Pulpotomy) at random, using a computerised randomisation tool that dentists will access using the online study database.

Root Canal Treatment is the current treatment available on the NHS to try and save your tooth. It aims to remove all of the nerve from the tooth and replace it with a filling material. An alternative treatment for this kind of toothache is called a Full Pulpotomy. A Full Pulpotomy removes only the top portion of the nerve which includes the damaged part caused by decay but aims to keep the lower (healthy) portion alive. Full Pulpotomy is already used on children's teeth and on adult teeth when they are being treated in dental hospitals. However, it is not yet routinely used on adult teeth in NHS dental practices.

The treatments will take a varying amount of time to complete depending on individual circumstances, however a Full Pulpotomy usually takes 1-2 appointments and a Root Canal Treatment can take 3-4 appointments. The duration of the appointment will vary.

We will measure how well the treatments work over one year, with participant questionnaires completed at baseline, 7 days and also at 12 months. We will look at treatment success, levels of pain and if the tooth needs more treatment using CRFs as well as the questionnaires. We will measure patients' anxiety, how their oral health affects their quality of life, treatment costs and satisfaction. X-rays taken at baseline, after treatment and at 12-months along with data from dentists to compare the two treatments. We will also interview dental staff and patients.

## **Intervention Type**

Procedure/Surgery

## **Primary outcome(s)**

1. Clinical success of Full Pulpotomy with Root Canal Treatment (no re-intervention or symptoms of pulpitis or apical periodontitis) at 1 year. Assessed using radiograph at 1-year post-randomisation: Success is the absence of apical radiolucency in the periapical radiograph following participant's initial intervention
2. Pain is measured using the 10-point visual analogue score (VAS) at baseline, 7 days and 12 months
3. Oral Health Related Quality of Life will be assessed at 7 days and 1 year using the Oral Health Impact Profile 14 (OHIP-14)
4. Health status will be assessed using the generic EQ5D at 7 days and 1 year, consisting of five dimensions of HRQoL
5. Dental anxiety will be assessed using the validated MDAS criteria at 1 year
6. Patient Satisfaction will be assessed using the NHS England Commissioning of Dental Services Guidelines Patient Reported Experience Measures (PREMs) at 1 year

## **Key secondary outcome(s)**

1. Radiographic success (absence of radiolucency in participant's initial intervention)
2. Patient health related quality of life (OHIP-14) at 1 year
3. Dental pain (10-point visual analogue score (VAS)) at 1 year
4. Dental anxiety (assessed using the validated MDAS criteria) at 1 year
5. Satisfaction with care/treatment and adverse events at 1 year in 12-month questionnaire and 12 month follow up CRF
6. Patient health related quality of life and dental pain at 7 days measured using VAS and OHIP-

14 in 7-day questionnaire

7. General population preferences for treatment and outcomes and predict uptake, NHS and patient perspective costs; incremental net benefit over the study follow up; and incremental cost per QALY over the 1-year follow-up and extrapolated over a modelled lifetime horizon, measured using the 12-month questionnaire, follow up CRF at 12-months and participant interviews.

8. To explore the implementation of technologies and the mechanisms of impact including acceptability of interventions to patients and clinicians by interviews at 1 year

**Completion date**

31/12/2025

## Eligibility

**Key inclusion criteria**

Adults (16 years and older) with symptoms indicative of irreversible pulpitis (as defined by the European Society of Endodontology) in a premolar or molar tooth with deep caries and or a deep restoration.

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Sex**

All

**Key exclusion criteria**

1. Tooth with immature roots, clinical or radiographic signs of a necrotic pulp, or a poor prognosis (e.g., internal or external resorption)
2. Presence of a sinus, tenderness to percussion, buccal tenderness, pathological mobility or evidence of pathology on a periapical radiograph
3. Insufficient tooth tissue for a restoration
4. All treatment delivered under a private contract

**Date of first enrolment**

01/05/2023

**Date of final enrolment**

31/10/2024

## Locations

**Countries of recruitment**

United Kingdom

England

Scotland

**Study participating centre**

**Dundee Dental Hospital & Research School**

Level 9

DHSRU

Park Place

Dundee

United Kingdom

DD1 4HN

## **Sponsor information**

**Organisation**

University of Dundee

**ROR**

<https://ror.org/03h2bxq36>

## **Funder(s)**

**Funder type**

Government

**Funder Name**

Health Technology Assessment Programme

**Alternative Name(s)**

NIHR Health Technology Assessment Programme, Health Technology Assessment (HTA), HTA

**Funding Body Type**

Government organisation

**Funding Body Subtype**

National government

**Location**

United Kingdom

# Results and Publications

## Individual participant data (IPD) sharing plan

The data-sharing plans for the current study are unknown and will be available at a later date.

## IPD sharing plan summary

Data sharing statement to be made available at a later date

## Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Study website</a>	Study website	11/11/2025	11/11/2025	No	Yes