

# Can supplementation with a probiotic reduce the crying time and modify the gut microbiome in a group of infants with colic?

<b>Submission date</b> 14/05/2020	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 16/05/2020	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 02/12/2020	<b>Condition category</b> Signs and Symptoms	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

Colic is when a baby cries a lot and is difficult to console, without an obvious cause. This is a common problem that should get better on its own, but can be distressing and exhausting for parents. The exact causes of infantile colic are not fully understood. Likely causes could be problems with gut motility and food intolerance. Growing evidence suggests that the gut microbiota (the microorganisms that live in the gut) could be also involved in colic.

Certain compositions of the gut microbiota, where a type of bacteria called Coliforms are in high concentrations and a type of bacteria called Lactobacilli are in a low concentration, have been found to be more common in infants with colic than in infants without colic.

Probiotic supplementation with Lactobacilli has been shown to be effective in treating this condition in some clinical trials of breastfed infants. However, changing the composition of the gut bacteria during the early stages of gut development can induce lasting shifts in immunity as a result of inflammation of the gut.

A specific bacterial species called *Lactobacillus rhamnosus* may have anti-inflammatory properties. Supplementation with a probiotic of this bacteria may improve colic and reduce inflammation. The aim of this study was to investigate how effective a *Lactobacillus rhamnosus* probiotic was in treating infantile colic, in addition to maternal diet avoidance of cow's milk.

### Who can participate?

Newborns with colic aged between 2 and 10 weeks who are breastfed.

### What does the study involve?

Participants are randomly allocated to one of two groups. Infants will either have normal care and breastfeeding or they will take 5 drops of the probiotic each day for 28 days and their mother will avoid consuming cow's milk over this period. Fecal samples will be collected from nappies before the trial starts and after 28 days. Their mother will also report how often the infant is crying at the start and end of the study.

What are the possible benefits and risks of participating?

This probiotic is well known and safe. There could be a reduction in colic symptoms of crying and fussing of the baby and parental discomfort.

Where is the study run from?

University Hospital City of Health and Science of Turin (Italy)

When is the study starting and how long is it expected to run for?

April 2017 to October 2019

Who is funding the study?

Office of the Governor (USA)

Who is the main contact?

Dr Francesco Savino

francesco.savino@unito.it

### **Study website**

<https://www.cittadellasalute.to.it/>

## **Contact information**

### **Type(s)**

Scientific

### **Contact name**

Dr Francesco Savino

### **ORCID ID**

<https://orcid.org/0000-0001-9163-1016>

### **Contact details**

Department of Pediatrics

Ospedale Infantile Regina Margherita

Azienda Ospedaliera Città' Della Salute E Della Scienza Di Torino

Piazza Polonia, 94

Italy

Italy

10126

+39 (0)113135618

francesco.savino@unito.it

## **Additional identifiers**

### **EudraCT/CTIS number**

2020-002496-35

### **IRAS number**

### **ClinicalTrials.gov number**

Nil known

## **Secondary identifying numbers**

Nil known

# **Study information**

## **Scientific Title**

Supplementation with the probiotic L rhamnosus GG to reduce the crying time and modify gut microbiota, or fecal calprotectin values in a group of infants with colic (Lactobacillus GG): a randomized controlled pilot study

## **Acronym**

Lactobacillus GG

## **Study objectives**

The aim of this study is to investigate whether supplementation with the probiotic Lactobacillus rhamnosus ATCC 53103 and maternal dietary elimination of cow's milk is effective in treating infantile colic of breastfed infants and reducing gut inflammation by measuring fecal calprotectin levels and composition of gut microbiota compared to placebo control.

## **Ethics approval required**

Old ethics approval format

## **Ethics approval(s)**

Approved 05/04/ 2017, Ethics Committee of the University Hospital City of Health and Science of Turin (Ospedale Mauriziano di Torino , Corso Spezia 60, Torino 10126 Italy; +39 (0)113134079), ref: 7/2017

## **Study design**

Randomised controlled trial

## **Primary study design**

Interventional

## **Secondary study design**

Randomised controlled trial

## **Study setting(s)**

Hospital

## **Study type(s)**

Treatment

## **Participant information sheet**

See additional file in Italian ISRCTN16554977\_PIS\_Italian (added 08/06/2020)

## **Health condition(s) or problem(s) studied**

Infantile colic

## Interventions

At enrollment, each infant underwent a medical examination by a pediatrician, and parents completed a questionnaire to obtain data concerning type of delivery, birth weight, and gestational age. The pediatrician asked the parents to report full-force crying by means of a well-validated three-day diary before the faecal sample collection at baseline and again after the 28 day intervention. The mean daily minutes of crying were calculated as a sum of crying, fussing, and unsuitable crying using a parental diary.

Eligible participants will be randomized to either receive the intervention or placebo control:

1. The intervention arm received *Lactobacillus rhamnosus* GG ( $5 \times 10^9$  colony units per day). The probiotic study product consisted of a suspension of freeze-dried *L. rhamnosus* ATCC53103 in a mixture of mais oil and mono and diglyceride oil supplied in a 5 mL dark bottle fitted with a dropper cap.
2. The control arm received a placebo. The placebo was a mixture of mais oil and mono and diglyceride oil supplied in a 5 mL dark bottle fitted with a dropper cap, without probiotic.

Both arms were instructed to take 5 drops of either the probiotic or placebo in the morning 30 minutes before feeding each day for 28 days. For both arms, maternal avoidance of cow's milk was requested for the 28 days study period.

Randomization was performed by the random-digit method, on the basis of a computer-generated numbers. A two treatments randomization scheme with random block of varying size was used (Stata Statistical Software: Release 9. StataCorp LP, College Station, TX. Ralloc Procedure).

## Intervention Type

Supplement

## Primary outcome measure

Colic measured by changes in the parent-reported crying time per day of infants at baseline and 28 days

## Secondary outcome measures

Gut inflammation assessed using real-time PCR analysis of gut microbiota from fecal samples, and fecal calprotectin levels measured via a quantitative assay at baseline and 28 days

## Overall study start date

05/04/2017

## Completion date

30/11/2019

## Eligibility

### Key inclusion criteria

1. Aged between 2 and 10 weeks
2. Gestational age between 37 and 40 weeks
3. Birth weight between 2700 g and 4200 g,
4. Exclusively breastfed
5. Colic, as diagnosed according to the Wessel definition modified by Roma IV criteria (infants have at least three episodes of unexplained full-force crying lasting more than three hours per

day on at least three days a week for at least one week)  
6. Maternal willingness and ability to avoid dietary cow's milk

**Participant type(s)**

Patient

**Age group**

Neonate

**Sex**

Both

**Target number of participants**

20

**Total final enrolment**

57

**Key exclusion criteria**

1. Clinical evidence of chronic illness or gastrointestinal disorders
2. Antibiotic or probiotic treatment in the week preceding recruitment

**Date of first enrolment**

01/03/2018

**Date of final enrolment**

31/10/2019

**Locations****Countries of recruitment**

Italy

**Study participating centre**

**Ospedale Infantile Regina Margherita (ORIM)**

Children's Hospital Regina Margherita

University Hospital City of Health and Science of Turin

Piazza Polonia, 94

Torino

Italy

10126

**Sponsor information****Organisation**

Azienda Ospedaliera Citta' Della Salute E Della Scienza Di Torino

**Sponsor details**

Piazza Polonia, 94

Torino

Italy

10126

+39 (0)113135059

fsavino@cittadellasalute.to.it

**Sponsor type**

Hospital/treatment centre

**Website**

<https://www.cittadellasalute.to.it/>

**ROR**

<https://ror.org/001f7a930>

## Funder(s)

**Funder type**

Government

**Funder Name**

Office of the Governor

**Alternative Name(s)****Funding Body Type**

Government organisation

**Funding Body Subtype**

Local government

**Location**

United States of America

## Results and Publications

**Publication and dissemination plan**

Planned publication in a high-impact peer-reviewed journal.

**Intention to publish date**

01/06/2020

## Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be available upon request from Dr Francesco Savino, francesco.savino@unito.it.

## IPD sharing plan summary

Available on request

## Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Participant information sheet</a>			08/06/2020	No	Yes
<a href="#">Results article</a>	results	05/06/2020	02/12/2020	Yes	No