

Synthetic versus autologous reconstruction of the medial patellofemoral ligament

Submission date 20/10/2016	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 03/03/2017	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 14/03/2023	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

The medial patellofemoral ligament (MPFL) helps to keep the kneecap in the right position. If the MPFL is torn (ruptured), the knee cap is displaced from its normal position. There are many ways to address the unstable knee cap. One such way is to operate, and there are different procedures that can be performed. One method is to use the patient's own hamstring tendons to restore the knee cap (autologous hamstring graft). An alternative is to use a synthetic mesh to restore stability. Both methods have been successfully used, with good results. However, no study has been done to determine if one method is better than the other. The aim of this study is to address this issue, and compare the short-term results using both of these methods.

Who can participate?

Patients aged 14-25 with MPFL rupture. Only patients with 2 or more dislocations are considered for inclusion.

What is involved from you as a participant?

Participants are randomly allocated to one of two groups. One group undergoes MPFL reconstruction surgery with an autologous hamstring graft. The other group undergoes the same surgery but with a synthetic ligament instead. Participants attend outpatient appointments to track their recovery. Initial follow up is performed at 6 weeks after the operation as per routine follow up. Participants' knees are assessed at 3, 6, 12, 18 and 24 months after the operation by a physiotherapist.

What are the benefits and risks of participating?

The information gathered may help assist and guide what surgical method we use and how we subsequently treat patients. The risks of surgery are no different between the groups, as the surgery is almost identical. Using the patient's own hamstring tendons may slightly increase the duration of the surgery and may result in some additional bruising.

Where is the study run from?

Altnagelvin Area Hospital (UK)

When is the study starting and how long is it expected to run for?
September 2016 to January 2022 (updated 11/08/2020, previously: July 2022)

Who is funding the study?
Western Health and Social Care Trust (UK)

Who is the main contact?
Mr Danny Acton

Contact information

Type(s)
Scientific

Contact name
Mr Danny Acton

Contact details
c/o Miss Ellen Kelly
Altnagelvin Area Hospital
Glenshane Road
Londonderry
United Kingdom
BT47 6SB
+44 (0)287 134 5171
danny.acton@westerntrust.hscni.net

Additional identifiers

Clinical Trials Information System (CTIS)
Nil known

Integrated Research Application System (IRAS)
211907

ClinicalTrials.gov (NCT)
Nil known

Protocol serial number
IRAS Project ID 211907

Study information

Scientific Title
Synthetic versus autologous reconstruction (Syn-VAR) study for recurrent patellar instability: a prospective randomised pilot study

Study objectives

To compare autologous hamstring medial patellofemoral ligament (MPFL) reconstruction with vastus medii obliquus (VMO) advancement against synthetic graft MPFL reconstruction with VMO advancement.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Office for Research Ethics Committees of Northern Ireland, 21/08/2017, ref: ORECNI ID 17/NI/0129

Study design

Randomised control prospective single-centre study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Recurrent patellar instability with MRI-proven MPFL injury

Interventions

Patients will be randomised to receive either:

1. MPFL reconstruction with autologous hamstring graft (h-MPFL)
2. Synthetic woven ligament (s-MPFL)

Both are in combination with VMO advancement. Once randomised, the patients will have the procedure explained, and have the option to proceed, or opt out pre-operatively. The operating surgeon will not be blinded to the operative procedure. All operations will be performed by a single soft tissue knee specialist surgeon, using their default surgical technique. The only difference in surgical procedure will be the choice of graft used

Pre-operative clinical assessment will be performed by a registrar or consultant. Pre-operative AP, lateral and skyline view radiographs will be reviewed. Measurements for patella alta will be performed (Insall-Salvati and Gaton-Deschamp indices) and congruence angles will be measured. Pre-operative MRI scans will also be reviewed and Tibial Tuberosity to Trochlear groove (TTTG) distances measured.

Patients will be excluded if they require tibial tuberosity transfer or other osseous reconstructive procedures, or a lateral Z-lengthening procedure, as will the requirement for a multi-ligamentous knee reconstruction. This aims to keep the study population homogenous for the purposes of direct group comparison.

Initial outpatient follow up will be performed at 6 weeks post-operatively as per routine follow up. Post-operative rehabilitation will be in accordance with departmentally agreed post-operative physiotherapy rehabilitation protocol. Functional score assessment will be performed at 3/6/12/18/24 months post-operatively by an outpatient physiotherapist associated with the

study. Validated scoring questionnaires will be performed at each visit, to include Tegner-Lysholm activity scores, Kujala stability scores, and the Banff Patellar Instability Instrument (BPPII).

Primary end point will be two years, or MRI-confirmed re-rupture requiring revision surgery.

Patients may opt out upon recruitment, or at any time throughout the study. Data collected at clinical review will be included up to the point of opt out for statistical analysis.

Intervention Type

Procedure/Surgery

Primary outcome(s)

Functional outcomes, measured using the Tegner activity scale, Lysholm Score and the Banff Patellar Instability Index Score, pre-operatively followed by post-operative checks at 6 weeks, and then again at 3/6/12/24 months post-operatively

Key secondary outcome(s)

1. Revision surgery
2. Recurrent dislocation

The timepoints for revision and recurrent dislocation are not specified, as these will be defined by these events occurring in the patient after surgery.

Completion date

31/01/2022

Eligibility

Key inclusion criteria

Current inclusion criteria as of 29/01/2018:

1. Aged 14-25 years
2. Confirmed medial patellofemoral ligament rupture on MRI
3. Pre-operative AP, lateral and skyline radiographs
4. Unilateral or bilateral symptoms
5. Clinical signs of patellar instability
6. Closed proximal tibial physes
7. Only patients with 2 or more dislocations will be considered for inclusion

Previous inclusion criteria from 08/09/2017 to 29/01/2018:

1. Aged 14-25 years
2. Confirmed medial patellofemoral ligament rupture on MRI
3. Pre-operative AP, lateral and skyline radiographs
4. Unilateral or bilateral symptoms
5. Clinical signs of patellar instability
6. Closed proximal tibial physes

Original inclusion criteria:

1. Aged 14-25 years
2. Confirmed medial patellofemoral ligament rupture on MRI
3. Pre-operative AP, lateral and skyline radiographs
4. Unilateral or bilateral symptoms

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Mixed

Sex

All

Total final enrolment

20

Key exclusion criteria

Current exclusion criteria as of 14/02/2018:

1. Patients requiring bony transfer procedures (e.g., tibial tuberosity transfer)
2. Patients requiring a lateral release or Z-lengthening of the lateral structures
3. Patients with lateral compression signs on the skyline radiographs
4. Patients with confirmed multi-ligamentous knee injury
5. Patients with confirmed osteochondral lesions on MRI or plain radiograph pre-operatively
6. Patients who have previously underwent surgery
7. Abnormal hip or hindfoot pathology
8. Excessive femoral anteversion (>30 degrees) and/or excessive external tibial torsion (>40 degrees)
9. Excessive coronal plane deformity
10. Dejour grade C/D trochlear dysplasia
11. Excessive patellar tilt as measured by the Angle or Laurin and Fulkerson Angle methods.
12. Skeletally immature patients
13. Establish radiological changes of arthritis
14. Patients with confirmed hypersensitivity to synthetic graft material
15. A TT-TG distance >20mm on MRI/CT

Previous exclusion criteria from 08/09/2017 to 14/02/2018:

1. Patients requiring bony transfer procedures (e.g., tibial tuberosity transfer)
2. Patients requiring a lateral release or Z-lengthening of the lateral structures
3. Patients with lateral compression signs on the skyline radiographs
4. Patients with confirmed multi-ligamentous knee injury
5. Patients with confirmed osteochondral lesions on MRI or plain radiograph pre-operatively
6. Patients who have previously underwent surgery
7. Abnormal hip or hindfoot pathology
8. Excessive femoral anteversion (>30 degrees) and/or excessive external tibial torsion (>40 degrees)
9. Excessive coronal plane deformity
10. Dejour grade D trochlear dysplasia
11. Excessive patellar tilt as measured by the Angle or Laurin and Fulkerson Angle methods.
12. Skeletally immature patients

13. Establish radiological changes of arthritis
14. Patients with confirmed hypersensitivity to synthetic graft material
15. Insall-Salvati >1.2, Gaton Dechamp <1.3 or TT-TG >20mm

Original exclusion criteria:

1. Patients requiring bony transfer procedures (e.g., tibial tuberosity transfer)
2. Patients requiring a lateral release or Z-lengthening of the lateral structures
3. Patients with lateral compression signs on the skyline radiographs
4. Patients with confirmed multi-ligamentous knee injury
5. Patients with confirmed osteochondral lesions on MRI or plain radiograph pre-operatively
6. Patients who have previously undergone surgery

Date of first enrolment

01/10/2017

Date of final enrolment

31/07/2020

Locations

Countries of recruitment

United Kingdom

Northern Ireland

Study participating centre

Altnagelvin Area Hospital

Glenshane Road

Londonderry

United Kingdom

BT47 6SB

Sponsor information

Organisation

Western Health and Social Care Trust

ROR

<https://ror.org/00sb42p15>

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Western Health and Social Care Trust

Results and Publications

Individual participant data (IPD) sharing plan

Data will be kept on a secure computer account on the trust server, with password access only for the study investigators. This will be kept before being electronically destroyed after completion of the study, in line with local data protection policies. There are no plans to make this freely available in the interests of patient anonymity and confidentiality.

IPD sharing plan summary

Not expected to be made available

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Protocol article	protocol	03/05/2018		Yes	No
HRA research summary			26/07/2023	No	No
Participant information sheet		14/11/2016	03/03/2017	No	Yes
Poster results			14/03/2023	No	No