

Impact of brewer's yeast (beta-glucan supplementation) on body status and muscle damage after exercise stress

| | | |
|--|---|---|
| Submission date 01/04/2020 | Recruitment status No longer recruiting | <input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol |
| Registration date 02/04/2020 | Overall study status Completed | <input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results |
| Last Edited 15/10/2020 | Condition category Other | <input type="checkbox"/> Individual participant data |

Plain English summary of protocol

Background and study aims

Strenuous exercise has been shown to result in a temporary suppression of immune system activity. If this exercise-induced immunosuppression is continued at a high frequency, those engaging in regular strenuous exercise are at a higher risk for upper respiratory infections and other infections that may require a cessation of exercise or lead to reductions in overall performance. Exercising in the heat appears to exacerbate these physiological responses, in particular gastrointestinal distress, which encompasses a significant majority of immune cell activity.

Because of the apparent detrimental effects on immune system functioning following strenuous exercise, different strategies have been developed to attenuate these decrements in immune function. Specifically, different nutritional interventions and dietary supplement strategies have been utilized to serve as countermeasures. Beta-glucans may play a vital role in supporting innate immune system activity following strenuous exercise. Beta-glucans are a polysaccharide found within the cellular membranes of different plants, such as yeast, fungi and oats. Beta-glucan ingestion has been shown to enhance immune system activity following exercise; however, mixed results exist within the literature regarding the effectiveness of different types of beta-glucans and their ability to support immune function. Supplementation with beta-glucans derived from yeast may afford an improved level of protection from infection during periods of rigorous training. Therefore, the purpose of this study is to examine the efficacy of yeast-derived beta-glucan supplementation on reducing exercise-induced immunosuppression.

Who can participate?

Adults over 18 years, performing some form of aerobic exercise at least twice per week for the last 12 months and have a VO₂Peak between the 50th and 80th percentile for their age and gender.

What does the study involve?

Participants will be randomly allocated to receive supplement of either 250 mg/day of yeast beta-glucan or a maltodextrin placebo for 13 days. Participants will arrive fasted and complete a

bout of treadmill exercise at 55% VO₂Max in a hot and humid environment. Prior to and 0, 2, and 72 hours after completing exercise changes in white blood cell counts, pro- and anti-inflammatory cytokines, markers of muscle damage, markers of muscle function, soreness, and profile of mood states will be assessed.

What are the possible benefits and risks of participating?

The results of this study can be utilized to develop effective immune-boosting supplements to reduce exercise-induced damage and improve recovery from exercise and sport participation.

Benefits can be translated to the general population.

Risks of the blood sampling procedure include bruising, hematoma, dizziness, fainting, pain upon needle stick, and the remote risk of infection. These risks will be minimized by having trained personnel obtain blood samples using standard procedures and, sterile, single-use phlebotomy supplies. Only personnel trained and cleared to perform blood sampling in research populations will perform the blood draws.

Yeast-derived beta-glucan utilized in this study is considered to be safe in long-term studies in a healthy human population.

Each exercise bout will consist of up to 60 minutes of treadmill running in a hot (35 – 40°C), humid (40 – 45%) environment. Throughout the exercise trial and to control for clinical safety while also standardizing physiological load, continuous core body temperature measurements will be made using orally ingested temperature sensors that communicate via a radio frequency to sensors integrated with software to measure core body temperature. Throughout each exercise bout, ad libitum water intake will be permitted.

Where is the study run from?

Lindenwood University (USA)

When is the study starting and how long is it expected to run for?

November 2017 to August 2019

Who is funding the study?

Leiber GmbH (Germany)

Who is the main contact?

Prof Chad Kerksick, ckerksick@lindenwood.edu

Contact information

Type(s)

Scientific

Contact name

Prof Chad Kerksick

ORCID ID

<https://orcid.org/0000-0003-0458-7294>

Contact details

Lindenwood University
209 S Kingshighway St
Saint Charles
United States of America

63301
+1 636.627.4629
ckersick@lindenwood.edu

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

LIN-INC-2018-0014

Study information

Scientific Title

The effect of yeast beta-glucan (*Saccharomyces cerevisiae*) supplementation on muscle damage, inflammatory markers, changes in mood and muscle function following a bout of treadmill exercise in a hot and humid environment

Study objectives

The purpose of this study was to examine the efficacy of yeast beta-glucan supplementation on reducing exercise-induced immunosuppression after an extended bout of treadmill exercise in a hot and humid environment for its ability to impact inflammation, muscle damage, muscle function, and mood state.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 18/06/2018, Lindenwood University IRB (Lindenwood University, 209 S Kingshighway St, St Charles, MO 63301, USA; +1 636-949-4730; IRB@lindenwood.edu), ref: 1242265-3

Study design

Interventional randomized double-blind crossover

Primary study design

Interventional

Study type(s)

Other

Health condition(s) or problem(s) studied

Recovery from stressful exercise

Interventions

Subjects will supplement with either 250 mg/day of yeast beta-glucan or a maltodextrin placebo for 13 days. Participants will arrive fasted and complete a bout of treadmill exercise at 55%

VO2Max in a hot and humid environment. Prior to and 0, 2, and 72 hours after completing exercise changes in white blood cell counts, pro- and anti-inflammatory cytokines, markers of muscle damage, markers of muscle function, soreness, and profile of mood states will be assessed.

The order of supplementation (beta-glucan or placebo) was determined using <http://www.randomizer.org/>

Intervention Type

Supplement

Primary outcome(s)

1. Indicators of muscle damage pre, immediately after, 2h and 72 hours post-exercise:
 - 1.1. Perceived soreness measured using a 100-mm visual analog scale anchored with 0 – ‘No Soreness At All’ and 100-‘Extreme, Debilitating Soreness’
 - 1.2. Plasma creatine kinase measured using blood sample
 - 1.3. Plasma myoglobin measured using blood sample
2. Cytokines (IL-1 β , IL-6, IL-12, GM-CSF, TNF- α , IL-2, IL-7, IL-13, IFN-, IL-4, IL-8, IL-17, MCP-1, IL-6, IL-10, G-CST, and MIP-1 β) measured using blood sample assay pre, immediately after, 2h and 72 hours post-exercise
3. Muscle function (maximal voluntary isometric contraction tests and a 50-repetition isokinetic muscle fatigue test): pre, immediately after, 2h and 72 hours post-exercise
4. Mood state measured using profile of mood states (POMS): pre, immediately after, 2h and 72 hours post-exercise
5. Indicators of physiological stress (heart rate, RPE, VO₂, and core temperature values): up to 60 minutes or until reaching a core temperature of 39.2°C

Key secondary outcome(s)

Safety (Complete Blood Count and Comprehensive Metabolic Panel): pre, immediately after, 2h and 72 hours post-exercise

Completion date

03/08/2019

Eligibility

Key inclusion criteria

1. Performing some form of aerobic exercise at least twice per week for the last 12 months
2. Have a VO₂Peak between the 50th and 80th percentile for their age and gender

Participant type(s)

Healthy volunteer

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Total final enrolment

31

Key exclusion criteria

1. Any individual who is currently being treated for or diagnosed with a cardiac, respiratory, circulatory, musculoskeletal, metabolic, immune, autoimmune, psychiatric, hematological, neurological or endocrinological disorder or disease.
2. Any individual being obese (defined as body mass index > 30 kg/m² and body fat greater than 30%)
3. Any female who is pregnant or breast-feeding
4. Any individual less than 18 and greater than 50 years of age

Date of first enrolment

21/06/2018

Date of final enrolment

01/08/2019

Locations

Countries of recruitment

United States of America

Study participating centre

Lindenwood University

209 S. Kingshighway

St. Charles

United States of America

63301

Sponsor information

Organisation

Increnovo LLC

Funder(s)

Funder type

Industry

Funder Name

Leiber GmbH

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

IPD sharing plan summary

Available on request

Study outputs

| Output type | Details | Date created | Date added | Peer reviewed? | Patient-facing? |
|---------------------------------|---------|--------------|------------|----------------|-----------------|
| Results article | results | 19/04/2020 | 15/10/2020 | Yes | No |