# Effect of first trimester crown rump length measurement on rates of induction of labour for postdates.

Submission date	Recruitment status	<ul><li>Prospectively registered</li></ul>		
23/01/2004	No longer recruiting	☐ Protocol		
Registration date	Overall study status	Statistical analysis plan		
23/01/2004	Completed	[X] Results		
Last Edited	Condition category	[] Individual participant data		
03/12/2008	Pregnancy and Childbirth			

## Plain English summary of protocol

Not provided at time of registration

# Contact information

### Type(s)

Scientific

#### Contact name

Mr I Mackenzie

#### Contact details

Nuffield Department of Obstetrics and Gynaecology John Radcliffe Hospital Oxford United Kingdom OX3 9DU

# Additional identifiers

**Protocol serial number** SPGS 757

# Study information

Scientific Title

Study objectives

The project is an RCT of the effect of first trimester ultrasound on rates of induction of labour for postdates. Currently it is widespread practice in most obstetric units in the United Kingdom to induce labour 14 days after the estimated date of delivery (EDD). While this has been shown to have a beneficial effect on perinatal outcome, it may cause consumer dissatisfaction as it is perceived to be a painful intervention in an otherwise normal pregnancy. Measurement of the biparietal diameter at 20 weeks gestation has been shown to improve the accuracy of gestational age assessment, and this should lead to a reduction in rates of induction of labour for postdates. Gestational age assessment using CRL in the 1st trimester is more accurate than the biparietal diameter and may further reduce the numbers requiring induction for postdates. There is an increasing shift to perform prenatal diagnosis in the first trimester. Those providing antenatal care may have to choose the best time at which a routine scan in pregnancy can be offered. If a first trimester ultrasound scan is shown to improve the customisation and therefore the appropriateness of obstetric care this would provide further evidence to justify its introduction as well as its use in screening for Down's syndrome, either with nuchal fold or as an adjunct to serum screening.

The objective is to determine whether an ultrasound scan in the first trimester of pregnancy will lead to more accurate assessment of gestational age and consequently lead to a reduction in the number of pregnancies induced at term and fourteen days. If this is the case then there will be cost savings for the NHS, as well as the reduction in unnecessary and potentially painful interventions.

#### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

Not provided at time of registration

#### Study design

Randomised controlled trial

#### Primary study design

Interventional

#### Study type(s)

**Not Specified** 

#### Health condition(s) or problem(s) studied

Pregnancy and childbirth: Childbirth

#### **Interventions**

The control group will continue with routine antenatal care with ultrasound performed for the usual indications.

The intervention group will have an ultrasound scan between 8 and 12 weeks gestation to measure the crown-rump length (CRL) and if there is a discrepancy of greater than 5 days between the estimated date of delivery (EDD) calculated form the last menstrual period and that by CRL the EDD will be revised to that derived by scan. The EDD will be entered in the patient's obstetric notes and all subsequent management decisions will be based on this assessment of gestational age.

#### Intervention Type

Other

#### Phase

**Not Specified** 

#### Primary outcome(s)

Rates of induction of labour for postdates

#### Key secondary outcome(s))

- 1. The characteristics of labour with respect to the length of labour, the mode of delivery, the immediate perinatal outcome, analgesic requirements for labour and the overall client satisfaction with labour
- 2. Rates of gestation dependent events in pregnancy false positive and negative outcomes from maternal serum screening
- 3. The management of early pregnancy complications

#### Completion date

30/09/2002

# Eligibility

#### Key inclusion criteria

Women in the first trimester of pregnancy who do not require a first trimester ultrasound for any other clinical indication.

#### Participant type(s)

Patient

#### Healthy volunteers allowed

No

#### Age group

Adult

#### Sex

Female

#### Key exclusion criteria

Women will be excluded if there is a clinical indication for a first trimester ultrasound scan or if they refuse consent. Examples of the former would be if there was vaginal bleeding, uncertainty about viability of the pregnancy or the possibility of an ectopic pregnancy.

#### Date of first enrolment

01/02/1999

#### Date of final enrolment

30/09/2002

## Locations

#### Countries of recruitment

**United Kingdom** 

England

Study participating centre
Nuffield Department of Obstetrics and Gynaecology
Oxford
United Kingdom
OX3 9DU

# Sponsor information

## Organisation

NHS R&D Regional Programme Register - Department of Health (UK)

# Funder(s)

# Funder type

Government

#### **Funder Name**

NHS Executive South East (UK)

# **Results and Publications**

Individual participant data (IPD) sharing plan

# IPD sharing plan summary

Not provided at time of registration

# **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results of uncompleted trial	01/02/2006		Yes	No