

Investigating pelvic floor training as a treatment for severe bloating in gastrointestinal disorders

Submission date 30/12/2020	Recruitment status No longer recruiting	☐ Prospectively registered ☐ Protocol
Registration date 08/01/2021	Overall study status Completed	☐ Statistical analysis plan ☒ Results
Last Edited 21/09/2021	Condition category Digestive System	☐ Individual participant data

Plain English summary of protocol

Background and study aims

The cause of bloating, one of the most common and bothersome complaints in patients with functional gastrointestinal disorders (FGIDs), is complicated and partially understood.

The aim of this study is to investigate, in patients with severe bloating, the relationship between the defecation pattern, the severity of bloating, and the abdominal size changes, and the effectiveness of biofeedback treatment on bloating. Biofeedback training involves using electrical sensors to train

Who can participate?

Patients with FGIDs reporting bloating as their main complaint.

What does the study involve?

Participants will undergo a measurement of abdominal size (girth) using a belt around the abdomen. Participants will also be provided with dietary management with NICE advice with the addition of a lactose-free diet for two weeks. Over the two week period, participants will be asked complete a daily diary to assess their bowel function and to record girth at fasting (before eating) and two hours after lunch.

After two weeks of following diet changes, participants will be asked to fill in a questionnaire on the improvement of their bloating and abdominal girth measurements two hours after a meal.

Patients who do not report fair or major improvement at two weeks will undergo a standardized test. This will be the balloon expulsion test (BET) which uses a balloon to simulate a bowel movement and is scored as successful if the balloon is evacuated within two minutes.

Furthermore, participants will be invited to take part in biofeedback training provided at single center located in Verona, Northern Italy. This training has been previously validated for constipation due to dyssynergic defecation (DD) and will be provided by a registered nurse. The aim of this training is to improve the effort used during defecation with the purpose of relieving

bloating. To confirm the diagnosis of outlet dysfunction, all the subjects allocated to biofeedback training will undergo an evaluation of pelvic floor muscle function on straining using electromyography (EMG) testing. EMG testing uses electrodes to stimulate the pelvic muscle to measure electrical activity.

Clinical visits will occur 1, 3, and 6 months after treatment. During these visits, patients will be required to answer the questionnaires on their bloating that were completed after two weeks, and undergo physiological testing (BET and EMG evaluation).

What are the possible benefits and risks of participating?

There is a possible beneficial effect of biofeedback training on bloating. There are no anticipated risks for taking part in this study.

Where is the study run from?

University of Salerno (Italy)

When is the study starting and how long is it expected to run for?

From December 2015 to June 2019

Who is funding the study?

Investigator initiated and funded

Who is the main contact?

Prof Paola Iovino

piovino@unisa.it

Contact information

Type(s)

Scientific

Contact name

Prof Paola Iovino

ORCID ID

<https://orcid.org/0000-0002-9568-0680>

Contact details

Gastrointestinal Unit

Department of Medicine, Surgery and Dentistry

Scuola Medica Salernitana

University of Salerno

S. Allende

Baronissi

University Hospital "San Giovanni di Dio e Ruggi D'Aragona"

Piazzale Ippocrate 1

Salerno

Italy

84100

+39 335 7822672

piovino@unisa.it

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

Studio FgiD

Study information

Scientific Title

Investigating pelvic floor biofeedback as a treatment for severe bloating in functional gastrointestinal disorders with outlet dysfunction

Study objectives

1. Bloating and abdominal distension could be secondary to impaired gas or non-gas emptying possibly due to disordered defecation
2. Pelvic floor biofeedback is an effective treatment for severe bloating in functional gastrointestinal disorders with outlet dysfunction

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 06/05/2013, revision approved 13/06/2019, Ethical Committee-Azienda Sanitaria Locale Napoli 3 Sud (Via Marconi 66, Torre del Greco, Napoli, 80049, Italy), ref: n°3/2013, revised approval ref: 0089264

Study design

Multicenter observational cohort study

Primary study design

Observational

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Bloating treatment in functional gastrointestinal disorders

Interventions

All patients are screened for the severity of bloating that is their main complaint and diagnosed according to Rome III criteria for FGIDs. If the bloating is > 24 mm on 0-100 mm-VAS scale they undergo a girth measurement abdominal girth measured by a belt around the abdomen at

standardized sites and start 2 weeks of dietary management with NICE advice with the addition of lactose-free diet and complete a daily diary to assess bowel function as well as record girth measurements at fasting and 2 hours after lunch.

After two weeks at the first visit, participants will be asked to fill in a questionnaire on the subjective improvement of bloating on a 5-point Likert scale (worse to major improvement), a further assessment of abdominal bloating (using a 0-100 visual analogue scale), and abdominal girth measurements 2 h after a meal. Patients who did not report fair or major improvement /cure will undergo a standardized balloon expulsion test (BET) which will be scored as either successful or failed if the balloon could not be evacuated within 2 min. Furthermore, they will be invited to take part in a biofeedback protocol to improve the defecation effort with the purpose of ameliorating bloating that was provided at single center located in Verona, Northern Italy. To confirm the diagnosis of outlet dysfunction, all the subjects allocated to biofeedback will undergo a basal evaluation of pelvic floor muscle function on straining by pelvic floor electromyography (EMG) testing according to a previously described protocol. A registered nurse provided the BT previously validated for constipation due to dyssynergic defecation (DD). Clinical visits were scheduled at 1, 3, and 6 months post-treatment. During these visits patients will be required to answer a question on subjective perception of symptom improvement on a 5-point Likert scale (worse, no improvement, mild, fair, and major improvement/cure), will be assessed for bloating on the (0-100) VAS, and undergo physiological testing (BET and EMG evaluation).

Intervention Type

Behavioural

Primary outcome(s)

1. Response to diet intervention or pelvic floor biofeedback treatment (BT) measured using patient response to the question: "Compared to the interval preceding the treatment (diet intervention or biofeedback) how would you score your bloating improvement: worse (0), no improvement (1), mild (2), fair (3), or major improvement/cure (4)?" , where all patients reporting fair or major improvement/cure will be considered as responding, measured at 2 weeks , and 1, 3, and 6 months after BT

Key secondary outcome(s)

1. Effect of diet intervention on bloating severity measured using a 0-100 visual analogue scale (VAS), and abdominal girth (using a belt around the abdomen at standardized sites) at baseline and 2 weeks
2. Effect of diet intervention and pelvic floor biofeedback treatment (BT) on quality of life changes measured using the Short Form 36 (SF36) at 2 weeks and after BT
3. Effect of pelvic floor biofeedback treatment (BT) on bloating severity measured using a 0-100 visual analogue scale (VAS), balloon expulsion test (BET), and electromyography (EMG) testing on straining at 2 weeks (before BT), and then 1, 3, and 6 months after BT

Completion date

30/06/2019

Eligibility

Key inclusion criteria

1. Aged between 18 and 75 years
2. Able to understand and the willingness to comply with the study procedures

3. Average daily abdominal bloating score ≥ 24 on a 100-mm VAS with/without visible abdominal distension

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

180

Key exclusion criteria

1. Pregnancy
2. Serious, or unstable medical condition
3. Insulin-dependent diabetes mellitus
4. Major psychiatric diagnosis
5. Endocrine diseases
6. History of eating disorders
7. History of drug or alcohol abuse
8. Previous abdominal surgery, except appendectomy or cholecystectomy

Date of first enrolment

01/03/2017

Date of final enrolment

30/06/2018

Locations

Countries of recruitment

Italy

Study participating centre

Gastrointestinal Unit, University Hospital "San Giovanni di Dio e Ruggi D'Aragona"

Piazzale Ippocrate 1

Salerno

Italy

84100

Study participating centre
Pio Albergo Trivulzio Hospital
Via Trivulzio, 15
Milan
Italy
20146

Study participating centre
S. Giovanni Addolorata Hospital
Via dell'Amba Aradam 9
Rome
Italy
00184

Study participating centre
Division of Gastroenterology B, Azienda Ospedaliera Universitaria Integrata Verona
Piazzale Aristide Stefani, 1
Verona
Italy
37126

Sponsor information

Organisation
University of Salerno

ROR
<https://ror.org/0192m2k53>

Funder(s)

Funder type
Other

Funder Name
Investigator initiated and funded

Results and Publications

Individual participant data (IPD) sharing plan

The data sharing plans for the current study are unknown and will be made available at a later date.

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		16/09/2021	21/09/2021	Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes