

# The humeral shaft fracture trial: surgical versus non-surgical interventions for humeral shaft fractures in patients aged 18 years or older

<b>Submission date</b> 03/03/2020	<b>Recruitment status</b> No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered
		<input checked="" type="checkbox"/> Protocol
<b>Registration date</b> 13/05/2020	<b>Overall study status</b> Completed	<input checked="" type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
<b>Last Edited</b> 16/05/2025	<b>Condition category</b> Injury, Occupational Diseases, Poisoning	<input type="checkbox"/> Individual participant data
		<input checked="" type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

A humeral shaft fracture is a break in the long bone of the upper arm. It occurs mainly in two groups of individuals; young men and older women, as their bones are more fragile. Currently, the most common treatment for these fractures is non-operative. About 70% of cases are treated using a cast for 2 weeks and then a brace until the bone begins to heal properly – although there is a large variation in treatments between and in hospitals. The risk of complications is low and the cost is also relatively low at £1,100. The disadvantages are that the patient is immobilised for a prolonged period and the cumbersome cast can lead to significant pain and discomfort in some patients. There is also a 20% chance that the break will not heal. This then requires surgery and involves additional costs of approximately £15,500. There appears to be a worldwide trend towards treating these fractures with surgery (rather than a cast and a brace), however, there is no high-quality evidence that this is indeed a better option. Various reviews of the current evidence have recognised the need for further trials. Surgery is the more expensive route, and has a higher risk of complication e.g. infection and nerve damage. However, there is a better chance of the bone healing successfully and the patient is likely to recover more quickly allowing them to regain their independence sooner. The aim of this study is to directly compare these two methods of treating fractures of the humeral shaft. The researchers want to find out whether arm function and quality of life in patients with this fracture is better with the more conservative cast-and-brace treatment, or with surgery. They also need to compare the cost-effectiveness of both approaches. They want to produce sound evidence to establish if the drawbacks of surgery are balanced by improved results and acceptable costs.

### Who can participate?

Patients aged 18 over who have a broken upper arm

### What does the study involve?

Participants will be randomly allocated to receive either surgical or non-surgical (a brace) treatment for their broken upper arm. The technique of surgery used for those patients allocated to the surgery group will be chosen by the surgeon. Surgery will typically be followed

by 2 weeks in a sling. Patients treated non-surgically will have a cast applied in the Emergency Department which they will use for 2 weeks. They will then change to a brace which is usually worn for a further 8-10 weeks. Both groups will be given a structured rehabilitation programme. The researchers will collect data with regards to pain, time off work and driving, and functionality for 12 months after treatment.

What are the possible benefits and risks of participating?

Both treatments are used across the NHS currently for this type of fracture and are not new or experimental. There is a small risk of complications for participants who have the operation, such as infection or prominent metalwork as there would be with any surgery, both of which might require further treatment. The main potential risk of the brace treatment is that in about 1 of every 5 patients the bones do not heal properly. If this happens your fracture may require further treatment, which could be an operation. Both treatments are used across the NHS so there is no specific advantage for taking part in the study.

Where is the study run from?

University of Oxford (UK)

When is the study starting and how long is it expected to run for?

January 2020 to February 2025

Who is funding the study?

National Institute for Health Research Health Technology Assessment Programme (NIHR HTA) (UK)

Who is the main contact?

Miss Hannah Crook

[hush@ndorms.ox.ac.uk](mailto:hush@ndorms.ox.ac.uk)

**Study website**

<http://www.hushstudy.org>

## Contact information

**Type(s)**

Scientific

**Contact name**

Miss Hannah Crook

**Contact details**

Oxford Trauma Kadoorie Centre

Level 3 John Radcliffe Hospital

Headley Way

Headington

Oxford

United Kingdom

OX3 9DU

+44 (0)1865 227912

[hush@ndorms.ox.ac.uk](mailto:hush@ndorms.ox.ac.uk)

# Additional identifiers

## EudraCT/CTIS number

Nil known

## IRAS number

277059

## ClinicalTrials.gov number

Nil known

## Secondary identifying numbers

HTA - NIHR127817, IRAS 277059

# Study information

## Scientific Title

The HUmeral SHAft fracture trial (HUSH): a multi-centre prospective randomised superiority trial of surgical versus non-surgical interventions for humeral shaft fractures in patients aged 18 years or older

## Acronym

HUSH

## Study objectives

The aim of this pragmatic randomised controlled trial is to evaluate the clinical and cost-effectiveness of functional bracing, compared to surgical fixation for the treatment of humeral shaft fractures in patients over the age of 18.

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

Approved 02/06/2020, East of England - Cambridge Central Research Ethics Committee (Royal Standard Place Nottingham NG1 6FS, UK; +44 (0)207 104 8388; cambridgecentral.rec@hra.nhs.uk), ref: 20/EE/0127

## Study design

Pragmatic randomised controlled trial

## Primary study design

Interventional

## Secondary study design

Randomised controlled trial

## Study setting(s)

Hospital

## Study type(s)

## Treatment

### Participant information sheet

Not available in web format, please use contact details to request a participant information sheet

### Health condition(s) or problem(s) studied

Humeral shaft fracture

### Interventions

Participants will be randomised (online using RRAMP) in a 1:1 ratio stratified by centre; whether they are aged under 50 or 50 and over and whether they had a nerve injury at presentation or not, to either surgery or a brace. They will be randomised to be treated surgically or non-surgically.

The technique of surgery used for those patients allocated to the surgery group will be chosen by the surgeon. Surgery will typically be followed by 2 weeks in a sling.

Patients treated non-surgically will have a cast applied in the Emergency Department which they will use for 2 weeks. They will then change to a brace which is usually worn for a further 8-10 weeks.

Both groups will be given a structured rehabilitation programme. The trial will last for 12 months. Patients will be followed up at 6 months and 12 months after their injury. They will be asked about their quality of life, daily activities, pain, physiotherapy treatment and any complications. The researchers will also look at the resources and services they have used to determine the costs involved in both treatments.

### Intervention Type

Procedure/Surgery

### Primary outcome measure

Function measured using the Disabilities of Arms Shoulders and Hand (DASH) patient-reported outcome questionnaire at 12 months

### Secondary outcome measures

1. Function assessed using the Disabilities of Arms Shoulders and Hand (DASH) at baseline, 8 weeks, 3 and 6 months
2. Early pain recovery assessed using pain visual analogue scale (VAS) weekly in the first 8 weeks
3. Sport and performing arts functioning assessed using the DASH sports/performing arts at baseline, 8 weeks, 3, 6 and 12 months
4. Function assessed using the patient-reported outcome measurement information system (PROMIS) upper extremity at baseline, 4 weeks, 8 weeks, 3, 6 and 12 months
5. Function assessed using the PROMIS pain interference at baseline, 4 weeks, 8 weeks, 3, 6 and 12 months
6. Quality of life measured using the EQ-5D-5L at baseline, 8 weeks, 3, 6 and 12 months
7. Complications reported at 8 weeks, 3, 6 and 12 months
8. Cost-effectiveness measured using the Work Productivity and Activity Impairment Questionnaire at 3, 6 and 12 months
9. Time off work and driving recorded in the first 8 weeks

**Overall study start date**

01/01/2020

**Completion date**

28/02/2025

## Eligibility

**Key inclusion criteria**

1. Adult patients aged 18 years and older with a fracture of the humeral shaft (diaphysis)
2. Fracture of the humeral diaphysis which the surgeon believes may benefit from surgical fixation. 'Diaphysis' defined as the section of bone outside 1 Muller-square of the proximal and distal ends of the humerus (Müller 1990)
3. Participant is willing and able to give informed consent for participation in the study
4. Male or female, aged 18 years or above

**Participant type(s)**

Patient

**Age group**

Adult

**Lower age limit**

18 Years

**Sex**

Both

**Target number of participants**

334

**Total final enrolment**

334

**Key exclusion criteria**

1. The fracture is open
2. The fracture is complicated by local tumour deposits
3. The index injury occurred more than 16 days prior to recruitment
4. The patient is unable to adhere to trial procedures
5. Other upper limb injuries which may reasonably be expected to affect responses to outcome PROMs

**Date of first enrolment**

01/07/2020

**Date of final enrolment**

17/11/2024

## Locations

**Countries of recruitment**

England

United Kingdom

**Study participating centre**

**Oxford University Hospitals NHS Foundation Trust**

John Radcliffe Hospital

Headley Way

Headington

Oxford

United Kingdom

OX3 9DU

## **Sponsor information**

**Organisation**

University of Oxford

**Sponsor details**

Joint Research Office

1st Floor

Boundary Brook House

Churchill Drive

Oxford

England

United Kingdom

OX3 7GB

+44 (0)1865 289886

ctrng@admin.ox.ac.uk

**Sponsor type**

University/education

**Website**

<https://researchsupport.admin.ox.ac.uk/ctrng>

**ROR**

<https://ror.org/03sbpja79>

## **Funder(s)**

**Funder type**

Government

**Funder Name**

Health Technology Assessment Programme

**Alternative Name(s)**

NIHR Health Technology Assessment Programme, HTA

**Funding Body Type**

Government organisation

**Funding Body Subtype**

National government

**Location**

United Kingdom

## Results and Publications

**Publication and dissemination plan**

The protocol will be available prior to the completion of recruitment. The Statistical Analysis Plan and Health Economics Analysis Plan will be prepared before the final data has been collected. It is planned that each of these will be published in open-access journals.

The researchers will aim to publish the study findings as widely, speedily and efficiently as possible to allow their introduction to clinical practice and for evidence-based medicine to be used in the treatment of these patients. The research team has experience in translating research findings into clinical practice through the development of clear, evidence-based care pathways to improve patient care. A collaborative effort between health care practitioners, researchers and patients will help to make this possible. The involvement of service users (i.e. Patient Advisory Group) will ensure that project outputs are patient-orientated and relevant to the end-users.

Through the planned outputs, the study is expected to play a key role in enhancing the evidence base on efficacy, safety and cost-effectiveness of the treatment of humeral shaft fractures. Importantly, the executive summary and copy of the trial report will be sent to the National Institute of Health and Clinical Excellence (NICE) and other relevant bodies, including Clinical Commissioning Groups, so that the study findings can inform their deliberations and be translated into clinical practice nationally via the NICE non-complex fracture guidelines (<https://www.nice.org.uk/guidance/ng38>). The researchers will also work with the relevant National Clinical Director in the Department of Health to help ensure the findings of the trial are considered when implementing policy.

Finally, the researchers will work with the relevant Specialty Advisory Committees (SAC) to incorporate the findings into the training curriculum for clinicians who will undertake treatment humeral shaft fractures.

**Intention to publish date**

28/02/2026

**Individual participant data (IPD) sharing plan**

The data-sharing plans for the current study are unknown and will be made available at a later date; please contact [hush@ndorms.ox.ac.uk](mailto:hush@ndorms.ox.ac.uk) for more information.

**IPD sharing plan summary**

Data sharing statement to be made available at a later date

**Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">HRA research summary</a>	version 3.0		28/06/2023	No	No
<a href="#">Protocol article</a>		22/04/2024	03/12/2024	Yes	No
<a href="#">Statistical Analysis Plan</a>		16/05/2025	16/05/2025	No	No