

Enterogel® in the treatment of Irritable Bowel Syndrome with Diarrhoea

Submission date 14/11/2017	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 15/11/2017	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 20/09/2022	Condition category Digestive System	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Irritable Bowel Syndrome (IBS) is a common condition with symptoms causing around 1 in 10 people to seek help from their GP at some point in their lives. It is a combination of abdominal discomfort or pain and altered bowel habits. Depending on the stool consistency, IBS can be classified as IBS with diarrhoea (IBS-D), IBS with constipation (IBS-C), or mixed IBS (IBS-M). IBS is often a chronic condition that can significantly impair quality of life and cause absence from work and school. The emergence of new drugs for IBS has been slow and there is a need for new treatment alternatives, including drug-free treatments, which are easy to use and can safely be used in different types of patients, including in children and pregnant women. Currently available drug-free treatments include Enterogel®, which is an intestinal adsorbent that has been approved for use in patients with IBS-D and is available over-the-counter in the UK. It is an orally consumed gel-like product classified as a medical device, which has been shown to adsorb harmful substances like bacterial toxins in the gut. There have been studies suggesting that Enterogel® and other intestinal adsorbents can improve diarrhoea, bloating and abdominal pain in patients with IBS-D. However, larger studies are needed to further investigate the potential benefits of Enterogel® in the treatment of IBS-D. This study aims to recruit 430 participants with IBS-D. The goal is to investigate whether Enterogel® can improve IBS-D symptoms, including diarrhoea and abdominal pain. In addition, the study aims to investigate the impact of this treatment on the quality of life, work productivity and activity and general health. The safety and tolerability of Enterogel® in this condition will also be evaluated.

Who can participate?

Adults aged 16-75 who have been diagnosed with IBS-D and have diarrhoea symptoms

What does the study involve?

The study involves in total four visits and one follow-up call over 26 weeks. At each visit and at given time points between the visits, the participants complete questionnaires asking about their IBS symptoms, quality of life, work productivity and activity and general health. In addition, the participants are asked to complete a study diary throughout the study to record their symptoms and treatment use. The diary is available as an electronic and paper diary. The diary should be completed daily for the first 18 weeks. At the first visit (week 0), potential participants are provided with the study diary to record their diarrhoea and abdominal symptoms for 2

weeks. Those who had sufficient symptoms over this period are eligible to take part in the study and are randomly allocated to one of two treatment groups at the second visit (week 2). The first group receive dummy treatment and the second group receive Enterosgel® for 8 weeks. Neither the participants nor the research team know which treatment the participant receives. After 8 weeks, the participants attend the third visit (week 10) and all participants are given Enterosgel® for the next 8 weeks. This time, the participants are informed about the treatment they receive. After 8 weeks of treatment with Enterosgel®, the participants attend the fourth study visit (week 18) and return to standard-of-care. A final follow-up call takes place 8 weeks later (week 26).

What are the possible benefits and risks of participating?

To allow all patients to try Enterosgel®, this study includes an 8-week period where all participants receive it. Some participants may find that Enterosgel® helps with their IBS symptoms while others might not experience any benefit. The aim of this study is to evaluate what proportion of participants can benefit from this treatment, and what the benefits are. In the future, this information can be helpful to the patients and healthcare professionals when they are considering different treatment alternatives. Enterosgel® has been available in some countries for 30 years and is commonly used as a treatment for gastrointestinal conditions, including acute diarrhoea and IBS-D. According to the manufacturer, possible known side effects are nausea and constipation. Enterosgel® is completely excreted from the gut and does not get absorbed into the blood circulation.

Where is the study run from?

This study will be conducted at about 30 medical practices, secondary care centres and private clinics in the UK and a virtual site. In addition to the participating research sites, the study is advertised through local pharmacies and through an IBS research register called ContactME-IBS (<https://www.contactme-ibs.co.uk>)

When is the study starting and how long is it expected to run for?

March 2017 to October 2021

Who is funding the study?

Bioline Products s.r.o

Who is the main contact?

Dr Carol Howell

Contact information

Type(s)

Scientific

Contact name

Dr Carol Howell

Contact details

Enteromed Ltd
85 Great Portland Street
London
United Kingdom
W1W 7LT

+44 (0)7596887046
research@enteromed.co.uk

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

ENT04UK

Study information

Scientific Title

Randomised, double-blind, placebo-controlled multi-centre study to assess the efficacy, tolerability and safety of Enterogel® in the treatment of Irritable Bowel Syndrome with Diarrhoea (IBS-D) in adults (RELIEVE IBS-D)

Acronym

RELIEVE IBS-D

Study objectives

The hypothesis is that Enterogel® is superior compared with placebo in terms of patient-reported outcomes for stool consistency and abdominal pain, in patients with IBS-D.

Ethics approval required

Old ethics approval format

Ethics approval(s)

North East - Tyne & Wear South Research Ethics Committee, 19/02/2018, REC ref: 18/NE/0023

Study design

Randomised double-blind placebo-controlled multi-centre trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Irritable Bowel Syndrome with diarrhoea (IBS-D)

Interventions

Using a centralised computer-based randomisation tool, patients will be randomised 1:1 to receive intestinal adsorbent Enterogel® or placebo. The patients will receive blinded study treatment for 8 weeks according to study-specific dosage instructions, which allow adjusting the

dosage based on IBS symptoms. The treatment is provided in 90g tubes, each containing a single treatment dose. The Enterosgel® tubes contain 22.5g Enterosgel® pre-diluted in water. The placebo tubes contain water with chalk added to mimic the appearance and consistency of Enterosgel®. Updated 17/10/2018: The control group will receive identical tubes containing placebo.

Treatments are taken orally up to 6 doses per day, depending on the symptoms.

After 8 weeks of blinded treatment, all patients will receive open-label Enterosgel® treatment for 8 weeks to take if they experience diarrhoea.

Following the open-label treatment period, all patients will return to standard-of-care and will be followed up for 8 weeks.

Intervention Type

Device

Phase

Not Applicable

Primary outcome(s)

Percentage of patients defined as responders for abdominal pain and stool consistency based on daily study diary data, during at least 4 weeks in the 8-week blinded treatment period, where:

1. An Abdominal Pain Intensity Weekly Responder is defined as a patient who experiences a decrease in the weekly average abdominal pain score of at least 30 percent compared with baseline. The weekly average abdominal pain score is derived by scoring the worst pain experienced each day and taking the average for one week

AND

2. A Stool Consistency Weekly Responder is defined as a patient who experiences a 50 percent or greater reduction in the number of days per week with at least one stool that has a consistency of Bristol Stool Form Scale (BSFS) Type 6 or 7 compared with baseline

Key secondary outcome(s)

Original secondary outcome measures:

Double-blind treatment phase and open-label treatment phase:

1. Stool frequency based on daily study diary data; mean over 8 weeks and the last 4 weeks of each treatment period
2. Stool consistency assessed as average number of days/week with Bristol Stool Scale type >5 based on daily study diary data and percentage of responders over 8 weeks and the last 4 weeks of each treatment period
3. Abdominal pain based on daily study diary data; mean on a VAS scale from 0 to 10 over 8 weeks and the last 4 weeks, percentage of responders over 8 weeks and the last 4 weeks of each treatment period
4. Bloating based on weekly study diary data; mean weekly score on a numerical scale from 0 to 6 over 8 weeks and the last 4 weeks of each treatment period
5. Urgency based on weekly study diary data; mean weekly score on a numerical scale from 0 to 6 over 8 weeks and the last 4 weeks of each treatment period
6. Percentage of patients reporting adequate relief of global IBS symptoms based on weekly study diary data weekly from baseline until week 8 of each treatment period
7. Irritable Bowel Syndrome Severity Scoring System (IBS-SSS) score weekly from baseline until

week 8 of each treatment period

8. Irritable Bowel Syndrome Quality of Life (IBS-QOL) score every 4 weeks from baseline until week 8 of each treatment period
9. Patient Health Questionnaire-12 Somatic Symptom scale (PHQ-12 SS) score to measure severity of somatic symptoms excluding gastrointestinal symptoms every 4 weeks from baseline until week 8 of each treatment period
10. IBS-related Work Productivity and Activity Impairment (WPAI:IBS) score; weekly from baseline until week 8 of each treatment period
11. Use of rescue medication, i.e. loperamide, based on weekly study diary data; number of days over each 8 weeks of treatment
12. Adverse events over each 8 weeks of treatment

Follow-up:

1. Satisfaction with bowel symptoms based on weekly study diary data; mean weekly score on a numerical scale from 0 to 6, over 8 weeks
2. Satisfaction with abdominal pain based on weekly study diary data; mean weekly score on a numerical scale from 0 to 6, over 8 weeks
3. Loperamide use based on weekly study diary data; mean number of days per week over 8 weeks
4. Irritable Bowel Syndrome Severity Scoring System (IBS-SSS) score at baseline and 8 weeks
5. Patient Health Questionnaire-12 Somatic Symptom scale (PHQ-12 SS) score to measure severity of somatic symptoms excluding gastrointestinal symptoms at baseline and 8 weeks
6. Irritable Bowel Syndrome Quality of Life (IBS-QOL) score at baseline and 8 weeks
7. IBS-related Work Productivity and Activity Impairment (WPAI:IBS) score at baseline and 8 weeks
8. Percentage of patients who maintained treatment benefit over 8 weeks after cessation of treatment based on weekly study diary data over 8 weeks

Updated 17/10/2018:

Follow-up:

In patients who reported adequate relief in the last 4 weeks of open-label period:

1. Maintenance of treatment benefit (percentage of patients who report increased or maintained treatment benefit at 8 weeks)
2. Enterosgel® use (percentage of patients who report having used Enterosgel® during the follow-up period; frequency of use in these patients)
3. Loperamide use (percentage of patients who report having used less loperamide during the follow-up period than before the trial)
4. Irritable Bowel Syndrome Severity Scoring System (IBS-SSS) score at baseline and 8 weeks
5. Patient Health Questionnaire-12 Somatic Symptom scale (PHQ-12 SS) score to measure severity of somatic symptoms excluding gastrointestinal symptoms at baseline and 8 weeks
6. Irritable Bowel Syndrome Quality of Life (IBS-QOL) score at baseline and 8 weeks
7. IBS-related Work Productivity and Activity Impairment (WPAI:IBS) score at baseline and 8 weeks
8. Percentage of patients who maintained treatment benefit over 8 weeks after cessation of treatment based on weekly study diary data over 8 weeks

Updated 07/01/2020:

Double-blind treatment phase and open-label treatment phase:

1. Stool frequency based on daily study diary data; mean over 8 weeks and the first and last 4 weeks of each treatment period
2. Stool consistency assessed as average number of days/week with Bristol Stool Scale type >5 based on daily study diary data (over 8 weeks and the first and last 4 weeks of each treatment

period) and percentage of responders

3. Abdominal pain based on daily study diary data; mean on a VAS scale from 0 to 10 over 8 weeks and the last 4 weeks, percentage of responders over 8 weeks and the last 4 weeks of each treatment period
4. Bloating based on weekly study diary data; mean weekly score on a numerical scale from 0 to 6 over 8 weeks and the last 4 weeks of each treatment period
5. Urgency based on weekly study diary data; mean weekly score on a numerical scale from 0 to 6 over 8 weeks and the last 4 weeks of each treatment period
6. Percentage of patients reporting adequate relief of global IBS symptoms based on weekly study diary data over 8 weeks and the first and last 4 weeks of each treatment period
7. Irritable Bowel Syndrome Severity Scoring System (IBS-SSS) score average over 8 weeks and the first and last 4 weeks
8. Irritable Bowel Syndrome Quality of Life (IBS-QOL) 4-weekly score average over 8 weeks and week 4 and week 8 scores
9. Patient Health Questionnaire-12 Somatic Symptom scale (PHQ-12 SS) score to measure severity of somatic symptoms excluding gastrointestinal symptoms 4-weekly score average over 8 weeks and week 4 and week 8 scores
10. IBS-related Work Productivity and Activity Impairment (WPAI:IBS) weekly score average over 8 weeks of each treatment period
11. Use of rescue medication, i.e. loperamide, based on weekly study diary data; number of days over 8 weeks and the last 4 weeks
12. Adverse events over each 8 weeks of treatment

Completion date

31/10/2021

Eligibility

Key inclusion criteria

Current inclusion criteria as of 04/04/2019:

1. Written informed consent
2. Irritable Bowel Syndrome with diarrhoea (IBS-D) according to Rome IV criteria
3. Aged 16-75
4. Considered suitable to take part in the study by the consenting doctor/nurse
5. Diary completed on at least 11 of 14 days ($\geq 75\%$) during the screening period

Previous inclusion criteria from 17/10/2018 to 04/04/2019:

1. Written informed consent
2. Irritable Bowel Syndrome with diarrhoea (IBS-D) according to Rome IV criteria
3. Aged 18-75
4. Considered suitable to take part in the study by the consenting doctor/nurse
5. Diary completed on at least 11 of 14 days ($\geq 75\%$) during the screening period

Original inclusion criteria:

1. Irritable Bowel Syndrome with diarrhoea (IBS-D) according to Rome IV criteria
2. Negative calprotectin test or adequate exclusion of Inflammatory Bowel Disease (IBD)
3. Aged 18+
4. Considered suitable to take part in the study by the consenting physician

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Mixed

Lower age limit

16 years

Upper age limit

75 years

Sex

All

Total final enrolment

439

Key exclusion criteria

Current exclusion criteria as of 07/01/2020:

1. Loose stools (BSFS 6 or 7) on less than 3 days during the 14 days after Screening Visit
2. Average abdominal pain <2.5 during the 14 days after Screening Visit (scale 0–10: 0 = no pain; 10 = worst possible pain)
3. Previously diagnosed coeliac disease
4. Previously diagnosed Inflammatory Bowel Disease
5. Previous bowel cancer or bowel resection
6. Other previously known gastrointestinal disorder contributing to the diarrhoea
7. Unexplained weight loss
8. Unexplained rectal bleeding
9. Previous use of Enterosgel®
10. Use of antidepressant agents, unless used at a stable dose for at least 6 weeks
11. Use of any probiotic supplements, other intestinal adsorbents, slow-release medications or strong opioids
12. Participation in any research where treatment is provided, or was provided in the last three months
13. Pregnancy

Previous exclusion criteria from 04/04/2019 to 07/01/2020:

1. Loose stools (BSFS 6 or 7) on less than 3 days during the 14 days after Screening Visit
2. Average abdominal pain <3 during the 14 days after Screening Visit (scale 0–10: 0 = no pain; 10 = worst possible pain)
3. Previously diagnosed coeliac disease
4. Previously diagnosed Inflammatory Bowel Disease
5. Previous bowel cancer or bowel resection
6. Other previously known gastrointestinal disorder contributing to the diarrhoea
7. Unexplained weight loss
8. Unexplained rectal bleeding
9. Previous use of Enterosgel®
10. Use of antidepressant agents, unless used at a stable dose for at least 6 weeks
11. Use of any probiotic supplements, other intestinal adsorbents, slow-release medications or

strong opioids

12. Participation in any research where treatment is provided, or was provided in the last three months

13. Pregnancy

Previous exclusion criteria from 17/10/2018 to 04/04/2019:

1. Loose stools (BSFS 6 or 7) on less than 3 days during the 14 days after Screening Visit
2. Average abdominal pain <3 during the 14 days after Screening Visit (scale 0–10: 0 = no pain; 10 = worst possible pain)
3. IBS-SSS \geq 400
4. Previously diagnosed coeliac disease
5. Previously diagnosed Inflammatory Bowel Disease
6. Previous bowel cancer or bowel resection
7. Other previously known gastrointestinal disorder contributing to the diarrhoea
8. Unexplained weight loss
9. Unexplained rectal bleeding
10. Previous use of Enterosgel®
11. Use of antidepressant agents, unless used at a stable dose for at least 6 weeks
12. Use of any probiotic supplements, other intestinal adsorbents, slow-release medications or strong opioids
13. Participation in any research where treatment is provided, or was provided in the last three months
14. Pregnancy

Original exclusion criteria:

1. Loose stools (BSFS 6 or 7) on less than 3 days during the last 14 days before Baseline
2. IBS-SSS \geq 400
3. Organic gastrointestinal disorder
4. Celiac disease (existing diagnosis)
5. Lactose malabsorption (existing diagnosis)
6. Bile acid malabsorption (existing diagnosis)
7. Bowel cancer
8. Previous use of Enterosgel®
9. Pregnancy
10. Diary completed on at least 11 of 14 days (\geq 75%) during the screening period

Date of first enrolment

15/10/2018

Date of final enrolment

21/04/2021

Locations

Countries of recruitment

United Kingdom

England

Study participating centre
University Hospital of North Durham
United Kingdom
DH1 5TW

Study participating centre
Newcastle Upon Tyne Hospitals
United Kingdom
NE7 7DN

Study participating centre
Bodey Medical Centre
United Kingdom
M14 6WP

Study participating centre
The Village Practice
United Kingdom
FY5 2TZ

Study participating centre
Blackpool Victoria Hospital
United Kingdom
FY3 8NR

Study participating centre
Vauxhall Primary Health Care
United Kingdom
L5 8XR

Study participating centre
Ashgate Medical Practice
United Kingdom
S40 4AA

Study participating centre

Newton Place Surgery

United Kingdom

ME13 8FH

Study participating centre

Pound Hill Medical Group

United Kingdom

RH10 7DX

Study participating centre

West Walk Surgery

United Kingdom

BS37 4AX

Study participating centre

Pickering Medical Centre

United Kingdom

YO18 8BL

Study participating centre

Functional Gut Clinic

United Kingdom

W1G 6NB

Study participating centre

University Hospital Coventry and Warwickshire

United Kingdom

CV2 2DX

Study participating centre

Warrington Hospital

United Kingdom

WA5 1QG

Study participating centre

Queen's Road Surgery

United Kingdom

DH8 0BW

Study participating centre

Well Close Medical Group

United Kingdom

TD15 1LL

Study participating centre

Well Close Medical Group

United Kingdom

TD15 1LL

Study participating centre

Redburn Park Medical Centre

United Kingdom

NE29 6HT

Study participating centre

Derriford Hospital

United Kingdom

PL6 8DH

Study participating centre

Royal Stoke University Hospital

United Kingdom

ST4 6QG

Study participating centre

Hadrian Primary Care Alliance

United Kingdom

NE43 7LL

Study participating centre

Castlegate and Derwent Surgery
United Kingdom
CA13 9HT

Study participating centre
Carmel Medical Practice
United Kingdom
DL3 8SQ

Study participating centre
Stepping Hill Hospital
United Kingdom
SK2 7JE

Study participating centre
Royal Bournemouth Hospital
United Kingdom
BH7 7DW

Study participating centre
St George's University Hospital
United Kingdom
SW17 0QT

Study participating centre
King's College Hospital
United Kingdom
SE5 9RS

Study participating centre
Doncaster Royal Infirmary
Armthorpe Road
Doncaster
United Kingdom
DN2 5LT

Study participating centre

Wythenshawe Hospital
Southmoor Road
Manchester
United Kingdom
M23 9LT

Sponsor information

Organisation

Enteromed Ltd

ROR

<https://ror.org/013bz8f47>

Funder(s)

Funder type

Industry

Funder Name

Bioline Products s.r.o

Results and Publications

Individual participant data (IPD) sharing plan

After publication, individual anonymised participant data that underlie the results reported in the publication will be available to researchers upon request from Dr Carol Howell at Enteromed Ltd. Participants have provided their consent for the sharing of anonymised data for research purposes.

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		27/06/2022	28/06/2022	Yes	No
Protocol article		30/01/2020	19/10/2021	Yes	No
HRA research summary			28/06/2023	No	No
Protocol file		29/06/2020	20/09/2022	No	No
Study website	Study website	11/11/2025	11/11/2025	No	Yes