# A study on complete vs. incomplete small bowel examination using single-balloon enteroscopy in obscure gastrointestinal bleeding

<b>Submission date</b> 04/04/2025	<b>Recruitment status</b> No longer recruiting	[X] Prospectively registered		
		☐ Protocol		
Registration date 10/04/2025	Overall study status Completed	Statistical analysis plan		
		Results		
Last Edited	Condition category	Individual participant data		
14/04/2025	Digestive System	[X] Record updated in last year		

#### Plain English summary of protocol

Background and study aims

Obscure gastrointestinal bleeding (OGIB), accounting for about 5% of all gastrointestinal bleeding cases, often originates in the small intestine. Single-balloon enteroscopy (SBE) has recently been introduced as a diagnostic and therapeutic tool for OGIB but has a lower rate of complete small bowel examination. It remains unclear whether complete enteroscopy with SBE leads to better clinical outcomes. Some studies suggest that complete examinations significantly reduce rebleeding rates. Based on this, we hypothesize that complete SBE results in better clinical outcomes regarding rebleeding events compared to incomplete SBE. To investigate this, we conducted a multicenter observational study comparing rebleeding rates between patients undergoing complete and incomplete small bowel examination with SBE, aiming to evaluate the diagnostic value and clinical impact of complete enteroscopy in OGIB management.

#### Who can participate?

Patients who are undergoing single-balloon enteroscopy for obscure gastrointestinal bleeding.

#### What does the study involve?

Before undergoing the single-balloon enteroscopy (SBE) procedure, patients receive standard bowel preparation with 3 liters of polyethylene glycol solution administered 8 hours prior to the procedure. SBE is performed under conscious sedation or general anesthesia by experienced endoscopists, each of whom has performed at least 50 SBE procedures. The SIF-Q260 enteroscope (Olympus, Tokyo, Japan) is used in all cases. Data on clinical presentation, diagnosis, and whether the SBE is classified as complete or incomplete are collected. Patients are then categorized into two groups: complete and incomplete SBE. They are followed for re-bleeding events, with data obtained from medical records and telephone interviews. Clinical presentation, diagnosis, and interventions at the time of re-bleeding are recorded and compared between the two groups.

What are the possible benefits and risks of participating?

There are no direct benefits or risks to patients participating in this study, as all participants will

continue to receive the standard best care. However, this study may provide valuable information on the clinical impact of complete and incomplete SBE in patients with OGIB.

Where is the study run from?

- 1. The First Affiliated Hospital of Zhengzhou University, Zhengzhou, China
- 2. Henan Provincial People's Hospital, Zhengzhou, China
- 3. Qinghai Provincial People's Hospital, Xining, China

When is the study starting and how long is it expected to run for? November 2022 to April 2025

Who is funding the study? Investigator initiated and funded

Who is the main contact? Fangbin Zhang, MD, PhD, fcczhangfb@zzu.edu.cn

### **Contact information**

#### Type(s)

Public, Scientific, Principal investigator

#### Contact name

Dr Fangbin Zhang

#### **ORCID ID**

https://orcid.org/0000-0002-8680-6413

#### Contact details

No. 1 Jianshe East Road Zhengzhou China 450052 (+86)13526655133 fcczhangfb@zzu.edu.cn

# Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

Nil known

# Study information

Scientific Title

Clinical outcomes of complete versus incomplete small bowel examination by single-balloon enteroscopy in obscure gastrointestinal bleeding: a prospective multicenter study

#### **Study objectives**

A complete small bowel examination leads to better clinical outcomes than an incomplete examination, including more accurate diagnosis and a lower risk of re-bleeding.

#### Ethics approval required

Ethics approval required

#### Ethics approval(s)

approved 16/11/2022, Medical Ethics Committee of the First Affiliated Hospital of Zhengzhou University (No. 1 Jianshe East Road, Zhengzhou, 450052, China; +86 37166295219; yangzhgcp@163.com), ref: 2022-KY-1127

#### Study design

Multicenter observational study (retrospective analysis of prospectively collected data)

#### Primary study design

Observational

#### Study type(s)

Diagnostic

#### Health condition(s) or problem(s) studied

Patients with obscure gastrointestinal bleeding are defined as those who have overt bleeding such as hematemesis, melena, or hematochezia, and have negative results on both upper endoscopy and colonoscopy.

#### **Interventions**

This study follows patients with obscure gastrointestinal bleeding who undergo either complete or incomplete small bowel examination using single-balloon enteroscopy between January 2016 and December 2024. Follow-up data on re-bleeding events are collected from medical records and telephone interviews. The aim is to compare the clinical outcomes between the two groups.

#### Intervention Type

Procedure/Surgery

#### Primary outcome(s)

Re-bleeding events, defined as the recurrence of overt gastrointestinal bleeding such as hematemesis, melena, or hematochezia after the initial complete or incomplete single balloon enteroscopy. Clinical presentation, diagnosis, and interventions at the time of re-bleeding are collected from patient records

#### Key secondary outcome(s))

Clinical findings and diagnoses from the initial single balloon enteroscopy, either complete or incomplete, are collected from patient records

#### Completion date

25/04/2025

# **Eligibility**

#### Key inclusion criteria

- 1. A confirmed diagnosis of obscure gastrointestinal bleeding (OGIB)
- 2. Admission to one of the following institutions: The First Affiliated Hospital of Zhengzhou University, Henan Provincial People's Hospital, or Qinghai Provincial People's Hospital
- 3. Underwent single-balloon enteroscopy (SBE) between January 2016 and December 2024

#### Participant type(s)

Patient

#### Healthy volunteers allowed

No

#### Age group

Adult

#### Lower age limit

18 years

#### Upper age limit

80 years

#### Sex

All

#### Key exclusion criteria

- 1. Declined to participate in the follow-up
- 2. Were lost to follow-up during the study period

#### Date of first enrolment

10/04/2025

#### Date of final enrolment

20/04/2025

## Locations

#### Countries of recruitment

China

# Study participating centre The First Affiliated Hospital of Zhengzhou University

No. 1 Jianshe East Road Zhengzou China 450052

#### Study participating centre Henan Provincial People's Hospital

No.7 Weiwu Road Zhengzhou China 450003

Study participating centre Qinghai Provincial People's Hospital

No.2 Gonghe Road Xining China 810007

# Sponsor information

#### Organisation

The First Affiliated Hospital of Zhengzhou University

# Funder(s)

#### Funder type

Other

#### **Funder Name**

Investigator initiated and funded

# **Results and Publications**

#### Individual participant data (IPD) sharing plan

The current data sharing plans for this study are unknown and will be available at a later date

#### IPD sharing plan summary

Data sharing statement to be made available at a later date

#### **Study outputs**

Output type **Details** Date created Date added Peer reviewed? Patient-facing? Participant information sheet

Participant information sheet

11/11/2025 11/11/2025 No