Better care pathways in pregnancies after stillbirth

Submission date	Recruitment status No longer recruiting	[X] Prospectively registered		
12/02/2018		[X] Protocol		
Registration date 13/02/2018	Overall study status Completed	Statistical analysis plan		
		[X] Results		
Last Edited 17/08/2022	Condition category Pregnancy and Childbirth	[] Individual participant data		
11/00/2022	r regulaticy and entiablian			

Plain English summary of protocol

Background and study aims

In the UK, 13 babies die shortly before or soon after birth every day, causing long lasting grief for parents. Most women who have had a stillbirth or neonatal death conceive again, often soon after the loss. Subsequent pregnancies are associated with high stress and anxiety for parents, increasing complications such as the baby being born too early or too small. Difficult feelings and emotions often persist, even after the birth of a healthy baby and can disrupt early bonding, making family and social problems more likely later in life. Sensitive and appropriate support is vital for parents in pregnancy after the death of a baby. However, our recent research demonstrated that not all parents had good care all of the time; tactless and insensitive communication and lack of emotional support were

common issues. This study examines whether it is possible to conduct research testing a new package of care to improve support for parents who have previously experienced death of a baby. Women would have a named midwife care-coordinator, who would deliver antenatal care alongside their doctor, maintain a relationship with the family in pregnancy and during the early days after birth and provide access to extra support. This change will be introduced with a small group of women in two hospitals in North-West England and compared with similar women who received care immediately prior to the change. This study assesses whether parents are willing to take part and stay in the research study, whether the change works as planned and the best ways of assessing the effect on well-being and maternity services. If this is study is successful then a larger study will be undertaken to assess whether this change would benefit women, represents good value for money and should be introduced to the NHS.

Who can participate?

Women aged 16 and older who are less than 20 weeks gestation and have experienced a stillbirth or a neonatal death of a previous baby and their partners. Health workers involved directly or indirectly involved in those providing care to women can also participate.

What does the study involve?

This study includes different phases. The first one to three to months include the study set up. The next three to eight months participants are recruited. During this time, they receive the existing care provided for women after previous death of a baby. Months nine to twenty consists of the second phase of recruitment where the new care package is introduced. The women

recruited during this phase receive a midwife care coordinator and a buddy widwife that provides support. Participants are able to contact their buddy and have access to a WhatsApp group. During the study, all participants are followed up at 30-37 weeks of pregnancy and 4-6 weeks after birth. Participants are asked about their anxiety, depression, health status and maternal-infant bonding. Women are also provided with a diary for them to record their thoughts and feelings. There are also interviews with participants, their partners and health care workers.

What are the possible benefits and risks of participating?

There are no direct benefits intended for participants during phase 1 of this study. During phase 2 it is hoped that the additional support delivered for the intervention will address deficiencies in care identified for women and families in previous exploratory work but this is cannot be guaranteed. Perinatal loss is an extremely sensitive area of maternity care with potential for women, partners and families affected by the death of baby and participating in research to suffer emotional distress. However, evidence suggests that women and families who have experienced the death of a baby around the time of birth find participation in research, particularly interviews helpful in allowing them to discuss events and their care in depth.

Where is the study run from?

- 1. Burnley General Hospital (Burnley)
- 2. Royal Oldham Hospital (Manchester)

When is the study starting and how long is it expected to run for? January 2018 to December 2020

Who is funding the study? National Institute for Health Research (UK)

Who is the main contact? Dr Tracey Mills (Scientific) tracey.mills@lstmed.ac.uk

Contact information

Type(s)

Scientific

Contact name

Dr Tracey Mills

Contact details

Room 4.334 Jean Macfarlane Building School of Nursing, Midwifery and Social Work The University of Manchester Oxford Road Manchester United Kingdom M13 9PL +44 161 306 7709 tracey.mills@lstmed.ac.uk

Additional identifiers

Protocol serial number 36987

Study information

Scientific Title

Better maternity care pathways in pregnancies after stillbirth or neonatal death: A feasibility study

Study objectives

The aim of this study is to assess the feasibility of conducting a large-scale multicentre randomised controlled trial to test an intervention to improve psychological well being of parents who have previously experienced a

stillbirth (a baby born after 24 weeks of pregnancy, showing no signs of life) or neonatal death (a baby born alive, but died within first 28 days of life), through an improved maternity care pathway for the next pregnancy. Improving emotional and psychological support during subsequent pregnancies was ranked among the top ten research priorities identified by the recent James Lind Alliance Stillbirth Priority Setting Partnership. This multicomponent intervention, developed following our exploratory studies with parents and professionals, combines provision of a named midwife care coordinator and access to additional individual and group support activities during pregnancy after stillbirth or neonatal death. This package has been designed to improve continuity of care, foster emotional well being and enhance peer networks for parents during subsequent pregnancies.

Ethics approval required

Old ethics approval format

Ethics approval(s)

North West Greater Manchester West Research Ethics Committee, 17/01/2018, ref: 18/NW/0010

Study design

Non-randomised; Both; Design type: Process of Care, Complex Intervention, Management of Care, Cohort study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Maternity care during pregnancy after stillbirth or neonatal death

Interventions

This prospective, observational mixed-methods study determines the feasibility of conducting a large trial of a new care package to improve the emotional and psychological support for

parents in pregnancies after stillbirth or neonatal death. A pre- and post-observational cohort study over 36 months is conducted to allow assessment of important elements of the processes involved in a definitive trial within the limits of a feasibility study.

A total sample of 60 women (20 in the control phase (phase 1) and 40 in the intervention phase (phase 2) are recruited.

Partners (estimated up to 40 in total) are also recruited to explore experiences of subsequent pregnancy and the intervention through questionnaires and interviews. Birth partners (not woman's spouse or domestic partner) are recruited to take part in interviews where they are the main source of support for the woman. Health professionals (midwives, obstetricians, service managers) involved in care of women in pregnancy after stillbirth or neonatal death and in the delivery of the intervention in the two sites will also be recruited for interview (up to 15).

The first 1-3 months include the study set up. Months 3-8 phase 1 recruitment includes the usual care.

Over a six month period 10 women and partners in each of the two study sites (up to 20 weeks gestation who have previously experienced a stillbirth/neonatal death) are identified and approached via the clinical care team and recruited by the research midwife.

No national care pathway exists for women over pregnancy after stillbirth or neonatal death, therefore understanding of 'usual care' is limited. Planned data collection during the control phase allows detailed documentation of usual care in the participating sites. Participants recruited during months 3-8 have existing care pathway provided for women after previous perinatal death. Antenatal care normally be consultant-led, with appointments in the hospital and community clinics. Additional monitoring and investigations are requested according to clinical need, following national guidance. The standard antenatal education programme are available. Intrapartum care are provided in the obstetric unit and postnatal care by the community midwifery team, until transfer to primary care, up to 28 days after birth.

Recruitment is completed by end of month 8.

Months 9-20 consists of the Phase 2 recruitment: Intervention introduced, new care package

Over a twelve month period, 20 women per site (up to 20 weeks gestation who have previously experienced a stillbirth/neonatal death) are identified, approached and recruited as above to receive a new care package.

This involves being allocated a midwife care co-ordinator, with experience in caring for bereaved families and training in the intervention to improve continuity of care. The care co-ordinator works with a 'buddy' midwife also trained to provide the intervention to support and cover annual leave sickness and absence. The care co-ordinator also gets to know the woman and family through providing midwifery care during pregnancy as often as possible. They help the woman to make a plan for her pregnancy and birth, work with her obstetrician and other professionals to communicate clearly about care. They are also contactable at specified times should the woman need additional support or information for non-urgent concerns. The care co-ordinator remains in contact after the birth and during the early postnatal period to provide support eg by phone, although she may not directly provide postnatal care. The care co-ordinator midwives are trained by the study health psychologist to provide self-help advice to support women in coping with difficult feelings through a 'Coping Strategies Toolbag'. Women in the intervention group are also offered access to a monthly support group facilitated by midwives from the research and clinical teams on each study site to promote peer support.

Session content will be flexible according to women's needs but includes targeted antenatal education appropriate to this group. A study 'WhatsApp' group are also available for women, moderated by the research team.

Data collection: During the study all the women participants (phase 1 and 2) meets with the research midwife at recruitment, 30-37 weeks and 4 to 6 weeks after birth. During pregnancy the appointments takes no more than 1.5 hours, the postnatal appointment would not be expected to last more than two hours.

Recruitment and retention data: Data on numbers of women who fulfill the eligibility criteria, who are invited to participate, the number recruited and women who leave the study before completion are kept. With permission, reasons for non-recruitment will also be recorded for women/ partners who have given contact details.

Demographic and clinical data: At the appointments the research midwife study completes case report forms using medical records and self-report to collect women's information including demographics, medical and social history, previous and current pregnancy data including all healthcare use and maternal and infant outcomes after the birth.

Psychological assessments: Women and partners (if participating) are also asked to complete a questionnaire containing several validated tools to assess a range of relevant psychological constructs including anxiety, depression, worry, health status and in the postnatal period, maternal-infant bonding and maternal self-efficacy.

Experiences of maternity care subsequent pregnancy, the care package and participation in the research.

Women are provided with a diary at recruitment. This has sections for recording thoughts and feelings about their pregnancy and care on a daily or weekly basis. Participants are informed that it is their choice whether to use these. They are collected at the end of the study.

Views and experiences are explored through semi-structured qualitative interviews with a research assistant, in person, in a place chosen by the participant (e.g. home or private room in the hospital) or on the telephone if preferred.

Experience with previous studies indicates interviews with up to 12 women participating in phase 1, and up to 25 women participating in phase 2 are required to achieve data saturation, where no new ideas emerge. For partners up to 8 participating in phase 1, and 12 participating in phase 2 are recruited. Interviews with women and partners

are scheduled for 4-6 weeks after the birth and if both partners agree they are interviewed separately. As not all women and partners will wish to participate or be available for an interview, all participants are approached at recruitment to ascertain their willingness. The study team will recruit up to 15 health professionals caring for women or

managing services during the study, they are approached directly by the research team towards the end of the data collection period in year 3.

Intervention Type

Behavioural

Primary outcome(s)

1. Recruitment rates of women and partners recorded as the number of eligible participants who consent to take part, measured using the participant log, at the end of the study.

2. Retention of women and their partners recorded as the number of participants who remain in the study until completion, measured using the participant log, at the end of the study

Key secondary outcome(s))

- 1. Experiences of study participation and the intervention of parents and healthcare staff to determine the acceptability of trial processes and of the intervention. Explored through qualitative interviews, 4-6 weeks after birth (women and partners) and at the end of data collection (health professionals)
- 2. The characteristics of proposed psychological, cost-effectiveness, utility and clinical outcome measures will be examined in preparation for a definitive trial. These will be assessed by completion of validated self-report questionnaires (Generalised Anxiety Disorder 7 item scale (GAD-7) completed by women and partners at recruitment, 30-37 weeks and 4-6 weeks after the birth. Cambridge Worry Scale completed by women only at recruitment and 30-37 weeks gestation. Edinburgh Postnatal Depression Score completed by women only at recruitment, 30-37 weeks and 4-6 weeks after birth. Maternal-Infant Bonding scale completed by women only at 4-6 weeks after the birth. Maternal Self Efficacy Scale, (MSES) an 8-item scale to assess women's self-belief in their abilities to meet the needs of their infant, will be completed by women at 4-6 weeks after birth. Investigator designed case report forms (clinical outcomes) completed at recruitment, 30-37 weeks gestation and 4-6 weeks postnatal.
- 3. Elements of process evaluation will assess fidelity and quality of implementation, clarify causal mechanisms and identify contextual factors impacting on outcomes. Fidelity will be assessed using the intervention logs completed by care co-ordinators during phase two, case report forms completed at 30-37 weeks gestation and 4-6 weeks after the birth and interviews with women and partners 4-6 weeks after birth and health professionals at the end of data collection

Completion date

31/12/2020

Eligibility

Key inclusion criteria

- 1. Women and Partners:
- 1.1. Pregnant women over 16 years old and ≤20 weeks' gestation and have experienced a stillbirth or a neonatal death of any previous baby
- 1.2. Partners, approached through the woman after she has agreed (a partner's unwillingness to participate will not affect the woman's continued participation)
- 1.3. Booked and planning have antenatal care at East Lancashire Hospitals NHS Trust or Pennine Acute Trust
- 1.4. For feasibility, sufficient command of English to participate in interviews and complete questionnaires will be required. Translation of materials will be explored with the aim of including parents lacking fluency in English in the definitive trial.
- 2. Health workers:
- 2.1. Midwives, obstetricians, service managers directly/indirectly involved in providing care to women participating in the study

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Total final enrolment

78

Key exclusion criteria

- 1. Pregnant women/partners under 16 years
- 2. Women/partners who are already receiving care through a specialist antenatal service (eg diabetes, haematology clinics) as they would be receiving specialist midwifery/obstetric care relevant to their complex history.
- 3. Participants who are unable or unwilling to consent

Date of first enrolment

01/03/2018

Date of final enrolment

31/10/2019

Locations

Countries of recruitment

United Kingdom

England

Study participating centre Burnley General Hospital

Casterton Avenue Burnley United Kingdom BB10 2PQ

Study participating centre Royal Oldham Hospital

Rochdale Road Oldham Manchester United Kingdom OL1 2JH

Sponsor information

Organisation

The University of Manchester

ROR

https://ror.org/027m9bs27

Funder(s)

Funder type

Government

Funder Name

National Institute for Health Research

Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

The data sharing plans for the current study are unknown and will be made available at a later date.

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		10/08/2022	12/08/2022	Yes	No
HRA research summary			28/06/2023	No	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes