

Efficacy of non-thermal gas plasma on sub-clinical wound infection (biofilm) in patients with diabetic ulcers

Submission date 14/10/2016	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 20/10/2016	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 02/06/2023	Condition category Circulatory System	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

People living with diabetes often have to live with long-term complications of the disease. One of the most common, disabling and costly complications are foot ulcers which do not heal. Infections of diabetic foot ulcers are a major cause of amputation and death. In many cases, the infection is resistant to antibiotic treatment, which can make them very difficult to treat. The infected wounds are also slow to heal due to a layer of bacteria on the surface of the wound called biofilms. Non-thermal gas plasma (NTGP) is known to be an effective and safe treatment for chronic (long term) infected wounds and does not lead to antibiotic resistance. This type of treatment consists of a five minute 'blast' of gas plasma held over the infected wound. The treatment head on the device contains a patented ionization chamber that bombards Argon gas (an unreactive gas) with electrons (negatively charged particles) emitted from multiple hot electric filaments. The resulting plasma ions mix with air creating reactive agents to generate a wide, constant treatment field that is capable of treating large tissue areas. The aim of this study is to find out if NTGP is an effective treatment for diabetic foot ulcers and establish a solution to a real life situation.

Who can participate?

Adults with diabetes, who have foot ulcers where healing is stalled for at least 4 weeks.

What does the study involve?

Participants are randomly allocated to one of two groups. Those in the first group receive standard wound care dressing twice per week. This involves having the wound cleaned with sterile water or normal saline (salt water) and then dressed with the usual dressing used at that site. Those in the second group also receive standard wound care dressing twice per week with the addition of NTGP. This involves five minutes of applying argon gas directly over the wound after it has been cleaned, before it is dressed. Participants in both groups have samples taken from the wound at the start of this study and then after 4 weeks in order to measure the amount of bacteria present on the surface of the wound. Photographs are also taken every week for four weeks in order to assess healing.

What are the possible benefits and risks of participating?

Participants who receive the NTGP treatment may benefit from improved quality of life as the technology should reduce stress, trauma and recovery time. There are no notable risks involved with participating.

Where is the study run from?

Leeds Teaching Hospital NHS Trust, UK

(updated 17/07/2019, previously: Salford Royal NHS Foundation Trust (UK))

When is study starting and how long is it expected to run for?

August 2016 to January 2018

Who is funding the study?

1. Innovate UK (UK)

2. Adtec Europe Limited (EU)

Who is the main contact?

Mrs Mary McGovern

clinical@adtec.eu.com

Contact information

Type(s)

Public

Contact name

Mrs Mary McGovern

Contact details

Adtec Europe Limited

Unit 1 Alice Way

Hounslow

London

United Kingdom

TW3 3UD

+44 208 737 5500

clinical@adtec.eu.com

Additional identifiers

Protocol serial number

59178-445167

Study information

Scientific Title

A randomised controlled trial to evaluate the efficacy of non-thermal gas plasma (NTGP) on sub-clinical wound infection (biofilm) in patients with diabetic foot ulcers compared to those treated with standard of care dressings

Acronym

Biofilm Project

Study objectives

The healing of chronic wounds that are stalled by sub clinical wound infection (biofilm) can be accelerated following intervention with non-thermal gas plasma.

Ethics approval required

Old ethics approval format

Ethics approval(s)

East Midlands - Leicester South Research Ethics Committee, 20/01/2017, ref: 16/EM/0476
Health Research Authority, 25/01/2017, IRAS project ID: 198288

Study design

Single-centre prospective randomised controlled trial

Primary study design

Interventional

Study type(s)

Prevention

Health condition(s) or problem(s) studied

Chronic diabetic foot ulcers

Interventions

Participants are randomised to one of two groups using sealed, opaque sequentially numbered envelopes (SNOSE)

Intervention group: Participants receive local standard of care wound dressing (SOCD) plus 5 minutes treatment of NTGP twice weekly over a period of 30 days (4 weeks). SOCD involves cleansing the wound with sterile water or normal saline and then dressed with the local standard of care dressing e.g. Aquacel Foam non-adhesive, Aquacel Foam adhesive, Biatain foam non adhesive, Biatain Foam adhesive. NTGP involves a two minute treatment of argon gas plasma directly over the wound.

Control group: Participants only receive local standard of care wound dressing (SOCD) twice weekly. This involves cleansing the wound with sterile water or normal saline and then dressed with the local standard of care dressing e.g. Aquacel Foam non-adhesive, Aquacel Foam adhesive, Biatain foam non adhesive, Biatain Foam adhesive.

Follow up for all study participants will occur after the 4 weeks, last samples and digital photographs will be taken at this time and participants will be informed when the results of the study will be completed and published.

Intervention Type

Device

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

-

Primary outcome(s)

Wound volume is measured using digital photographic imaging that will record, and measure the wound bed tissue type, wound dimensions and wound volume at baseline, 1, 2, 3 and 4 weeks.

Key secondary outcome(s)

1. The effect of gas plasma in regards to accelerated healing is measured using wound tissue biopsy's and wound slough samples taken from the 60 patients at baseline and week 4
2. Presence of biofilms is measured as the numbers of bacteria and comparison to bacteria quantified from the wound surface, using wound tissue biopsy's and wound slough samples, at baseline and week 4
3. Microbial populations are measured using digital photographic imaging, at baseline week 1, week 2, week 3, week 4 as well as wound tissue biopsy and wound slough samples at baseline and week 4

Completion date

31/05/2021

Eligibility

Key inclusion criteria

Inclusion criteria as of 04/04/2017:

1. Male or female,
2. Aged 18 years and over
3. Diabetics with HbA1c less than 10% (<86mm/mol) (Amended 12/09/2017 to Diabetics with HbA1c less than 12% (<108mm/mol)) that has been recorded within the previous 3 months
4. Foot ulcers with University of Texas grade/size A1 or A2 or B1* sited below the ankle including plantar, dorsal and heel ulcers
*Texas grade B1 ulcers which have symptoms consistent with mild diabetic foot infection (IDSA 2012)¹⁴ including:
 - 4.1. Pus or inflammation
 - 4.2. Inflammation extends less than 2cm from the wound
 - 4.3. Infection is limited to skin/soft tissue
 - 4.4. Systemically well
5. Ulcers with dimension 1.0cm² – 30cm²
6. Ulcer where there is less than 40% decrease in wound surface area in the previous 4 weeks
7. Ulcer has not been present for more than 2 years
8. If foot pulses not palpable the toe brachial pressure index (TBPI) should be greater than 0.5
9. If TBPI cannot be assessed then the ankle brachial pressure index (ABPI) should be 0.8 or above
10. If ABPI is 1.3 or above the Doppler sounds should be at least biphasic

Original inclusion criteria:

1. Aged 18 years and over
2. Diabetics with HbA1c less than 10% (<86mm/mol) that has been recorded within the previous 3 months.
3. Foot ulcers with University of Texas grade/size A1, A2, B1 sited below the ankle including plantar, dorsal and heel ulcers.

4. Diabetics whose wound/ulcer currently, or in the last 7 days, has symptoms consistent with mild diabetic foot infection (IDSA 2012)¹⁴ including:
- 4.1. Pus or inflammation
 - 4.2. Inflammation extends less than 2cm from the wound
 - 4.3. Infection is limited to skin/soft tissue
 - 4.4. Systemically well
5. Ulcers with dimension 1.0cm² – 30cm²
6. Ulcer where there is less than 40% decrease in wound surface area in the previous 4 weeks
7. Ulcer has not been present for more than 2 years
8. Adequate blood supply determined by palpable foot pulses or if not palpable a TBPI greater than 0.7
9. Loss of protective sensation to a 10g monofilament
10. Male or female

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

21

Key exclusion criteria

Exclusion criteria as of 04/04/2017:

1. HbA1c greater than 10% (>86mm/mol) Amended 12/09/2017: to HbA1c greater than 12% (>108mm/mol)
2. Those with malignancy or other immunosuppressive diseases
3. Those receiving radiotherapy or medications that actively delay healing (e.g. steroids, antimetabolites)
4. Those whose wound/ulcer currently has symptoms consistent with moderate diabetic foot infection (IDSA 2012¹⁴) – including
 - 4.1. Pus or inflamed wound in a patient who is systemically well and/or one of the following:
 - 4.2. Inflammation extends greater than 2cm from wound
 - 4.3. Lymphangitis
 - 4.4. Localised necrosis/ gangrene
 - 4.5. Involvement of muscle, tendon, joint or bone (active osteomyelitis)
5. Those whose wound/ulcer currently has symptoms consistent with severe diabetic foot infection (IDSA, 2012) as demonstrated by:
 - 5.1. Extensive cellulitis,
 - 5.3. Deep abscess with or without signs of systemic toxicity (fever, vomiting, hypotension, confusion, acidosis, renal failure, severe hyperglycaemia, leucocytosis)

6. Pregnant, or breast feeding
7. Women who are of child-bearing age not using reliable contraception
8. TBPI less than 0.5
9. ABPI less than 0.8
10. If ABPI above 1.3 and vessel sounds are monophasic
11. Those who have clinical evidence of gangrene at any location
12. Those who has a medical condition that in the opinion of the investigator would make the patient an inappropriate candidate for the study
13. Those who have necrotic toes on the study ulcer foot
14. Those who have undergone surgical procedure other than debridement on the study ulcer foot within 3 weeks prior to screening
15. Study ulcer over active Charcot's joint
16. Non study ulcer within 5.0 cm from the study ulcer at enrolment
17. Participation in other clinical study in the last 4 weeks

Original exclusion criteria:

1. HbA1c greater than 10% (>86mm/mol)
2. Those with malignancy or other immunosuppressive diseases
3. Those receiving radiotherapy or medications that actively delay healing (e.g. steroids, NSAIDS, antimetabolites)
4. Those whose wound/ulcer currently (or in the last 7 days) has symptoms consistent with moderate diabetic foot infection (IDSA 2012/14) – including:
 - 4.1. Pus or inflamed wound in a patient who is systemically well and/or one of the following:
 - 4.2. Inflammation extends greater than 2cm from wound
 - 4.3. Lymphangitis
 - 4.4. Localised necrosis/ gangrene
 - 4.5. Involvement of muscle, tendon, joint or bone (active osteomyelitis)
5. Those whose wound/ulcer currently (or in the last 7 days) has symptoms consistent with severe diabetic foot infection (IDSA, 2012) as demonstrated by:
 - 5.1. Extensive cellulitis,
 - 5.2. Deep abscess with or without signs of systemic toxicity (fever, vomiting, hypotension, confusion, acidosis, renal failure, severe hyperglycaemia, leukocytosis)
6. Pregnant, or breast feeding
7. Women who are of child-bearing age not using reliable contraception
8. TBPI less than 0.7
9. Those who have clinical evidence of gangrene at any location
10. Those who have a medical condition (e.g. diabetic nephropathy) that in the opinion of the investigator would make the patient an inappropriate candidate for the study
11. Those who have necrotic toes on the study ulcer foot
12. Those who have undergone surgical procedure other than debridement on the study ulcer foot within 3 weeks prior to screening
13. Study ulcer over active Charcot's joint
14. Non study ulcer within 5.0 cm from the study ulcer at enrolment
15. Participation in other clinical study in the last 4 weeks

Date of first enrolment

06/02/2017

Date of final enrolment

31/12/2020

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

St James's Hospital

Leeds Teaching Hospital NHS Trust

Beckett Street

Leeds

United Kingdom

LS9 7TF

Sponsor information

Organisation

Adtec Europe Limited

ROR

<https://ror.org/059gjhp96>

Funder(s)

Funder type

Government

Funder Name

Innovate UK

Alternative Name(s)

Technology Strategy Board

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Funder Name

Adtec Europe Limited

Results and Publications

Individual participant data (IPD) sharing plan

The current data sharing plans for the current study are unknown and will be made available at a later date

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		08/12/2022	02/06/2023	Yes	No
HRA research summary			28/06/2023	No	No