# Early presbyopia population characterisation

Submission date 17/07/2025	Recruitment status No longer recruiting	Prospectively registered	
		□ Protocol	
Registration date 29/07/2025	Overall study status Completed	Statistical analysis plan	
		Results	
<b>Last Edited</b> 29/07/2025	<b>Condition category</b> Eye Diseases	[] Individual participant data	
		[X] Record updated in last year	
Plain English summary of protocol			
Background and study aims			
The onset of presbyopia is a continuous physiological phenomenon. Typically, the timing of the			
		yith contact lenses or spectacles, is when patients	
report near vision-associated symptoms. This study aims to characterise the early presbyopic			
population with a view to identify factors likely to contribute to the difficulty in transferring			

## Who can participate?

Early presbyopes aged between 40 to 50 years old with single vision correction or no vision correction.

from single vision to multifocal vision correction and to differentiate between asymptomatic

# What does the study involve?

The study involves attending one study visit and completing a set of remote questionnaires in 10 days following the study visit.

#### What are the possible benefits and risks?

The study results will contribute to additional information on early presbyopia.

adaptors during the period preceding the use of a multifocal correction.

No risks given at registration

Where is the study running from?
Ocular Technology Group - International (OTG-i) (UK)

When is the study starting and how long is it expected to run for? March 2025 to September 2025

Who is funding the study? CooperVision International Limited (UK)

Who is the main contact?

Deborah Moore, dmoore@otq.co.uk

# Contact information

#### Type(s)

Scientific, Principal investigator

#### Contact name

Dr Michel Guillon

#### Contact details

66 Buckingham Gate London United Kingdom SW1E 6AU +44 (0)2072224224 MGuillon@otg.co.uk

#### Type(s)

**Public** 

#### Contact name

Miss Deborah Moore

#### Contact details

66 Buckingham Gate London United Kingdom SW1E 6AU +44 (0)2072224224 dmoore@otg.co.uk

# Additional identifiers

## Clinical Trials Information System (CTIS)

Nil known

# Integrated Research Application System (IRAS)

354245

# ClinicalTrials.gov (NCT)

Nil known

#### Protocol serial number

CV24-52

# Study information

#### Scientific Title

Early presbyopia program phase 1 population characterisation

# **Study objectives**

The study will be an observational study to increase knowledge on early presbyopia, therefore, no specific hypothesis will be tested, however, the study results will make it possible to develop specific hypotheses to be tested to improve the management of early presbyopia.

## Ethics approval required

Ethics approval required

## Ethics approval(s)

approved 30/05/2025, London-Brent Research Ethics Committee (2 Redman Place, Stratford, London, E20 1JQ, United Kingdom; +44 (0)2071048128, 207 104 8117, 2071048131; brent. rec@hra.nhs.uk), ref: 25/PR/0578

#### Study design

Single-visit in-clinic prospective observational study

### Primary study design

Observational

#### Study type(s)

Other

### Health condition(s) or problem(s) studied

Early presbyopia

#### **Interventions**

The study will be a single-visit, in-clinic, prospective observational study followed by three remote time-controlled questionnaires implemented on three different days within the following 10-day period.

The participants will attend the clinic wearing their habitual visual correction and will bring their electronic devices (e.g. laptops, tablets, electronic books, smartphones) to quantify their near environment needs. The potential participants will be screened and enrolled on the study. Once enrolled, the participant will complete a set of questionnaires, and then the investigator will carry out a set of non-invasive clinical measurements. At the end of the visit, a remote electronic controlled questionnaire will be set up.

The participants will complete three remote electronic questionnaires each on three different days within the following 10-day period.

#### Intervention Type

Other

## Primary outcome(s)

- 1. Optical and visual characteristics of the early presbyopic population measured using time-controlled visual acuity at two points during the study visit
- 2. Identify differentiating factors between symptomatic and asymptomatic early presbyopes measured using subjective questionnaires during the study visit and remotely after the study visit

## Key secondary outcome(s))

There are no secondary outcome measures

# Completion date

30/09/2025

# Eligibility

#### Key inclusion criteria

The entry criteria will be:

- 1. Age 40 to 50 years
- 2. Vision correction:
- 2.1. Distance Vision Correction only
- 2.1.1. Single vision distance spectacles
- 2.1.2. Single vision distance contact lenses
- 2.1.3. No vision correction
- 3. Spectacle refraction:
- 3.1. Distance: Sphere: -6.00D to + 3.00D
- 3.2. Astigmatism: 0.00D to -2.50
- 4. Best corrected distance visual acuity of at least 20/25 in each eye

#### Participant type(s)

Healthy volunteer

#### Healthy volunteers allowed

No

#### Age group

Adult

#### Lower age limit

40 years

#### Upper age limit

50 years

#### Sex

Αll

#### Key exclusion criteria

- 1. Acute and subacute inflammation or infection of the anterior chamber of the eye
- 2. Any eye disease, injury or abnormality that affects the cornea, conjunctiva or eyelids that would contraindicate contact lens wear
- 3. Corneal hypoesthesia (reduced corneal sensitivity), if not aphakic
- 4. Severe insufficiency of lacrimal secretion (dry eyes)
- 5. Any systemic disease that may affect the eye or may be exaggerated by wearing contact lenses (e.g. acne and eczema)
- 6. Allergic reactions of ocular surfaces or adnexa that may be induced or exaggerated by wearing contact lenses or use of contact lens solutions
- 7. Any active corneal infection (bacterial, fungal, protozoal or viral)
- 8. Newly prescribed (within the past 30 days) use of some systemic medications (such as antihistamines, decongestants, diuretics, muscle relaxants, tranquilizers, stimulants, antidepressants, anti-psychotics, oral contraceptives) or new prescription eyedrops which is not

rewetting/lubricating eyedrops for which contact lens wear could be contraindicated as determined by the investigator

- 9. Monocular participants (only one eye with functional vision) or participants fit with only one contact lens
- 10. Subjects with slit lamp findings greater than grade 1 (e.g. edema, infiltrates, corneal neovascularization, corneal staining, tarsal abnormalities, conjunctival, anterior segment inflammation) as recorded during the clinical visit, any previous history or signs of a contact lens related corneal inflammatory event (past corneal ulcers), or any other ocular abnormality that may contraindicate contact lens wear at the enrolment visit
- 11. History of corneal refractive surgery
- 12. Enrolment of the family members of the investigator, family members of the investigator's staff, or individuals living in the households of these individuals.

# Date of first enrolment 05/06/2025

Date of final enrolment 30/09/2025

# Locations

# **Countries of recruitment** United Kingdom

**England** 

Study participating centre
Ocular Technology Group International
66 Buckingham Gate
London
United Kingdom
SW1E 6AU

# Sponsor information

# Organisation

CooperVision International Limited

# Funder(s)

# Funder type Industry

#### **Funder Name**

CooperVision

# **Results and Publications**

# Individual participant data (IPD) sharing plan

The data-sharing plans for the current study are unknown and will be made available at a later date

# IPD sharing plan summary

Data sharing statement to be made available at a later date

# **Study outputs**

Output type Details Date created Date added Peer reviewed? Patient-facing?

Participant information sheet Participant information sheet 11/11/2025 11/11/2025 No Yes