Phase out as a treatment for chronic untreatable tinnitus: a double blind crossover trial

lan
t data
ast year
1

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr K M Heijneman

Contact details

Universitair Medisch Centrum Groningen Afd. Keel-,Neus-, Oorheelkunde P.O. Box 30001 Groningen Netherlands 9700 RB +31 (0)50 361 8053 k.m.heijneman@kno.umcg.nl

Additional identifiers

Protocol serial number N/A

Study information

Scientific Title

Study objectives

This study examines the effect of the Phase Out treatment on chronic, incurable tinnitus in adult subjects in comparison with placebo sound. The expectation of this study is that Phase Out treatment is effective for a longer duration and results in increased residual inhibition than placebo sound.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics approval received from the Medical Ethical Committee of Groningen on the 26th June 2007 (ref: METc2007/061).

Study design

Randomised, placebo controlled, crossover, double blinded trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Tinnitus

Interventions

A subject will receive Phase Out treatment for thirty minutes three times a week for one week and placebo sound treatment on the same regime during another. One month interval is in between these two sets of treatment. If a treatment is started, the subject fills in a report mark on the tinnitus loudness and tinnitus annoyance in the tinnitus diary every evening till three weeks after the treatment session. One week after each week of therapy a subject receives the evaluating questionnaires and will send them back after filling in.

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

The major aim of this study is disappearance (report mark) of the tinnitus lasting many hours (time).

Outcomes will be measured at weeks five and nine.

Key secondary outcome(s))

Besides the major aims, different questionnaires will be used to determine for which kind of tinnitus patients, this treatment is most effective:

1. Tinnitus Handicap Inventory (THI)

- 2. Tinnitus Reaction Questionnaire (TRQ)
- 3. Vital Exhaustion (VE) questionnaire
- 4. Hospital Anxiety and Depression Scale (HADS)
- 5. Short Form questionnaire (SF-36)
- 6. Eysenck Personality Questionnaire
- 7. Type D Personality Scale
- 8. Social Support Questionnaire (SSQ)
- 9. Tinnitus Coping Style Questionnaire (TCSQ)

Outcomes will be measured at weeks five and nine.

Completion date

01/05/2009

Eligibility

Key inclusion criteria

- 1. Subjects greater than 18 years
- 2. Unilateral or bilateral tinnitus
- 3. Predominant tone tinnitus by history
- 4. Tinnitus for minimum of three months

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

Not Specified

Key exclusion criteria

- 1. Acoustic neurinoma
- 2. Aortic/outflow tract stenosis
- 3. Pulsatile tinnitus
- 4. Pregnancy
- 5. Inability to correct use of test equipment: unable to cooperate during audiologic examination
- 6. Known tinnitus etiology, which would demand other treatment
- 7. Hearing loss greater than 60 decibel compared with standardised normal hearing on standard frequencies of a tone audiogram (250, 500, 1000, 2000, 4000 and 8000 hertz)

Date of first enrolment

01/05/2007

Date of final enrolment

01/05/2009

Locations

Countries of recruitment

Netherlands

Study participating centre
Universitair Medisch Centrum Groningen
Groningen
Netherlands
9700 RB

Sponsor information

Organisation

University Medical Centre Groningen (UMCG) (The Netherlands)

ROR

https://ror.org/03cv38k47

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

University Medical Centre Groningen (UMCG) (The Netherlands)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration