

Retrospective study in patient with soft tissue sarcoma of trunk wall and extremities who have been treated with surgery in association or not with chemotherapy and/or radiotherapy, to point out the response to preoperative treatment and outcome

Submission date 21/12/2016	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 21/02/2017	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 10/04/2017	Condition category Cancer	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Soft Tissue Sarcoma (STS) is a condition in which cancer cells form in the soft tissues of the body, such as the muscles, tendons (bands of tissue that connect muscles to bones), fat and blood vessels. STS most commonly affects tissue in the arms, legs and trunk (torso). In most cases, the first treatment used to treat STS is surgically removing the tumour. This can be done in combination with radiotherapy (using high-energy radiation beams to destroy cancer cells) or chemotherapy (anti-cancer drugs). The aim of this study is to look at the medical records of people who were diagnosed with STS and were treated surgically to see if outcomes are better for those who also had radiotherapy.

Who can participate?

Adults who have been diagnosed with STS between 2008 and 2012

What does the study involve?

Participating centres are asked to prepare a list of patients with STS who were diagnosed between 2008 and 2012. Patients would have been treated with surgery alone, surgery and radiotherapy, surgery and chemotherapy, surgery and chemo-radiation (a combination of radiotherapy and chemotherapy), or surgery and isolated limb perfusion (a type of chemotherapy injected directly into the affected limb) Information from patient charts is then collected in order to find out what treatment the participants had and whether it has been successful.

What are the possible benefits and risks of participating?

There are no direct benefits or risks involved with participating.

Where is the study run from?

1. Istituto Oncologico Veneto (Italy)
2. The Royal Marsden NHS Foundation Trust (UK)
3. Állami Egészségügyi Központ - Magyar Honvédség Központi Kórháza (Hungary)
4. Mount Sinai Hospital (Canada)
5. HM Hospitales (Spain)
6. Klinikum der Universität München (Germany)

When is the study starting and how long is it expected to run for?

October 2016 to May 2017

Who is funding the study?

Nanobiotix (France)

Who is the main contact?

Dr Eveline Boucher

eveline.boucher@nanobiotix.com

Contact information

Type(s)

Public

Contact name

Dr Eveline Boucher

Contact details

Nanobiotix

60 Rue de Wattignies

Building C

Paris

France

75012

+33 40469645

eveline.boucher@nanobiotix.com

Additional identifiers

Protocol serial number

N/A

Study information

Scientific Title

Multicenter retrospective observational chart review on treatment management of soft tissue sarcoma patients treated by surgery with or without perioperative treatment

Acronym

ROCR

Study objectives

The primary objective of this chart review is to investigate the prognostic significance of pathological complete response (pathCR*) to a pre-operative treatment in patients with STS of trunk and extremities on Disease Free Survival (DFS**) at three years.

*PathCR is defined as greater than or equal to 95% necrosis/fibrosis or less than or equal to 5% viable cells

**DFS is defined as time between diagnosis, and local or distant relapse of the disease or occurrence of any cancer or death

Ethics approval required

Old ethics approval format

Ethics approval(s)

This is a non interventional retrospective review of patient individual charts which does not need ethical approval. In the UK, the Ethics Committee of the Royal Marsden Foundation will review the study.

Study design

Retrospective observational cohort chart review

Primary study design

Observational

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Malignant soft tissue sarcoma

Interventions

Each center will be asked to prepare a list of eligible patients based on inclusion and exclusion criteria. If no database of soft tissue sarcoma (STS) patients is available, the institution data manager will be asked to prepare a list of all patients newly diagnosed between 2008 and 2012 with STS. From this list a selection of eligible patients should be performed according to inclusion and exclusion criteria.

All data are collected retrospectively. The charts are reviewed once and treatment data are collected in a eCRF. Patients must have been treated between 2008 and 2012 and followed up for 3 years, data of the treatment procedure and the follow up visits are collected at one time point.

Intervention Type

Other

Primary outcome(s)

Disease free survival is measured through medical record review 3 years after disease diagnosis.

Key secondary outcome(s)

1. Pathological response (% of necrosis, % of fibrosis, % of viable cells) is measured through review of the pathological report of the surgical specimen at the time of surgery

2. Tumor type and subtype is assessed through pathological report review at diagnosis and the time of surgery
3. Chemotherapy treatment (drug name, drug dose, treatment duration, number of cycles) is assessed by reviewing drug prescription form at each course of chemotherapy and at the end of chemotherapy treatment
4. Radiation treatment (Dose delivered, treatment duration) is assessed by reviewing medical radiation report at the end of Radiation treatment
5. Surgical procedure (conservative surgery, amputation, R0 resection, R2 resection) is assessed by reviewing the medical surgery report at the time of surgery
6. Quality of surgery (R0 resection, R1 resection, surgical margin, tumor size) is measured by reviewing pathological reports at the date of surgery
7. New tumors (local recurrence, distant recurrence, other cancer) are assessed by reviewing medical report and radiological MRI and CT scan report on the date of surgery, 6 months after surgery, 12 months after surgery, 18 months after surgery, 24 months after surgery, 30 months after surgery, 36 months after surgery
8. Quality of life (Performans status, walking distance, presence at work) is measured by reviewing medical reports on the date of diagnosis of the tumor, date of surgery, 6 months after surgery, 12 months after surgery, 18 months after surgery, 24 months after surgery, 30 months after surgery, 36 months after surgery

Completion date

15/05/2017

Eligibility

Key inclusion criteria

1. Primary Soft Tissue Sarcoma (STS) of the extremities and trunk wall newly pathologically diagnosed between 2008 and 2012 with STS.
2. Age \geq 18 years old
3. Any histology (except those specified in exclusion criteria below)
4. Any grade
5. Treated by any of these therapies for the primary tumor:
 - 5.1. Surgery alone
 - 5.2. Surgery + radiation therapy (pre, post or per-operative)
 - 5.3. Surgery + chemotherapy (pre or post-operative)
 - 5.4. Surgery + chemo-radiation (pre or post-operative)
 - 5.5. ILP (isolated limb perfusion) and surgery +/- radiation therapy

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

1. Patients with the following histological type: bone sarcoma, Kaposi's sarcoma, primitive neuroectodermal tumor, angiosarcoma, epithelioid hemangioendothelioma
2. Benign or intermediate histological subtypes: aggressive fibromatosis, desmoid tumor or dermatofibrosarcoma protuberans
3. Gastrointestinal stromal tumour (GIST)
4. STS localized in the intra-abdominal region and retroperitoneal region
5. Re-resection for non carcinological first resection (R1 or R2)
6. Patient with metastatic disease at diagnosis of STS
7. No pathological report available at diagnosis or at surgery
8. Less than 36 months (pilot and extended phase) of data available following diagnosis unless the patient died before
9. No possibility to have access to the data
10. Concurrent active therapy for another cancer (= patient receives active treatment for this cancer during the study period)

Date of first enrolment

13/01/2017

Date of final enrolment

30/04/2017

Locations

Countries of recruitment

United Kingdom

England

Belgium

Canada

France

Germany

Hungary

Italy

Poland

Spain

Study participating centre

Istituto Oncologico Veneto
Via Gattamelata 64
Padova
Italy
35128

Study participating centre
Fondazione IRCCS Istituto Nazionale dei Tumori
Via Venezian, 1
Milan
Italy
20133

Study participating centre
The Royal Marsden NHS Foundation Trust
Fulham Road
London
United Kingdom
SW3 6JJ

Study participating centre
Állami Egészségügyi Központ - Magyar Honvédség Központi Kórháza
Róbert Károly krt. 44
Budapest
Hungary
1134

Study participating centre
Mount Sinai Hospital
University of Toronto
00 University Avenue
Toronto
Canada
M5G 1X5

Study participating centre
HM Hospitales
Oña 10
Madrid
Spain
28050

Study participating centre
Klinikum der Universität München
Campus Großhadern
Medizinische Klinik Poliklinik III
Marchioninistr. 15
Munich
Germany
81377

Sponsor information

Organisation
Nanobiotix

ROR
<https://ror.org/047ts9g27>

Funder(s)

Funder type
Industry

Funder Name
Nanobiotix

Results and Publications

Individual participant data (IPD) sharing plan

The datasets collected and analysed in this study are available upon reasonable request from:
Deloitte Consulting
Clémentine Demaire
Senior consultant | Health Economics and Outcomes Research
Berkenlaan 8a, 1831 Diegem, Belgium
Mobile: + 32 470 26 93 76

IPD sharing plan summary
Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes