

Turning the tip of the stent upside down may ease urinary symptoms

Submission date 15/10/2024	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 22/10/2024	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 25/02/2025	Condition category Urological and Genital Diseases	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

This study aims to find out if changing the direction of a double-j stent placed in the ureter can reduce the uncomfortable urinary symptoms that patients often experience. The researchers want to see if turning the tip of the stent outward instead of inward makes a difference.

Who can participate?

Patients who have undergone a procedure called ureteroscopy for kidney or ureteral stones and have had a double-j stent inserted can participate in this study.

What does the study involve?

Participants will be divided into two groups based on the direction of their ureteral stent. They will fill out a questionnaire about their symptoms before and after the treatment. The researchers will compare the results to see if there is a difference between the two groups.

What are the possible benefits and risks of participating?

The possible benefit of participating is that it may help identify a better way to place stents that reduces urinary symptoms. The risks are minimal and are mainly related to the standard procedure of having a stent placed.

Where is the study run from?

The study is being conducted at Medical Park Gebze Hospital Urology Clinic (Turkey)

When is the study starting and how long is it expected to run for?

May 2023 to September 2024

Who is funding the study?

Investigator initiated and funded

Who is the main contact?

Dr. Goksel Bayar, goxelle@yahoo.com

Contact information

Type(s)

Public, Scientific, Principal Investigator

Contact name

Mr Göksel Bayar

ORCID ID

<https://orcid.org/0000-0003-1506-9732>

Contact details

Guzeller mah kavak cad medical park hastanesi

Kocaeli

Türkiye

34363

+90 5315849749

goksel.bayar@medicalpark.com.tr

Additional identifiers**EudraCT/CTIS number**

Nil known

IRAS number**ClinicalTrials.gov number**

Nil known

Secondary identifying numbers

Nil known

Study information**Scientific Title**

Reverse placement of a double J ureteral stent improve lower urinary tract symptoms

Study objectives

If we place the tip of the stent upside down, we can reduce contact with the bladder trigone. In this way, the patient's complaints can be reduced.

Ethics approval required

Ethics approval required

Ethics approval(s)

Approved 01/06/2024, Istinye university local ethics committee (Hamidiye, Kâğıthane, İstanbul, 34408, Türkiye; +90 8502836000; info@istinye.edu.tr), ref: 24-120

Study design

Prospective randomized controlled

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Quality of life, Treatment, Safety, Efficacy

Participant information sheet

No participant information sheet available

Health condition(s) or problem(s) studied

Improve lower urinary tract symptoms on patients that inserted double-j stent

Interventions

At the end of ureteroscopy, double-j stent was inserted as normal in control group and reverse in study group. Follow-up duration was until removed stent from kidney about a month. Randomisation was performed online on website:<https://clincalc.com/stats/samplesize.aspx>

Intervention Type

Procedure/Surgery

Primary outcome measure

Ureteral stent symptoms questionnaire score before stent removal

Secondary outcome measures

Pain measured as use of painkiller or not, measured by interview before stent removal

Overall study start date

01/05/2023

Completion date

01/09/2024

Eligibility**Key inclusion criteria**

Patient underwent ureteroscopy and inserted double-j stent

Participant type(s)

Patient

Age group

Adult

Lower age limit

18 Years

Upper age limit

65 Years

Sex

Both

Target number of participants

140

Total final enrolment

116

Key exclusion criteria

1. Bilateral operation performed
2. Have previous lower urinary tract symptoms (etc. prostate hyperplasia, overactive bladder)
3. Pregnant

Date of first enrolment

05/06/2024

Date of final enrolment

05/08/2024

Locations

Countries of recruitment

Türkiye

Study participating centre

Gebze Medical Park Hospital

Guzeller mah

Kocaeli

Türkiye

41400

Sponsor information

Organisation

Gebze Medical Park Hospital

Sponsor details

Guzeller mah kavak cad

Kocaeli

Türkiye
41400
+90 (262) 675 75 75
erol.turan@medicalpark.com.tr

Sponsor type

Hospital/treatment centre

Website

<https://www.medicalpark.com.tr/gebze-hastanesi/h-7>

Funder(s)

Funder type

Other

Funder Name

Investigator initiated and funded

Results and Publications

Publication and dissemination plan

Planned publication in a peer-reviewed journal

Intention to publish date

01/11/2024

Individual participant data (IPD) sharing plan

Not provided at time of registration

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		24/02/2025	25/02/2025	Yes	No