

Intervention for mothers during pregnancy to reduce exposure to second-hand smoke

| | | |
|--|---|---|
| Submission date 11/02/2024 | Recruitment status No longer recruiting | <input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol |
| Registration date 19/02/2024 | Overall study status Completed | <input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results |
| Last Edited 18/03/2025 | Condition category Other | <input type="checkbox"/> Individual participant data |

Plain English summary of protocol

Background and study aims

This study looks at a special program designed to help pregnant women in India and Bangladesh avoid second-hand smoke at home. Second-hand smoke can be harmful, especially to unborn babies, leading to health problems after birth. The researchers wanted to see if this program, which includes various supportive materials and advice, could make a difference in reducing smoke exposure.

Who can participate?

Pregnant women who don't use tobacco themselves but are around second-hand smoke, confirmed by a saliva test, were invited to join the study. They had to be within the first 20 weeks of their pregnancy to participate.

What does the study involve?

Participants were divided into two groups. One group received a comprehensive package that included educational booklets, personalized messages, feedback on their exposure levels, and one-on-one consultations. The other group got just an educational leaflet. The main goal was to see if this approach could lower the levels of smoke exposure, as measured by a saliva test, three months later. The study also looked at how much the women knew about the risks of second-hand smoke, their confidence in asking family members to smoke less, and how ready they thought their families were to make these changes.

What are the possible benefits and risks of participating?

The big hope was that this program would lead to healthier environments for pregnant women, reducing the risks associated with second-hand smoke. There weren't any significant risks involved in participating, as the study mainly focused on providing information and support.

Where is the study run from?

The research was carried out in two locations: Comilla in Bangladesh and Bangalore in India. These sites were chosen to recruit and study the women involved.

When is the study starting and how long is it expected to run for?

December 2015 to June 2017

Who is funding the study?

1. Medical Research Council (MRC) (UK)
2. Department of Biotechnology (DBT) (India)

It's important to note that these funders didn't influence how the study was conducted or reported.

Who is the main contact?

Atif Rahman, atif.rahman@liverpool.ac.uk

Contact information

Type(s)

Public, Scientific, Principal Investigator

Contact name

Prof Atif Eahman

Contact details

Waterhouse Building Block B
Liverpool
United Kingdom
L69 3BX
+44 (0)151 794 2000
atif.rahman@liverpool.ac.uk

Type(s)

Scientific

Contact name

Dr Veena Satyanarayana

ORCID ID

<http://orcid.org/0000-0002-9608-2228>

Contact details

Department of Clinical Psychology
National Institute of Mental Health and Neuro Sciences (NIMHANS)
Bangalore
India
560029
+91 (0)80 26995180
veenas@nimhans.ac.in

Additional identifiers

EudraCT/CTIS number

Nil known

IRAS number

ClinicalTrials.gov number

Nil known

Secondary identifying numbers

Protocol v1

Study information

Scientific Title

Intervention for Mothers during Pregnancy to Reduce Exposure to Second-hand Smoke (IMPRESS): a pilot randomized controlled trial in Bangladesh and India

Acronym

IMPRESS

Study objectives

The aims were to examine the feasibility and acceptability of

1. The study design, measures and methods including delivery of a multicomponent intervention to reduce home exposure to second-hand smoke (SHS) among non-smoking pregnant women
2. The intervention in reducing home exposure to SHS among non-smoking pregnant women from two low- and middle-income countries (LMICs), India and Bangladesh
3. To estimate the standard deviation of the proposed primary outcome measure, to inform sample size calculation for a prospective definitive trial

Ethics approval required

Ethics approval required

Ethics approval(s)

1. Approved 01/03/2016, University of Liverpool (Foundation Building, Brownlow Hill, Liverpool, L69 7ZX, United Kingdom; +44 (0)151 794 2000; iphresearch@liverpool.ac.uk), ref: N/A
2. Approved 16/12/2015, National Institute of Mental Health and Neuro Sciences (NIMHANS) (Hosur Road / Marigowda Road, (Lakkasandra, Wilson Garden), Bangalore, 560029, India; +91 (0) 80 26995000; dirstaff@nimhans.ac.in), ref: N/A

Study design

Pilot randomized controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Community, GP practice

Study type(s)

Prevention, Quality of life

Participant information sheet

Not available in web format, please use contact details to request a participant information sheet

Health condition(s) or problem(s) studied

Second-hand smoke

Interventions

Multicomponent behavioural intervention

Participants were divided into two groups. Simple 1:1 randomization using computer-generated sequence of random numbers was used. One group received a comprehensive package that included educational booklets, personalized messages, feedback on their exposure levels, and one-on-one consultations. The other group received an educational leaflet. The duration of the intervention was 2 months in the Bangalore site as this had an additional component of voice calls. The duration of follow-up was 3 months.

Intervention Type

Behavioural

Primary outcome measure

Saliva cotinine levels measured using NicAlert® test at baseline and 3-month follow-up

Secondary outcome measures

Women's knowledge of SHS; attitudes to SHS and behaviours to reduce SHS exposure; perceived confidence in negotiating change with husbands and other family members and perceived readiness of their husband to change, measured using researcher-administered knowledge, attitude and behaviour questionnaire at baseline and 3 months

Overall study start date

01/12/2015

Completion date

01/06/2017

Eligibility**Key inclusion criteria**

1. Women within 20 weeks of pregnancy attending the selected clinics
2. Over 18 years old, reported SHS exposure by their husbands at home
3. Resident in the area and were NicAlert positive

Participant type(s)

Healthy volunteer

Age group

Adult

Lower age limit

18 Years

Upper age limit

50 Years

Sex

Female

Target number of participants

100

Total final enrolment

101

Key exclusion criteria

Women using tobacco (smoking or chewing tobacco)

Date of first enrolment

01/11/2016

Date of final enrolment

01/06/2017

Locations**Countries of recruitment**

Bangladesh

India

Study participating centre

National Institute of Mental Health and Neuro Sciences

NIMHANS

Bangalore

India

560029

Study participating centre

Ark Foundation

Suite C3-4, House, 6 Rd 109

Dhaka

Bangladesh

1212

Sponsor information

Organisation

University of Liverpool

Sponsor details

Waterhouse Building, Block B
Liverpool
England
United Kingdom
L69 3BX
+44 (0)151 794 2000
iphresearch@liverpool.ac.uk

Sponsor type

University/education

Website

<http://www.liv.ac.uk/>

ROR

<https://ror.org/04xs57h96>

Funder(s)**Funder type**

Research council

Funder Name

Medical Research Council

Alternative Name(s)

Medical Research Council (United Kingdom), UK Medical Research Council, MRC

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Funder Name

DBT India

Results and Publications

Publication and dissemination plan

The publication will be published in peer-reviewed journals.

Intention to publish date

01/06/2024

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be available upon request from Dr Veena A. Satyanarayana (veenas@nimhans.ac.in). The dataset will be shared comprising baseline and follow-up data in Excel format. Written informed consent was sought from participants consenting to participate in the trial. Datasets were pseudo-anonymized before analysis. No individual-level data will be shared in the publication.

IPD sharing plan summary

Available on request

Study outputs

| Output type | Details | Date created | Date added | Peer reviewed? | Patient-facing? |
|---------------------------------|---------|--------------|------------|----------------|-----------------|
| Results article | | 17/04/2024 | 18/03/2025 | Yes | No |