

Tracking physical activity after total knee replacement

Submission date 08/12/2020	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 04/02/2021	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
Last Edited 20/06/2024	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Total knee arthroplasty surgery, also known as total knee replacement, is recognised as a successful procedure. However some patients find recovery more difficult than others. At present there is little evidence on the best type of exercise recovery plan to give patients following surgery. We think it likely that recovery strategies can be targeted more effectively if they take into consideration participant information that can be collected before surgery, such as expectations regarding recovery, pain, and psychological status.

Who can participate?

Adults over 18 years, scheduled for total knee arthroplasty for osteoarthritis.

What does the study involve?

In this study we ask participants undergoing surgery to wear an activity monitor, worn on the wrist like a watch for a week in the month preceding surgery, and then for six weeks following surgery. We will also collect data prior to surgery on:

- ability to do everyday tasks (using a validated questionnaire);
- functional tests (40m walking test, climbing 9 steps, and number of times participant can do sit-to-stand in 30s);
- pain levels measured on visual analogue scale;
- analgesic use;
- inflammation using level of C-reactive protein in blood (measured again day following surgery)
- pre-operative characteristics –socioeconomic data; expectations of surgery and pain; mental health status; frailty level; number of falls in last 3 months
- osteoarthritis classification (measured on routine x-rays)

Participants will be called once a week following surgery to record their pain level and analgesic use, and to check they are happy wearing the activity monitor. At six weeks, data will again be collected on the ability to do everyday tasks; functional tests; analgesic and resource use and pain.

In the analysis, changes in the primary outcome measure of activity over time, and distinct subtypes amongst TKA recovery trajectories will be examined using statistical modelling.

What are the possible benefits and risks of participating?

We cannot promise the study will help you but we hope that the information we obtain from this study will help improve treatment for patients having total knee replacement surgery in the future. The possible risks and/or discomforts associated with the knee replacement operation are as for a routine knee replacement operation. There is no increased surgical risk if you take part in this study. You will have study visits before surgery and at 6 weeks which should take place on the same day as your routine visits, and we will give you a phone call every week up to the 6 weeks following surgery, to monitor your pain levels and your use of painkillers.

Where is the study run from?

This study is a collaboration between Northumbria Healthcare NHS Foundation Trust and the Orthopaedic Research Institute (ORI) at Bournemouth University (UK).

When is the study starting and how long is it expected to run for?

January 2020 to September 2022

Who is funding the study?

Zimmer, Inc. (USA)

Who is the main contact?

Tom Wainwright (scientific), twainwright@bournemouth.ac.uk

Tikki Immins (public), timmins@bournemouth.ac.uk

Contact information

Type(s)

Scientific

Contact name

Mr Tom Wainwright

ORCID ID

<https://orcid.org/0000-0001-7860-2990>

Contact details

Orthopaedic Research Institute

Bournemouth University

89 Holdenhurst Road

Bournemouth

United Kingdom

BH8 8EB

+44 (0)1202 961656

twainwright@bournemouth.ac.uk

Type(s)

Public

Contact name

Mrs Tikki Immins

ORCID ID

<https://orcid.org/0000-0002-9797-6098>

Contact details

Orthopaedic Research Institute
Bournemouth University
89 Holdenhurst Road
Bournemouth
United Kingdom
BH8 8EB
+44 (0)1202 961656
timmins@bournemouth.ac.uk

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Integrated Research Application System (IRAS)

274998

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

CPMS 44153, IRAS 274998

Study information

Scientific Title

Tracking physical activity after total knee arthroplasty

Study objectives

This study aims to identify and differentiate high and low physical activity responders in the early recovery stage following total knee arthroplasty (TKA). This will enable further studies of interventions designed for groups of patients at risk of a poorer recovery of physical function, unlike earlier studies undertaken with “average patients” which have had a limited effect on postoperative recovery.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 02/03/2020, North of Scotland Research Ethics Committee (Summerfield House, 2 Eday Road, Aberdeen, AB15 6RE, UK; +44 (0)1224558458; nosres@nhs.net), ref: 20/NS/0035

Study design

Observational cohort study

Primary study design

Observational

Study type(s)

Other

Health condition(s) or problem(s) studied

Physical activity after total knee arthroplasty

Interventions

This is a prospective cohort study recruiting 120 participants from patients undergoing total knee arthroplasty surgery as part of their routine care at Northumbria Healthcare NHS Foundation Trust.

Participants will be assessed at baseline up to one month prior to surgery, and be asked to wear an activity monitor for 7 consecutive days prior to surgery. The activity monitor is an Actiwatch (CE marked. Phillips Respironics Murrysville, PA, USA). After routine surgery, at discharge, participants will be asked to again wear the activity monitor continuously for the six weeks following surgery, when they will be assessed again. They will be rung once each week up to 6 weeks to check on pain levels on activity, and on analgesic use.

Screening and consent

The clinical team will identify patients who are suitable to take part in the study. During a routine appointment a member of the clinical care team will explain the purpose of the study and invite potential participants to discuss the study with a member of the research team. The research team member will explain the purpose of the study, the procedures and the duration to the patient. If they would like to proceed they will be asked to complete an informed consent form. The research team member will ensure that the patient has sufficient time to consider taking part in the study before taking consent, in line with GCP guidelines.

Visit 1 – baseline assessment, up to one month before surgery

Participants will be given an activity monitor to wear for seven consecutive days before surgery. Data on demographics, medical history, physical examination (to include height and weight), current medication, name of GP, American Society of Anaesthesiologists classification will be recorded for the study. X-rays taken as part of routine care will be assessed for level of osteoarthritis using the Kellgren and Lawrence classification; and blood taken prior to surgery as part of pre-assessment testing, will be tested for C-reactive protein level (CRP). Participants will be asked to rate their pain on activity, and undertake functional tests – how many sit-to-stands they can perform in 30s, walk fast-paced for 40m, and a stair climb test. They will also be asked to complete questionnaires on how they find their pain and ability to do everyday activities, their expectations of surgery, their frailty level, how they perceive their pain will be following surgery, levels of anxiety and depression, use of analgesics, and number of falls in the past three months. Visit 2 – the day following surgery, the CRP protein level will be measured again, along with pain on activity level, and changes in analgesic use and other medications. At discharge, participants will be asked to wear their activity monitor again until they return for their 6 weeks follow-up visit. They will be asked to wear it continuously, except if they have hydrotherapy or go swimming.

Visit 3 – from discharge to 6 weeks, participants will be rung each week to check they are happy with the activity monitor, and to check their pain on activity level, changes in analgesic use and other medications. Visit 4 – 6 weeks following surgery. Participants will return their activity monitor. Their length of hospital stay will be recorded, and the reasons if the stay is over 4 days. Changes in analgesic use and pain level on activity will be recorded. Participants will be asked to repeat the functional tests; and to complete again a questionnaire on they find their pain and ability to do everyday activities.

Intervention Type

Behavioural

Primary outcome(s)

Physical activity, as measured using an Actiwatch device (Philips Respironics, Murrysville, PA, USA) from baseline (pre-surgery) to 8 weeks post-surgery

Key secondary outcome(s)

At baseline and 8-weeks post-surgery (unless otherwise stated):

1. Physical performance (PBOMs): 40m paced walk test, 30s sit-to-stand, 9 step stair-climb
2. Pain and function daily living subscales from Knee Injury and Osteoarthritis Outcome Score (KOOS)
3. Analgesic use measured using self report
4. Pain levels on activity measured using a numerical rating scale at baseline, and each week post-surgery until 8 weeks
5. C-reactive protein (CRP) measured using ... from baseline to one day following surgery (or at discharge if discharged on day of surgery)
6. Patient expectations (HSS Knee Replacement Expectation Survey) at baseline
7. Anxiety and depression measured using the Hospital Anxiety and Depression Score (HADS) at baseline
8. Pain catastrophizing measured using the Pain Catastrophizing Scale (PCS) at baseline
9. Frailty measured using the Edmonton Frail Scale at baseline
10. Number of falls in the past three months measured using self report at baseline
11. Classification of osteoarthritis measured using the Kellgren and Lawrence system at baseline

Completion date

30/09/2022

Eligibility

Key inclusion criteria

1. Participants capable of giving informed consent
2. Scheduled for primary unilateral total knee arthroplasty for osteoarthritis
3. Aged 18 years and over

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

120

Key exclusion criteria

Unable to complete follow-up (insufficient English, lives overseas, unable to return easily).

Date of first enrolment

12/04/2021

Date of final enrolment

30/04/2022

Locations**Countries of recruitment**

United Kingdom

England

Study participating centre**North Tyneside General Hospital City**

Northumbria Healthcare NHS Foundation Trust

Rake Lane

North Shields

United Kingdom

NE29 8NH

Sponsor information**Organisation**

Bournemouth University

ROR

<https://ror.org/05wwcw481>

Funder(s)**Funder type**

Industry

Funder Name

Zimmer; Grant Codes: IRE2020-06CH

Alternative Name(s)

Zimmer, Inc., Zimmer Biomet

Funding Body Type

Private sector organisation

Funding Body Subtype

For-profit companies (industry)

Location

United States of America

Results and Publications

Individual participant data (IPD) sharing plan

The current data sharing plans for this study are unknown and will be available at a later date.

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
HRA research summary			26/07/2023	No	No