Impact of a novel molecular tuberculosis (TB) diagnostic system in patients at high risk of TB mortality in rural South Africa

Submission date	Recruitment status No longer recruiting	[X] Prospectively registered		
15/06/2011		[X] Protocol		
Registration date	Overall study status	Statistical analysis plan		
17/06/2011	Completed	[X] Results		
Last Edited	Condition category	Individual participant data		
25/07/2017	Infections and Infestations			

Plain English summary of protocol

Background and study aims

HIV and tuberculosis (TB) infections are the leading causes of death for young adults in South Africa. TB bacteria that have developed resistance to first-line TB drugs (multidrug-resistant TB or MDR-TB) are now causing significant numbers of deaths. Diagnosis of MDR-TB is a laborious process which can take two to three months, by which time up to half of those with MDR-TB will have died. The Xpert MTB/RIF system is a new diagnostic test which can identify TB and can also recognise whether or not there is drug resistance within two hours. Hlabisa sub-district in KwaZulu-Natal is one place where MDR-TB has become very common in association with high levels of HIV infection. This study aims to investigate the impact of the Xpert MTB/RIF system, specifically whether outcomes are different when the system is positioned at different levels of the health system.

Who can participate?

Patients aged 18 or older with suspected TB, who have a confirmed HIV infection and/or are at high risk for MDR-TB

What does the study involve?

Participants are randomly allocated to one of two strategies: Xpert MTB/RIF testing at a district hospital laboratory or Xpert MTB/RIF testing at a primary health care clinic (point-of-care). The number of TB patients starting appropriate treatment within 30 days, the time taken to start TB treatment, and the number of deaths and hospital admissions in first 60 days are all measured.

What are the possible benefits and risks of participating? Not provided at time of registration

Where is the study run from?

Africa Centre for Health and Population Studies (South Africa)

When is the study starting and how long is it expected to run for? July 2011 to December 2012

Who is funding the study? Wellcome Trust (UK)

Who is the main contact? Dr Richard Lessells rlessells@africacentre.ac.za

Contact information

Type(s)

Scientific

Contact name

Dr Richard Lessells

Contact details

Africa Centre for Health and Population Studies PO Box 198 Mtubatuba South Africa 3935 +27 (0)35 550 7500 rlessells@africacentre.ac.za

Additional identifiers

Protocol serial number

Wellcome Trust grant number 090999/Z/09/Z, South African National Clinical Trials Registry DOH-27-0711-3568

Study information

Scientific Title

Impact of a novel molecular tuberculosis (TB) diagnostic system in patients at high risk of TB mortality in rural South Africa: a pragmatic cluster randomised controlled trial

Study objectives

Timely initiation of appropriate tuberculosis (TB) treatment will be improved when the diagnostic system is positioned at the primary health care clinic (point-of-care) compared to when it is positioned centrally at the district hospital laboratory.

Ethics approval required

Old ethics approval format

Ethics approval(s)

- 1. University of KwaZulu-Natal Biomedical Research Ethics Committee, ref: BF033/11
- 2. London School of Hygiene and Tropical Medicine Ethics Committee, ref: 5926 Approval pending as of 15/06/2011

Study design

Single-centre cluster randomised controlled trial

Primary study design

Interventional

Study type(s)

Diagnostic

Health condition(s) or problem(s) studied

Pulmonary tuberculosis

Interventions

Comparison between two delivery strategies for the GeneXpert system and Xpert MTB/RIF test: Positioning at district hospital vs positioning at primary health care clinic

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Proportion of TB cases initiated on appropriate TB treatment within 30 days of enrolment

Key secondary outcome(s))

- 1. Time to initiation of appropriate TB treatment
- 2. Time to initiation of appropriate MDR-TB treatment
- 3. All-cause mortality at 60 days
- 4. Hospital admissions in first 60 days
- 5. Time to initiation of antiretroviral therapy (for eligible HIV-infected participants)

Completion date

31/12/2012

Eligibility

Key inclusion criteria

- 1. TB suspect (defined as cough of any duration)
- 2. Age 18 years or older
- 3. Confirmed human immunodeficiency virus (HIV) infection and/or high risk for multi-drug-resistant tuberculosis (MDR-TB)

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

- 1. Severely unwell requiring admission to hospital
- 2. Previous MDR/extensively drug-resistant tuberculosis (XDR-TB) diagnosis or treatment
- 3. Diagnosis or suspicion of extra-pulmonary TB only
- 4. Unable to give informed consent

Date of first enrolment

04/07/2011

Date of final enrolment

31/12/2012

Locations

Countries of recruitment

South Africa

Study participating centre Africa Centre for Health and Population StudiesMtubatuba

South Africa 3935

Sponsor information

Organisation

London School of Hygiene and Tropical Medicine (UK)

ROR

https://ror.org/00a0jsq62

Funder(s)

Funder type

Charity

Funder Name

Wellcome Trust (UK) (090999/Z/09/Z)

Alternative Name(s)

Funding Body Type

Private sector organisation

Funding Body Subtype

International organizations

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created Date added	Peer reviewed?	Patient-facing?
Results article	results	01/10/2017	Yes	No
Protocol article	protocol	12/06/2013	Yes	No
Participant information sheet	Participant information sheet	11/11/2025 11/11/2025	No	Yes