

Brain imaging responses to food in insulin resistance - screening

Submission date 14/12/2010	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 14/12/2010	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 29/03/2018	Condition category Nutritional, Metabolic, Endocrine	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Obesity (being very overweight) and health problems related to obesity (including type 2 diabetes) are becoming more common, causing long-term ill health. As yet we do not understand why some people are particularly prone to weight gain and diabetes. One possibility is that people who are more prone to obesity and diabetes have a malfunction in the brain mechanisms that stop their desire to eat more after a meal. Gaining further knowledge of the way the brain controls eating will help the development of new ways to prevent and treat these diseases. The researchers are running a study to look at the way the brain controls appetite by using functional magnetic resonance imaging (fMRI), comparing the results from people who are "insulin resistant" and therefore at a higher risk of developing diabetes with people who are "insulin sensitive" and therefore at a lower risk of developing diabetes. Here, they are screening volunteers to see if they are able to join the study.

Who can participate?

Men aged between 18-65 years with a body mass index (BMI) of no more than 30 kg/m². Insulin sensitive participants should not have any family history of diabetes mellitus. Insulin resistant subjects must have first degree relatives (i.e. parent, sibling or child) with type 2 diabetes.

What does the study involve?

All participants are asked to complete a number of questionnaires about their eating habits to check that they don't have an eating disorder. They are also asked to provide their medical history and have a brief physical examination (which includes taking their height, weight, neck and waist measurements). Blood samples are taken after an overnight fast to check that they don't have diabetes and to see how well they respond to insulin. The results of this screening will be used to check if each participant can take part in the next part of the study (see <http://www.isrctn.com/ISRCTN51099878>)

What are the possible benefits and risks of participating?

Not provided at time of registration

Where is the study run from?

King's College Hospital NHS Trust

When is the study starting and how long is it expected to run for?
December 2010 to November 2013

Who is funding the study?
Diabetes UK

Who is the main contact?
Dr Yee Seun Cheah
y.cheah@nhs.net

Contact information

Type(s)
Scientific

Contact name
Dr Yee Seun Cheah

Contact details
Kings College London
10 Cutcombe Road
London
United Kingdom
SE5 9RJ
+44 (0) 20 7848 5654
y.cheah@nhs.net

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers
9515

Study information

Scientific Title
Brain imaging responses to food in insulin resistance: a single centre non-randomised interventional screening trial

Acronym
DRN 546

Study objectives

Obesity and related health problems including type 2 diabetes are becoming more common, causing long-term ill health. As yet, it is not understood why some people are particularly prone to weight gain and diabetes. One possibility is a malfunction in the brain mechanisms that stop our desire to eat more after a meal in people predisposed to obesity and diabetes. Gaining further knowledge of the way the brain controls eating will help the development of new ways to prevent and treat these diseases.

The project will look at the way the brain controls appetite by using functional magnetic resonance imaging (fMRI). This is a method of taking images of the brain that will allow us to see the activity of brain regions that control eating. Brain responses will be studied after eating in healthy relatives of people with diabetes, who are "insulin resistant", where the body is less responsive to insulin, a hormone normally produced by the body to control sugar (glucose) levels. These people will therefore be at higher risk of developing diabetes and obesity. They will be compared to people who are insulin sensitive, at lower risk of diabetes. The impact of treating insulin resistance on these brain responses will then be investigated. This will allow researchers to see if the brain controls eating differently in those at risk of diabetes and obesity, and whether it can be reversed. The imaging methods that are developed may also permit the early assessment of potential therapies to improve appetite control, aiding the development of new ways to prevent or treat obesity and diabetes in the future.

This screening study will identify appropriate subjects that will enter the subsequent intervention study (<http://www.isrctn.com/ISRCTN51099878>) described above.

Ethics approval required

Old ethics approval format

Ethics approval(s)

South East London REC3 (formally King's College Hospital REC), 18/06/2010, ref:10/H0808/47a

Study design

Single centre non-randomised interventional screening trial

Primary study design

Interventional

Secondary study design

Non randomised study

Study setting(s)

Hospital

Study type(s)

Screening

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Topic: Diabetes Research Network; Subtopic: Type 2; Disease: Obesity

Interventions

Volunteers will attend a screening visit to determine their eligibility to enter the intervention study (UKCRN 9117; DRN 518; ISRCTN51099878) according to the inclusion criteria. This will include completing validated questionnaires of eating behaviour to exclude the presence of eating disorders. Volunteers will also provide their medical history and undergo a brief physical examination, which will include measurements of their height, weight, neck and waist circumference. Blood samples will be collected after an overnight fast to exclude the presence of diabetes and to establish the degree of insulin sensitivity. The screening process will involve a maximum of 2 visits within a one month period.

Follow-up length: 1 months

Study entry: registration only

Intervention Type

Other

Phase

Not Applicable

Primary outcome measure

Insulin sensitivity and glycaemic status, results of intervention to be available within a maximum of one month of completing screening process

Secondary outcome measures

Eating behaviour, determined during screening visit

Overall study start date

01/12/2010

Completion date

01/11/2013

Eligibility

Key inclusion criteria

All subjects (insulin sensitive and insulin resistant):

1. Men
2. Age 18 - 65 years (inclusive at time of recruitment)
3. Right-handed
4. English speaking
5. No active medical illness including diabetes mellitus
6. Body mass index (BMI) less than or equal to 30 kg/m²

Insulin sensitive subjects:

7. No family history of diabetes mellitus
8. Insulin sensitive (determined by homeostatic model assessment - insulin resistance [HOMA2-IR] during screening process)

Insulin resistant subjects:

9. First degree relatives of patients with type 2 diabetes mellitus
10. Insulin resistance (determined by HOMA2-IR during screening process)

Participant type(s)

Patient

Age group

Adult

Lower age limit

18 Years

Sex

Male

Target number of participants

Planned sample size: 100; UK sample size: 100

Key exclusion criteria

1. Women
2. Left-handedness
3. Current or past history of significant substance abuse or eating disorder
4. Neurological or psychiatric disorders
5. Use of medication that may affect brain activity (e.g. antidepressants, anticonvulsants, antipsychotic drugs), drugs for obesity (orlistat or sibutramine) or drugs that lower glucose (e.g. metformin, sulphonylureas, thiazolidinediones, incretins or insulin)
6. Inability to understand spoken and/or written English
7. Claustrophobia (because of the small bore of the MR scanner)
8. BMI greater than 30 kg/m²
9. Contraindication to MRI (pacemaker in situ, extensive dental work, history of penetrating eye trauma, presence of metal surgical clips etc.)
10. Presence of diabetes

Date of first enrolment

01/12/2010

Date of final enrolment

01/11/2013

Locations

Countries of recruitment

England

United Kingdom

Study participating centre

King's College London
10 Cutcombe Road
London
United Kingdom
SE5 9RJ

Sponsor information

Organisation

Kings College London (KCL)

Sponsor details

c/o Mr Keith Brennan
Hodgkin Building
New Hunts House
Guy's Campus
London
England
United Kingdom
SE1 1UL

Sponsor type

University/education

Website

<http://www.kcl.ac.uk/index.aspx>

Organisation

King's College Hospital NHS Foundation Trust

Sponsor details

Denmark Hill
London
England
United Kingdom
SE5 9RS

Sponsor type

Hospital/treatment centre

Website

<http://www.kch.nhs.uk/home/>

Organisation

King's College London

Sponsor details**Sponsor type**

Not defined

Website

<http://www.kcl.ac.uk/index.aspx>

ROR

<https://ror.org/0220mzb33>

Funder(s)**Funder type**

Charity

Funder Name

Diabetes UK

Alternative Name(s)

DIABETES UK LIMITED, British Diabetic Association

Funding Body Type

Private sector organisation

Funding Body Subtype

Trusts, charities, foundations (both public and private)

Location

United Kingdom

Results and Publications**Publication and dissemination plan**

29/03/2018: results presented at European Association for the Study of Diabetes Annual meeting 2013 (<https://www.easd.org/virtualmeeting/home.html#!resources/increasing-homa-ir-modulates-brain-responses-to-meal-ingestion-in-insulin-sensitive-men-a-continuous-arterial-spin-labelling-functional-magnetic-resonance-imaging-study>)

Intention to publish date**Individual participant data (IPD) sharing plan**

IPD sharing plan summary

Not provided at time of registration