A multicentre randomised phase II study of hypofractionated bladder radiotherapy with or without image guided adaptive planning

Submission date	Recruitment status No longer recruiting	[X] Prospectively registered		
11/09/2013		[X] Protocol		
Registration date	Overall study status	Statistical analysis plan		
11/09/2013	Completed	[X] Results		
Last Edited	Condition category	[] Individual participant data		
27/11/2025	Cancer			

Plain English summary of protocol

http://www.cancerresearchuk.org/cancer-help/trials/a-trial-to-learn-more-about-weekly-radiotherapy-for-invasive-bladder-cancer-and-to-look-differnt-way-planning-treatment-hybrid

Contact information

Type(s)

Scientific

Contact name

Dr HYBRID Trial

Contact details

Section of Clinical Trials 15 Cotswold Road Sutton United Kingdom SM2 5NG +44 20 7352 8133 HYBRID-icrctsu@icr.ac.uk

Additional identifiers

ClinicalTrials.gov (NCT)

NCT01810757

Protocol serial number

15216

Study information

Scientific Title

A multicentre randomised phase II study of HYpofractionated Bladder Radiotherapy with or without Image guided aDaptive planning

Acronym

HYBRID

Study objectives

HYBRID aims to collect robust information about the side effects of weekly bladder radiotherapy, and is also investigating whether side effects can be reduced by using scans taken before each weekly treatment to tailor radiotherapy delivery (adaptive radiotherapy).

Ethics approval required

Old ethics approval format

Ethics approval(s)

11/09/2013, ref: 13/LO/1350

Study design

Randomised; Interventional; Design type: Treatment

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Topic: National Cancer Research Network; Subtopic: Bladder Cancer; Disease: Bladder (superficial)

Interventions

Participants will be randomly allocated between weekly radiotherapy and weekly adaptive radiotherapy. Both groups will receive the same radiotherapy dose.

- 1. Adaptive planning, 36 Gray of hypofractionated radiotherapy will be delivered in 6 fractions using the best fitting of 3 plans at each treatment. Plans will be selected and verified by accredited staff.
- 2. Standard planning (control), 36 Gray of hypofractionated radiotherapy will be delivered in 6 fractions using standard planning and delivery techniques.

Participants will visit the hospital 4 weeks, 3, 6, 12 and 24 months after radiotherapy to receive treatment for any symptoms and to check whether the cancer has returned. The main aim of HYBRID is to establish whether use of adaptive radiotherapy can reduce nonbladder side effects by reducing the amount of nonbladder tissue exposed to high doses of radiotherapy. HYBRID will also combine the results of both treatment groups to investigate how well weekly radiotherapy controls bladder cancer.

Added 27/11/2025:

Additional Data Linkage Information:

Participants from this trial will also be included in the INTERACT project which will link to their data held by NHS England. For more information, please see the INTERACT website: https://www.icr.ac.uk/interact.

Intervention Type

Other

Primary outcome(s)

Proportion of patients experiencing severe acute non-genitourinary side effects after radiotherapy; Timepoint(s): Up to 3 months

Key secondary outcome(s))

Not provided at time of registration

Completion date

31/12/2020

Eligibility

Key inclusion criteria

- 1. Written informed consent
- 2. Age =18 years
- 3. Histologically confirmed invasive bladder carcinoma (T2T4a N0 M0; any histological subtype)
- 4. Unsuitable for radical cystectomy or daily fractionated radiotherapy for any reason (including performance status, comorbidity, patient refusal)
- 5. Expected survival more than 6 months
- 6. WHO performance status 03
- 7. Willing to undergo post treatment cystoscopy
- 8. Able to attend post treatment follow up

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Mixed

Lower age limit

18 years

Upper age limit

100 years

Sex

All

Total final enrolment

65

Key exclusion criteria

- 1. Nodal or metastatic disease
- 2. Concurrent malignancy within 2 years of randomisation (not including non melanomatous skin carcinoma, previous non muscle invasive bladder tumours, NCCN low risk prostate cancer (T1 /T2a, Gleason 6 PSA <10), in situ carcinoma of any site)
- 3. Previous pelvic radiotherapy
- 4. Urinary catheter in situ
- 5. Any other contraindication to radiotherapy (e.g. inflammatory bowel disease)

Date of first enrolment

15/04/2015

Date of final enrolment

31/08/2016

Locations

Countries of recruitment

United Kingdom

England

Study participating centre Section of Clinical Trials

15 Cotswold Road Sutton England SM2 5NG

Sponsor information

Organisation

Institute of Cancer Research (UK)

ROR

https://ror.org/043jzw605

Funder(s)

Funder type

Charity

Funder Name

Cancer Research UK (UK)

Alternative Name(s)

CR_UK, Cancer Research UK - London, Cancer Research UK (CRUK), CRUK

Funding Body Type

Private sector organisation

Funding Body Subtype

Other non-profit organizations

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	11/12/2020	05/07/2023	Yes	No
Protocol article		26/05/2020	20/06/2022	Yes	No
HRA research summary			28/06/2023	No	No
Plain English results			12/02/2021	No	Yes