

Effects of ultraviolet A1 (UVA1) phototherapy in patients with systemic sclerosis (SSc)

Submission date 07/02/2008	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 08/02/2008	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
Last Edited 29/06/2016	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Prof Chak Lau

Contact details
University Division of Medicine & Therapeutics
Ninewells Hospital and Medical School
Dundee
United Kingdom
DD1 9SY

Additional identifiers

Protocol serial number
N/A

Study information

Scientific Title
Effects of high dose UVA1 phototherapy on cutaneous fibrosis and endothelial cell function in patients with systemic sclerosis

Study objectives

One of the characteristic features of systemic sclerosis (SSc) is excessive deposition of collagen within the skin. Such changes are believed to be the results of inappropriate activation of dermal fibroblasts by various inflammatory and pro-fibrotic cytokines, combined with damage to the endothelium. Numerous treatments, some such as immunosuppressive drugs with potentially hazardous side-effects, are currently used with only limited success. Recent pilot studies have reported successful treatment of patients with scleroderma (hard skin) by high dose ultraviolet A1 (UVA1) phototherapy. The hypothesis of this study is high dose UVA1 phototherapy is useful in the treatment of skin fibrosis in patients with SSc through the reduction of collagen deposition and improvement of endothelial functions.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Within patient, double blind, sham controlled study.

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Systemic sclerosis

Interventions

1. Intervention: UVA1 photo-irradiation therapy to the patient's arm
2. Control: sham treatment to the patient's arm

Patients with early stage SSc will receive a total of twenty phototherapy treatments. Both arms will be placed into similar light boxes but only one arm will receive the UVA1 light therapy treatment. Comparisons will be made by measuring elasticity and collagen content of the skin, but also by looking at skin biopsies, blood vessel function and release of various cytokines and expression of markers of scarring tissues in the skin biopsy.

The study will take a total of eight weeks per participant. There will be a total of twenty phototherapy sessions with the duration of treatment at each session being no longer than 50 minutes. This will depend upon the individual and the minimal erythema dose (MED) (the MED is defined as the minimum amount of irradiation at a waveband capable of producing a perceptible erythema) as to how long the phototherapy at each session will take.

Assessments will take place at the start of the study, during the treatment and again two weeks after the final treatment session. The following assessments will be performed:

Visit 1: day 0 - duration: no longer than 3 hours

1. Written consent
2. Medical history
3. Phototesting

4. Skin elasticity
5. Iontophoresis and scanning laser doppler imaging
5. Skin scan
6. Skin biopsy
7. Haematology and biochemistry blood tests
8. Photograph arms

Visit 2: day 6 - duration: no longer than 20 minutes

1. Skin elasticity
2. Skin scan
3. Visual analogue scale
4. 5-point likert scale

Visit 3: day 15 - duration: no longer than 2 hours

1. Skin elasticity
2. Iontophoresis and scanning laser doppler imaging
3. Skin scan
4. Skin biopsy (group 1)
5. Haematology and biochemistry blood tests
6. Visual analogue scale
7. 5-point likert scale

Visit 4: day 21 - duration: no longer than 20 minutes

1. Skin elasticity
2. Skin scan
3. Photograph arms
4. Visual analogue scale
5. 5-point likert scale

Visit 5: 14 days after final treatment - duration: no longer than 2 hours

1. Skin elasticity
2. Iontophoresis and scanning laser doppler imaging
3. Skin scan
4. Skin biopsy (group 2)
5. Haematology and biochemistry blood tests
6. Photograph arms
7. Visual analogue scale
8. 5-point likert scale

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Ultraviolet A1 (UVA1) phototherapy

Primary outcome(s)

To examine, within subjects, the effects of high dose UVA1 phototherapy in treating patients with SSc. Clinical and laboratory assessments of skin fibrosis, skin blood flow and endothelial cell function will be undertaken.

See interventions for details on when these points will be measured.

Key secondary outcome(s)

1. Measure the effect of UVA1 phototherapy on skin fibrosis using clinical assessment of the skin. The elasticity and tethering of the skin and one of the main components of the skin, collagen, will be analysed. The release of various noxious chemicals and expression of markers of scarring tissues will be examined in the skin biopsy.
2. Measure the effect of UVA1 phototherapy on cutaneous blood flow. The ability of the blood vessel to respond to local application of chemicals that are known to increase blood flow, a process called iontophoresis, will be taken as a measure of the function of the skin blood vessels.
3. Assess the affect of UVA1 phototherapy on the release of various noxious chemicals and expression of markers of scarring tissues in fibroblasts in the laboratory

See interventions for details on when these points will be measured.

Completion date

31/03/2011

Eligibility

Key inclusion criteria

1. Patients with SSc (both limited and diffuse disease pattern) as diagnosed according to the American College of Rheumatology preliminary classification criteria for this condition
2. SSc patients with cutaneous manifestations
3. SSc patients whose diagnosis was made within the previous 3 years
4. SSc patients who are capable of providing a written informed consent
5. Patients of either sex and aged 18 or above will be recruited

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

1. Patients who have received phototherapy in the previous three months
2. Patients who have been started on immunosuppressive treatment within the previous 6 months
3. Patients with localised or generalised morphea
4. Patients with other sclerodermas other than that associated with SSc. Examples of other forms of sclerodermas include occupational scleroderma (e.g. vinyl chloride disease), scleroporphyria (a generalised morphea picture arising due to porphyria cutanea tarda), acrosclerosis atrophicans (a late feature of some patterns of Lyme borreliosis, a tick-borne spirochaetal infection) and nephrogenic fibrosing dermopathy.
5. Pregnant or breastfeeding women
6. Subjects who are 18 years or under

Date of first enrolment

01/04/2008

Date of final enrolment

31/03/2011

Locations

Countries of recruitment

United Kingdom

Scotland

Study participating centre

University Division of Medicine & Therapeutics

Dundee

United Kingdom

DD1 9SY

Sponsor information

Organisation

University of Dundee (UK)

ROR

<https://ror.org/03h2bxq36>

Funder(s)

Funder type

University/education

Funder Name

University of Dundee (UK) - University Division of Medicine & Therapeutics Research Fund

Results and Publications

Individual participant data (IPD) sharing plan**IPD sharing plan summary**

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes